



Significance of PM_{2.5} Air Quality at the Indian Capital

Shovan Kumar Sahu, Sri Harsha Kota*

Department of Civil Engineering, Indian Institute of Technology Guwahati, Guwahati 781039, India

ABSTRACT

In New Delhi, the capital city of India, concentrations of regulated air pollutants often exceed the Indian national ambient air quality standards (INAAQS). As the sources of these pollutants differ, it is of utmost priority to understand the most dangerous air pollutant to formulate better control strategies in the city. In this study, regulated air pollutant concentrations in New Delhi during 2011 to 2014 were collected. Compared to other pollutants, PM_{2.5} concentrations exceeded the INAAQS quite often. While PM_{2.5} exceeded INAAQS during 85% of the days, NO₂, O₃, CO and SO₂ exceeded only on 37, 14, 11 and 0% of the days, respectively. Using air quality index approach, the most dominant pollutant was identified as PM_{2.5}, for 75 to 90% of the days. However, a seasonal variation in the percentage dominance of PM_{2.5} was observed. For example, PM_{2.5} was dominant during 95% of the winter and 68% of monsoon days. In addition to absolute concentrations, pollutants can also be ranked by studying their associated short term mortality impacts. However, such studies are rare in India. For the first time, the short term impact of PM_{2.5} concentrations on non-disease specific mortality in New Delhi was assessed using Poisson regression models. Results indicated that the excessive risk associated with PM_{2.5} estimated was 0.57, which was higher than the other regulated pollutants. This indicates a projected 6.2 and 6.5% decrease in mortality by meeting the PM_{2.5} Indian standards and WHO set limits, respectively.

Keywords: Air quality index; New Delhi; PM_{2.5}; Health impact assessment.

INTRODUCTION

A world health organization (WHO) report observes that air pollution resulted in around seven million deaths globally in 2012, of which South East Asian region, dominated by India, accounted for 2.3 million (WHO, 2014a). Outdoor particulate matter (PM) was the seventh highest killer in India during 1990–2010 (IHME, 2013). According to a recently released report (WHO, 2014a) by World Health Organization (WHO), among 1600 cities surveyed, thirteen of the Indian cities were among the top twenty worst polluted cities, with New Delhi leading the list. PM_{2.5} concentrations in New Delhi are atleast 10 times higher than Washington DC, and 3 times higher than Beijing (WHO, 2014a). In order to construe the status of air quality in India, National Ambient Air Quality Standards (INAAQS) were adopted in 1982 by the Central Pollution Control Board, with further revisions in 1994 and 2009. Predominant sources for these regulated pollutants differ. For example in New Delhi, residential, transport and industrial sectors are major sources for PM_{2.5} (Sahu *et al.*, 2011); transport sector is the major source of

CO and NO_x (Aneja *et al.*, 2001); and industries are major source for SO₂ (Sadavarte and Venkataraman, 2014). So this calls for knowledge about the most dangerous pollutant in the city to formulate stricter laws and better control strategies.

As the absolute concentrations of these pollutants differ, it is necessary to bring these pollutants onto a similar scale for direct comparison. To alleviate this, Ott (1978) suggested a scheme to transform the weighted values of air pollutant concentrations into a single or set of numbers, referred to as Air Quality Index (AQI), wherein bigger the AQI greater the pollution, higher is the health risk, and vice versa. The quality of air is reported as good, satisfactory, moderate, poor, very poor or severe depending on the overall AQI, calculated using individual AQI of pollutants considered. Several developed nations in the world, including the US, UK, Australia, and Canada, have their own AQI. These vary by the range of the index, and the methodology used to estimate species specific and overall AQI. For example, UK (COMEAP, 2011) and Canada (Chen and Copes, 2013) classify AQI into four categories, with AQI ranging from 0 to 10. However, the US uses six categories, ranging from 0 to 500. Moreover, while Canada uses a non-linear aggregate of AQI of different species based on their exposure-response relationships, the US and UK use maximum AQI of different species, to estimate the overall AQI. Few studies in India, tried to analyze air quality using AQI in cities.

*Corresponding author.

Tel.: +91-3612583331

E-mail address: harshakota@gmail.com

For example, Sharma *et al.* (2003) used Indian NAAQS and health standards by the US, for suspended particulate matter (SPM), CO, NO₂, O₃, SO₂, and PM₁₀ to estimate AQI in an Indian city. Bhaskar and Mehta (2010) used the US method to estimate AQI at thirteen different sites in Ahmadabad city in India, and observed strong seasonal effects on AQI. Banerjee and Srivastava (2011) estimated AQI proposed by Rao and Rao (1979), which uses the percentage mean of measured concentrations normalized by their corresponding NAAQS. Their analysis at an industrialized region, for PM₁₀, SPM, SO₂ and NO₂, indicated moderate to heavy air pollution, for most part of the year. Bishoi *et al.* (2009) used yearlong concentrations of CO, SO₂, PM₁₀, O₃ and NO₂, to compare AQI estimated using the US methodology and factor analysis. Results show that while both methodologies follow similar trends, the US methodology estimated higher AQI than factor analysis.

However, in order to quantify the health risk posed by a pollutant, it is essential to study the association between pollutant exposure and health outcome. These studies also aid regulatory agencies in fixing the pollution reduction targets to a level that would minimize the health risk of the exposed group. Health risk based studies are quite common in western countries. Cohen *et al.* (2005) observed that the relationship between relative risk and cardiopulmonary diseases, lung cancer, and acute respiratory infections in children was linear between PM_{2.5} concentrations of 7.5 µg m⁻³ to 50 µg m⁻³, and flattened thereafter. Pope *et al.* (2009) observed that a nonlinear power function expressed the relationship between relative risk of ischemic heart disease, cardiovascular disease and cardiopulmonary disease mortality from cigarette smoking. Results also indicated that initially cardiovascular disease mortality increased steeply with increase in concentration of fine particulate matter and flattened out at higher exposure concentrations. In India, mainly due to lack of mortality data, there are very few studies relating health risk and pollutant concentrations. Dholakia *et al.* (2014) studied the short term association between PM₁₀ concentration and mortality for five Indian cities. Balakrishnan *et al.* (2011) and Rajarathnam *et al.* (2011) did a time series study on Chennai and New Delhi respectively and obtained the risk co-efficient of PM₁₀, NO₂ and SO₂. Similarly, Guttikunda and Goel (2013) studied the health impacts of PM₁₀ and PM_{2.5} in New Delhi using exposure response coefficients from Atkinson *et al.* (2012). Chowdhury and Dey (2016) used satellite data for calculating nationwide PM_{2.5} exposure and a risk model to estimate cause specific premature death from ambient PM_{2.5} exposure. However, no short-term exposure response study of PM_{2.5}, which is the best indicator to the health risk levels from air pollution (WHO, 2014b), was carried out to our knowledge in India.

The objective of this study is to identify the most dangerous pollutant in New Delhi by AQI and health risk based approaches using ambient concentrations of regulated pollutants and health based mortality in New Delhi during 2011 to 2014.

MATERIAL AND METHODS

Indian AQI

CPCB set guidelines for Indian national ambient air quality standards of 12 pollutants (CPCB, 2009). Out of which 8 pollutants CO, NO₂, SO₂, PM_{2.5}, PM₁₀, O₃, Pb and NH₃ have short term standards. To inform people about the quality of air quickly so that people can take appropriate measures to protect themselves, IND-AQI was released in 2014. The details of IND-AQI are available elsewhere (CPCB, 2014), and only briefly summarized here. IND-AQI considers concentrations of PM₁₀, PM_{2.5}, NO₂, O₃, CO, SO₂, NH₃ and Pb. In a day, while maximum 8 hour running average concentrations of O₃ and CO are used, 24 hour averaged concentrations of other six pollutants are used in calculation. The concentration of each pollutant is converted to a number on a scale of 0–500. The sub AQI (AQI_i) for each pollutant (i) is calculated using Eq. (1)

$$AQI_i = \frac{I_{HI} - I_{LO}}{BR_{HI} - BR_{LO}} \times (C_i - BR_{LO}) + I_{LO} \quad (1)$$

where, C_i is the concentration of pollutant 'i'; BR_{HI} and BR_{LO} are breakpoint concentrations greater and smaller to C_i and I_{HI} and I_{LO} are corresponding AQI ranges.

The overall AQI, IND-AQI, can be estimated only if the concentrations of minimum three pollutants are available, with at least one of them being either PM_{2.5} or PM₁₀. The IND-AQI is then taken as the maximum AQI_i of the constituent pollutants, denoted as dominating pollutant. The IND-AQI is divided into five categories: good, satisfactory, moderate, poor, very poor and severe depending on whether the AQI falls between 0–50, 51–100, 101–200, 201–300, 301–400 or 401–500, respectively. IND-AQI calculation can be better understood by the following example: Consider a day in New Delhi with the 24-hr concentrations of PM_{2.5}, SO₂, NO₂, and maximum 8-hr concentration of CO, O₃ as 135 µg m⁻³, 13 µg m⁻³, 12 µg m⁻³, 3 mg m⁻³, and 84 µg m⁻³, respectively. Using the breakpoint concentrations in Table 1 and Eq. (1), the AQI_i of PM_{2.5}, SO₂, NO₂, CO and O₃ are calculated as 311.75, 16.25, 15, 112.27 and 84, respectively. The IND-AQI of that day would be 311.75, and PM_{2.5} would be termed as the dominant pollutant.

Health-Risk Associated with a Pollutant

This study uses excessive risk of the pollutants (ER_i) given by Cairncross *et al.* (2007), as shown in Eq. (2):

$$ER_i = \exp(\beta_i(C_i - C_{min,i})) - 1, \quad C_i > C_{min,i} \quad (2)$$

where, β_i is the exposure-response relationship coefficient, represents the increase in mortality per unit increase in concentrations.

The time series Poisson regression models are widely used to analyze the relation between pollutant concentrations and mortality (Dholakia *et al.*, 2014). The Poisson model used in this study, represented using Eq. (3), is described elsewhere (Bhaskaran *et al.*, 2013; Imai *et al.*, 2015).

Table 1. Breakpoints of different pollutants in IND-AQI (CPCB, 2014).

AQI Category (Range)	PM _{2.5} 24-hr	NO ₂ 24-hr	O ₃ 8-hr	CO 8-hr	SO ₂ 24-hr
Good (0–50)	0–30	0–40	0–50	0–1.0	0–40
Satisfactory (51–100)	31–60	41–80	51–100	1.1–2.0	41–80
Moderate (101–200)	61–90	81–180	101–168	2.1–10	81–380
Poor (201–300)	91–120	181–280	169–208	10.1–17	381–800
Very Poor (301–400)	121–250	281–400	209–748	17.1–34	801–1600
Severe (401–500)	250+	400+	748+	34+	1600+

Note: While CO concentrations are expressed in mg m⁻³; the other pollutants are expressed in µg m⁻³.

$$\log(\mu_t) = \gamma + \beta C_t + \alpha K C_t + f(t) \quad (3)$$

where t denotes time; α , γ and β are the regression coefficients; μ is related to mortality; K denotes temperature; and $f(t)$ is a smooth function of time. The spline function of time is used to remove seasonal and long term trends so that short term variation between concentration and mortality can be studied (Bhaskaran *et al.*, 2013).

In Eq. (2), C_i refers to the measured concentration and $C_{\min,i}$ refers to the concentration of a pollutant below which no adverse health effects can be expected. According to WHO (2005), adverse health effects can be expected for any concentration of PM_{2.5}, O₃, NO₂ and SO₂. Thus, in this study $C_{\min,i}$ of all pollutants except CO, is considered as 0. $C_{\min,i}$ for CO is considered as 2 mg m⁻³, as suggested by (CPCB, 2014), and shown in Table 1.

Study Area and Data Sources

New Delhi, in addition to being the capital of India, is also one of the most densely populated cities in the world. It has a population of 16.7 million with an annual average growth rate of 1.92%. The overall population density is 11,297 per sq. km. It faces extreme temperatures of as low as 7°C in winter to as high as 48°C in summer (WU, 2016). It received an average rainfall of 889.2 mm from 2011–2014. Such extreme temperatures can severely affect the already deteriorating air pollution in New Delhi. Four years data ranging from January 1st, 2011 to December 31st of 2014 was used for the analysis. The data was collected at a busiest traffic intersection, Bahadur Shah Zafar Marg (ITO) located at commercial down town of the city, as shown in Fig. 1. Hourly concentrations of CO, O₃, NO₂, SO₂, and PM_{2.5} were collected by Central Pollution Control Board (CPCB). Pollutant concentrations vary across a city. However, this is the longest period for which hourly data of these species is available in any location in New Delhi. In such situations studies resort to analysis of data collected at a single location. For example, Kim *et al.* (2015) studied the association of selected components of PM_{2.5} on mortality, using PM_{2.5} data obtained from a centrally located residential site in Denver. Similarly, Garrett and Casimiro (2011) studied the short term effect of PM_{2.5} and O₃ in Lisbon, using data obtained from a monitoring station in Olivais since this was the only station with data for both pollutants. Thus, in this study it is assumed that the concentrations observed in this location are representative of the entire city.

CO was measured using non-dispersive infrared

spectroscopy, O₃ and NO₂ using Chemiluminescence, SO₂ using ultra violet fluorescence, and PM_{2.5} using tapered element oscillating microbalance. As hourly PM₁₀ concentrations were not available in this location, PM_{2.5} is assumed to be the sole representative of particulate matter in this study. Due to unavailability of data, the percentage of days on which the IND-AQI was not calculated, was 19, 42, 41 and 60 in 2011, 2012, 2013 and 2014, respectively. The mortality data was obtained from the office of births and deaths registration in New Delhi. Due to unavailability of cause specific mortality, non-accidental mortality has been used in this study.

RESULTS AND DISCUSSION

Yearly Variation of Ambient Air Pollutant Concentrations

In INAAQS, CO and O₃ have both hourly and eight hour standards, and PM_{2.5}, SO₂ and NO₂ have daily and yearly standards. For comparison with INAAQS, the measured hourly data of PM_{2.5}, SO₂ and NO₂ were converted into daily averaged concentrations, and the hourly concentrations of O₃ and CO were used to estimate daily maximum 8 hour running average. Fig. 2 shows the change in concentrations of PM_{2.5}, SO₂, O₃, NO₂ and CO, from 2011 to 2014. The percentage of days for each pollutant exceeded INAAQS is also shown. Results indicate that in contrary to other pollutants, PM_{2.5} had maximum median concentrations in 2013. Panel (a) in Fig. 2 shows that median PM_{2.5} concentrations in all the four years exceeded the corresponding INAAQS of 100 µg m⁻³. PM_{2.5} concentrations exceeded INAAQS during 80, 82, 86 and 90% of days in 2011, 2012, 2013 and 2014, respectively.

SO₂ concentrations never exceeded INAAQS in the four years, as observed from panel (b) of the figure. Similar conclusion was derived by Goyal and Sidhartha (2003) from their air quality analysis during 1996 to 2001 at New Delhi. This indicates that SO₂ may not be a major air pollutant in New Delhi. However, the contribution of SO₂ in the formation of secondary sulfate particles might be significant, and should be explored further. Panel (c) shows that O₃ concentrations varied from 5 to 323 µg m⁻³ during the analysis period. O₃ concentrations exceeded the corresponding INAAQS of 100 µg m⁻³ for 16% in 2011, 26% in 2012, 0% in 2013, and 12% in 2014. Panel (d) shows that NO₂ concentrations exceeded the INAAQS of 80 µg m⁻³ limit 33.67% of the days in 2011, 62% of days in 2012, 15% of days in 2013, and 36% of days in 2014. CO concentrations reached a maximum of 10 mg m⁻³ during




Fig. 1. An interactive map showing the study region (Delhi) and the location of the monitoring station (ITO) in India.




Fig. 2. Change in concentrations of CO, NO₂, O₃, SO₂ and PM_{2.5} from 2011 to 2014 at New Delhi. While the left Y-axis shows the concentrations, represented using box whisker plots, the right Y-axis, represented using red crosses indicates the percentage of data which exceed the INAAQS. The box whisker plots show minimum, maximum, median, upper and lower quartiles. Note: The range of y-axis of all panels is curtailed for better representation of the figure.

the period. Panel (e) shows that CO exceeded corresponding INAAQS value of 2 mg m^{-3} ; 14%, 11%, 9.4% and 9.2% of days in 2011, 2012, 2013 and 2014, respectively. This implies that $\text{PM}_{2.5}$ is the major pollutant, followed by NO_2 and CO in New Delhi. However, more studies in New Delhi where both PM_{10} and $\text{PM}_{2.5}$ concentrations are available are necessary in future to support this conclusion.

Yearly Variation of IND-AQI

The yearly variation of frequency of days falling in the six IND-AQI categories, good, satisfactory, moderate, poor, very poor and severe, during 2011–2014 is shown in Fig. 3. Due to the lack of availability of data of other pollutants only concentrations of $\text{PM}_{2.5}$, CO, NO_2 , SO_2 and O_3 were used to estimate the IND-AQI in this study. The corresponding frequency of dominating species in each of those years is shown in Table 2.

$\text{PM}_{2.5}$ was the dominant species in 74 to 90% of the days in those four years. Maximum dominance of NO_2 was observed during 2011 (16%), and O_3 during 2011 (8%). While none of these years had days dominated by SO_2 , CO dominance was around 9% in most of the years, except 2012. Zero dominant days of SO_2 is expected as it never violates INAAQS, as observed from Fig. 2.

Fig. 3 indicates that at least 60% of the days in all the years were poor. Moreover, 70% of days in 2013 were either very poor or severe. In comparison, 44, 53 and 52% of such days exist in 2011, 2012 and 2014, respectively. This could be due to higher $\text{PM}_{2.5}$ concentrations in 2013 as observed in Fig. 2. Only 6% of the days in all the four years were either good or satisfactory.

Weekday-Weekend Variation of IND-AQI

Data analysts have reported weekday and weekend differences in air pollutant concentrations around the world (Altshuler *et al.*, 1995; Karar *et al.*, 2006; Tiwari *et al.*, 2015). To explore this, weekdays and weekend variation of IND-AQI was analyzed, and shown in Fig. 4. $\text{PM}_{2.5}$ was

the dominant pollutant during 83% and 82% on weekdays and weekends, respectively. Results indicate that air quality on weekends was only slightly better than weekdays. For example, 50% of weekends, in comparison to 52% of weekdays were in the IND-AQI category of very poor and severe. Similar conclusions were arrived by Kumar and Goyal (2011), who studied AQI, following the US methodology, at the same location during 2000 to 2006. However, these observations are in contrary to the assumption that at the sampling site, located at a busy traffic junction, where vehicle density is more on weekdays than weekends, air quality should be better on weekends.

Seasonal Variation of IND-AQI

Previous studies have shown higher concentrations of poly aromatic hydrocarbons, sulfates and nitrates in $\text{PM}_{2.5}$ in New Delhi (Pant *et al.*, 2015), which have a strong correlation with temperature and RH (Wang *et al.*, 2005). Additionally, studies in this region showed a strong correlation of reactive pollutants like O_3 , with temperature and RH (Gaur *et al.*, 2014). To explore this, seasonal differences in IND-AQI and dominant species as a function of relative humidity (RH) and temperature were studied, and shown in Fig. 5. The seasons were categorized as winter (December–February), pre-monsoon (March–May), monsoon (June–August) and post-monsoon (September–November). Analysis indicates that concentrations of all the pollutants decreased during monsoon. This is due to wet scavenging of pollutants due to higher precipitation during that season.

Panels (a) to (d), indicate that winter had highest, around 72%, and monsoon had least, 32%, very poor and severe days. Panels (e) to (h) show a clear decreasing trend of $\text{PM}_{2.5}$ dominance from winter to post-monsoon. While, $\text{PM}_{2.5}$ was dominant in 95% of days in winter, it dominated IND-AQI during 68 and 70% of days of monsoon and post-monsoon seasons, respectively. Moreover, CO dominated during 18% of days in monsoon, and both CO and NO_2 dominated post-monsoon by around 13–14% of days each.




Fig. 3. Change in frequency of days (%) in various IND-AQI categories from 2011–2014.

Table 2. Variation of domination of pollutants during 2011 to 2014.

Year	SO ₂	NO ₂	PM _{2.5}	CO	O ₃
2011	0	2	81	9	8
2012	0	17	75	4	4
2013	0	2	88	10	0
2014	0	0	90	9	1

Previous studies have indicated that in addition to wet deposition, variation in atmospheric mixing layer heights, air flow pattern can also be a main reason for seasonal differences of pollutant concentrations in this region (Sahu *et al.*, 2009; Bisht *et al.*, 2015). To examine this further, 24-hr back trajectories, originating from the sampling location, were generated using National Oceanic and Atmospheric Administration's Hybrid Single Particle Lagrangian Integrated Trajectory (HYSPPLIT) model. Fig. S1, in the supplementary material, shows the diurnal variation in atmospheric mixing layer heights in different seasons.

Analysis indicated that mixing layer heights peak in the afternoon, around 2 to 3 PM, with the least daily maximum mixing layer heights during winter.

Results in Fig. S2, in the supplementary material, show a clear seasonal pattern in air mass back trajectories in New Delhi. During winter and pre-monsoon, where more very poor and severe days are observed, air parcels might have originated in Punjab and Haryana, home of many coal-based power plants and industries. In contrary, air parcels could have originated in Rajasthan and Uttar Pradesh during monsoon, and Rajasthan and Punjab in post-monsoon. Thus, the seasonal differences and the negligible weekday-weekend difference observed in this study could be due to long range transport of air pollutants. Similar conclusions were arrived from a recent study (Ghosh *et al.*, 2015), which predicted significant influence of neighboring states in PM_{2.5} concentrations, during 2008 to 2010, in New Delhi.

Estimation of Health Risk Associated with Pollutants

To estimate the potential risk associated with the




Fig. 4. Days (%) in each IND-AQI category on weekdays and weekends. Note: Data in all the four years was used in analysis.




Fig. 5. Scatter plot of relative humidity versus temperature as a function of IND-AQI (panels a–d), and dominant species (panels e–h), for different seasons; winter, pre-monsoon, monsoon and post-monsoon.

concentrations of a pollutant, excessive risk associated with a pollutant was calculated using Eq. (2). Fig. 6 shows the monthly averaged $PM_{2.5}$ concentrations, temperature and non-accidental deaths at New Delhi during the analysis period. Estimated increase in non-accidental mortality, was 0.69% (95% CI: 0.17%, 1.21%), 0.88% (95% CI: 0.14%, 1.63%), and 3.77% (95% CI: -0.6%, 8.34%) per $10 \mu\text{g m}^{-3}$ increase of $PM_{2.5}$, NO_2 , and $1 \text{ mg m}^{-3} CO$, respectively.

The ER values were obtained as 0.57 (95% CI: 0.45, 0.69), 0.36 (95% CI: 0.24, 0.46), and 0.052 (95% CI: 0.046, 0.058) for $PM_{2.5}$, NO_2 , and CO , respectively. Non-significant risk factors were obtained for SO_2 and Ozone. This clearly indicates that $PM_{2.5}$ is associated with major health risk in New Delhi. Even though, Eq. (2) is commonly used (for example, see Hu *et al.* (2015)), it doesn't consider the possible non-linearities in the exposure-response curve observed by some previous studies (Cohen *et al.*, 2005; Pope *et al.*, 2009; Burnett *et al.*, 2014). Moreover, most of the health studies tend to depict the correlation between $PM_{2.5}$ concentrations and specific respiratory or heart diseases. However, in New Delhi disease specific mortality was not available during the analysis period. To study this further,

the RR for average $PM_{2.5}$ concentration of $133 \mu\text{g m}^{-3}$ at New Delhi was compared to that of Burnett *et al.* (2014). While, RR was 1.57 from the Eq. (2) used in this study, it was 1.4585, 1.658 and 2.662 for COPD, lung cancer and ALRI mortality, respectively from the integrated exposure function used in Burnett *et al.* (2014). Thus the overall risk of mortality obtained in this study lies within the range of values obtained using integrated risk function.

Table 3 shows the % increase in non-accidental mortality per $10 \mu\text{g m}^{-3}$ increase in $PM_{2.5}$ concentration estimated by different studies around the world. In comparison with other major countries, no such studies were done in India. This indicates that exposure-response relation for $PM_{2.5}$ is similar in major cities in the world. This value lies in the range of global estimate by Atkinson *et al.* (2014), and closer to several studies in China (Dai *et al.*, 2004; Chen *et al.*, 2011) and the US (Ostro *et al.*, 2006).

Furthermore, to better cogitate its associated health benefits, the possible number of non-accidental deaths averted by reducing $PM_{2.5}$ concentrations, to suggested levels by WHO and INAAQS was estimated following Shang *et al.* (2013). The expected number of premature deaths (PD_i)




Fig. 6. Time series of monthly average $PM_{2.5}$ ($\mu\text{g m}^{-3}$) and deaths in left y-axis and temperature (C) in right y-axis.

Table 3. Increase in non-accidental mortality (%) per $10 \mu\text{g m}^{-3}$ increase of $PM_{2.5}$ at different cities around the world.

Place (Study Year)	Reference	% increase in mortality
Global Review	(Atkinson <i>et al.</i> , 2014)	1.04 (0.52–1.56)
USA (1970–1986)	(Klemm, 2003)	1.2 (0.8–1.6)
USA (1999–2005)	(Zanobetti and Schwartz, 2009)	0.98 (0.75–1.22)
USA (1997–2002)	(Franklin <i>et al.</i> , 2007)	1.2 (0.3–2.1)
USA (2000–2008)	(Levy <i>et al.</i> , 2012)	1.2 (0.5–1.9)
California (1999–2002)	(Ostro <i>et al.</i> , 2006)	0.6 (0.2–1.0)
Spain (2003–2007)	(Ostro <i>et al.</i> , 2011)	1.4 (0.6–2.3)
Australia (1996–1999)	(Simpson <i>et al.</i> , 2005)	0.9 (-0.7–2.5)
Beijing (2007–2008)	(Chen <i>et al.</i> , 2011)	0.53 (0.37–0.69)
Guangzhou (2007–2008)	(Yang <i>et al.</i> , 2012)	0.90 (0.55–1.26)
Shenyang (2006–2008)	(Chen <i>et al.</i> , 2011)	0.35 (0.17–0.53)
Shanghai (2004–2005)	(Huang <i>et al.</i> , 2009)	0.30 (0.06–0.54)
Shanghai (2002–2003)	(Dai <i>et al.</i> , 2004)	0.85 (0.32–1.89)
Xian (2004–2008)	(HOU Bin, 2011)	0.20 (0.07–0.33)
New Delhi (2011–2014)	Current Study	0.69 (0.17–1.21)

Table 4. Projected reduction in non-accidental mortality and corresponding reduction in non-accidental deaths, due to proposed reduction in PM_{2.5} concentrations for New Delhi.

Proposed PM _{2.5} limit ($\mu\text{g m}^{-3}$)	Reduction in non-accidental mortality (%)	Reduction in non-accidental deaths (per 100000 people)
40	6.20	39
35	6.52	41
25	7.17	45
15	7.80	49
10	8.12	51

due to a pollutant exposure can be calculated by Eq. (4), using mortality rate (M).

$$PD_i = M \times \frac{ER_i}{ER_i + 1} \quad (4)$$

The excessive risk (ER_i), in Eq. (4), was calculated by Eq. (2), using C_i as the averaged PM_{2.5} concentration during the analysis period, and the $C_{\min,i}$ as the targeted PM_{2.5} limit.

Decrease of PM_{2.5} concentrations from current level to 10 $\mu\text{g m}^{-3}$ will result in a reduction of 8826 deaths in 2011. This is similar to the estimated deaths by (Guttikunda and Goel, 2013) due to PM_{2.5} in New Delhi as 7350–16200 in the year 2010. Table 4 gives the projected reduction in non-accidental mortality and number of reduction in non-accidental deaths due to decrease in current PM_{2.5} concentration to the levels suggested by INAAQS and WHO. The averaged non-accidental mortality rate data for the year 2011 to 2014 was obtained from statistical handbooks of Delhi during those years (DES, 2014) as 0.625%.

Results indicate that, when the current levels are reduced to meet INAAQS annual limits of PM_{2.5} i.e., 40 $\mu\text{g m}^{-3}$, the non-accidental mortality in New Delhi will be reduced by 6.20%, with 39 premature deaths avoided per 100000 people. Similarly, if the WHO levels of 35, 25, 15 and 10 $\mu\text{g m}^{-3}$ of PM_{2.5} are met, the number of premature deaths per 100000 people will be reduced by 41, 45, 49 and 51, respectively.

SUMMARY AND CONCLUSIONS

This paper analyzed data collected at a busy traffic junction during 2011 to 2014 at New Delhi, to determine the most dangerous pollutant in New Delhi. AQI was used to identify the dominant pollutant while a health risk study quantified the deaths due to high concentration of the pollutant. Investigation showed that PM_{2.5} was the dominant pollutant, during all the seasons, with dominant days being 24% higher in winter and pre-monsoon, than monsoon and post-monsoon. Moreover, significant differences in very poor and severe days were not observed on weekdays and weekends, with PM_{2.5} being the dominant pollutant in all days. However, this conclusion might vary if PM₁₀ data is included in the analysis, and more studies are needed to explore this further. Additional investigation revealed that, this could be due to long range transport of air pollutants from different regions. This shows that air quality in New Delhi can be ameliorated only due to better policies in

neighboring states.

The potential health risk associated with PM_{2.5} was greater than CO and NO₂. The excessive risk associated with PM_{2.5}, NO₂ and CO were obtained as 0.57 (95% CI: 0.45, 0.69), 0.36 (95% CI: 0.24, 0.46), and 0.052 (95% CI: 0.046, 0.058), respectively. This is in agreement with the AQI method. Finally, the impact of reducing the current PM_{2.5} concentrations, the most dominant pollutant in New Delhi was investigated. Results indicated that 39 and 41 premature deaths can be avoided per 100000 by bringing down the yearly averaged concentrations of PM_{2.5} to the levels suggested by INAAQS and WHO, respectively.

ACKNOWLEDGEMENTS

The authors would like to thank Indian Institute of Technology, Guwahati for providing facilities to carry out this research. Authors would also like to thank the Central Pollution Control Board at New Delhi for sharing the concentrations of pollutants measured at the site. The authors also thank the anonymous reviewers for their inputs.

SUPPLEMENTARY MATERIAL

Supplementary data associated with this article can be found in the online version at <http://www.aaqr.org>.

REFERENCES

- Altshuler, S.L., Arcado, T.D. and Lawson, D.R. (1995). Weekday vs. weekend ambient ozone concentrations: Discussion and hypotheses with focus on Northern California. *J. Air Waste Manage. Assoc.* 45: 967–972.
- Aneja, V.P., Agarwal, A., Roelle, P.A., Phillips, S.B., Tong, Q.S., Watkins, N. and Yablonsky, R. (2001). Measurements and analysis of criteria pollutants in New Delhi, India. *Environ. Int.* 27: 35–42.
- Atkinson, R.W., Cohen, A., Mehta, S. and Anderson, H.R. (2012). Systematic review and meta-analysis of epidemiological time-series studies on outdoor air pollution and health in Asia. *Air Qual. Atmos. Health* 5: 383–391.
- Atkinson, R.W., Kang, S., Anderson, H.R., Mills, I.C. and Walton, H.A. (2014). Epidemiological time series studies of PM_{2.5} and daily mortality and hospital admissions: A systematic review and meta-analysis. *Thorax* 69: 660–665.
- Balakrishnan, K., Ganguli, B., Ghosh, S., Sankar, S.,

- Thanasekaraan, V., Rayudu, V.N. and Caussy, H. (2011). Part 1. Short-term effects of air pollution on mortality: Results from a time-series analysis in Chennai, India. *Res. Rep. Health Eff. Inst.* 157: 7–44.
- Banerjee, T. and Srivastava, R.K. (2011). Assessment of the ambient air quality at the Integrated Industrial Estate-Pantnagar through the air quality index (AQI) and exceedance factor (EF). *Asia-Pac. J. Chem. Eng.* 6: 64–70.
- Bhaskar, B.V. and Mehta, V.M. (2010). Atmospheric particulate pollutants and their relationship with meteorology in Ahmedabad. *Aerosol Air Qual. Res.* 10: 301–315.
- Bhaskaran, K., Gasparrini, A., Hajat, S., Smeeth, L. and Armstrong, B. (2013). Time series regression studies in environmental epidemiology. *Int. J. Epidemiol.* 42: 1187–1195.
- Bishoi, B., Prakash, A. and Jain, V.K. (2009). A comparative study of air quality index based on factor analysis and US-EPA methods for an urban environment. *Aerosol Air Qual. Res.* 9: 1–17.
- Bisht, D.S., Dumka, U.C., Kaskaoutis, D.G., Pipal, A.S., Srivastava, A.K., Soni, V.K., Attri, S.D., Sateesh, M. and Tiwari, S. (2015). Carbonaceous aerosols and pollutants over Delhi urban environment: Temporal evolution, source apportionment and radiative forcing. *Sci. Total Environ.* 521: 431–445.
- Burnett, R.T., Pope, C.A., 3rd, Ezzati, M., Olives, C., Lim, S.S., Mehta, S., Shin, H.H., Singh, G., Hubbell, B., Brauer, M., Anderson, H.R., Smith, K.R., Balmes, J.R., Bruce, N.G., Kan, H., Laden, F., Pruss-Ustun, A., Turner, M.C., Gapstur, S.M., Diver, W.R. and Cohen, A. (2014). An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter Exposure. *Environ. Health Perspect.* 122: 397–403.
- Cairncross, E.K., John, J. and Zunckel, M. (2007). A novel air pollution index based on the relative risk of daily mortality associated with short-term exposure to common air pollutants. *Atmos. Environ.* 41: 8442–8454.
- Chen, H. and Copes, R. (2013). *Review of Air Quality Index and Air Quality Health Index*, Ontario Agency for Health Protection and Promotion Public Health Ontario, Toronto, Ontario.
- Chen, R., Li, Y., Ma, Y., Pan, G., Zeng, G., Xu, X., Chen, B. and Kan, H. (2011). Coarse particles and mortality in three Chinese cities: the China Air Pollution and Health Effects Study (CAPES). *Sci. Total Environ.* 409: 4934–4938.
- Chowdhury, S. and Dey, S. (2016). Cause-specific premature death from ambient PM_{2.5} exposure in India: Estimate adjusted for baseline mortality. *Environ. Int.* 91: 283–290.
- Cohen, A.J., Ross Anderson, H., Ostro, B., Pandey, K.D., Krzyzanowski, M., Kunzli, N., Gutschmidt, K., Pope, A., Romieu, I., Samet, J.M. and Smith, K. (2005). The global burden of disease due to outdoor air pollution. *J. Toxicol. Environ. Health Part A* 68: 1301–1307.
- COMEAP (2011). Review of UK Air Quality Index, Committee on the Medical Effects of Air Pollutants London.
- CPCB (2009). National Ambient Air Quality Standards, Central Pollution Control Board, New Delhi, India.
- CPCB (2014). National Air Quality Index Report, Central Pollution Control Board, New Delhi, India.
- Dai, H., Song, W., Gao, X. and Chen, L. (2004). Study on relationship between ambient PM₁₀, PM_{2.5} pollution and daily mortality in a district in Shanghai. *Wei Sheng Yan Jiu J. Hyg. Res.* 33: 293–297.
- DES (2014). Statistical Handbook of Delhi, 2011–2014, Directorate of economics & statistics
- Dholakia, H.H., Bhadra, D. and Garg, A. (2014). Short term association between ambient air pollution and mortality and modification by temperature in five Indian cities. *Atmos. Environ.* 99: 168–174.
- Franklin, M., Zeka, A. and Schwartz, J. (2007). Association between PM_{2.5} and all-cause and specific-cause mortality in 27 US communities. *J. Exposure Sci. Environ. Epidemiol.* 17: 279–287.
- Garrett, P. and Casimiro, E. (2011). Short-term effect of fine particulate matter (PM_{2.5}) and ozone on daily mortality in Lisbon, Portugal. *Environ. Sci. Pollut. Res.* 18: 1585–1592.
- Gaur, A., Tripathi, S.N., Kanawade, V.P., Tare, V. and Shukla, S.P. (2014). Four-year measurements of trace gases (SO₂, NO_x, CO, and O₃) at an urban location, Kanpur, in Northern India. *J. Atmos. Chem.* 71: 283–301.
- Ghosh, S., Biswas, J., Guttikunda, S., Roychowdhury, S. and Nayak, M. (2015). An investigation of potential regional and local source regions affecting fine particulate matter concentrations in Delhi, India. *J. Air Waste Manage. Assoc.* 65: 218–231.
- Goyal, P. and Sidhartha (2003). Present scenario of air quality in Delhi: A case study of CNG implementation. *Atmos. Environ.* 37: 5423–5431.
- Guttikunda, S.K. and Goel, R. (2013). Health impacts of particulate pollution in a megacity—Delhi, India. *Environ. Dev.* 6: 8–20.
- Hou, B., Dai, L.Z., Wang, Z. *et al.* (2011). Time-series analysis of acute mortality effects of air pollution in Xi'an. *J. Environ. Health* 28: 1039–1043 [in Chinese].
- Hu, J., Ying, Q., Wang, Y. and Zhang, H. (2015). Characterizing multi-pollutant air pollution in China: Comparison of three air quality indices. *Environ. Int.* 84: 17–25.
- Huang, W., Tan, J., Kan, H., Zhao, N., Song, W., Song, G., Chen, G., Jiang, L., Jiang, C., Chen, R. and Chen, B. (2009). Visibility, air quality and daily mortality in Shanghai, China. *Sci. Total Environ.* 407: 3295–3300.
- IHME (2013). India Global Burden of Disease Study 2010. Institute for Health Metrics and Evaluation, Seattle, United States.
- Imai, C., Armstrong, B., Chalabi, Z., Mangtani, P. and Hashizume, M. (2015). Time series regression model for infectious disease and weather. *Environ. Res.* 142: 319–327.
- Karar, K., Gupta, A.K., Kumar, A., Biswas, A.K. and Devotta, S. (2006). Statistical interpretation of

- weekday/weekend differences of ambient particulate matter, vehicular traffic and meteorological parameters in an urban region of Kolkata, India. *Indoor Built Environ.* 15: 235–245.
- Kim, S.Y., Dutton, S.J., Sheppard, L., Hannigan, M.P., Miller, S.L., Milford, J.B., Peel, J.L. and Vedal, S. (2015). The short-term association of selected components of fine particulate matter and mortality in the Denver Aerosol Sources and Health (DASH) study. *Environ. Health* 14: 1–11.
- Klemm, R.J. and Mason, R. (2003). Replication of reanalysis of harvard six-city mortality study. In *Revised Analyses of Time-Series of Air Pollution and Health*. Special Report. Health Effects Institute, Boston, MA, pp. 165–172.
- Kumar, A. and Goyal, P. (2011). Forecasting of daily air quality index in Delhi. *Sci. Total Environ.* 409: 5517–5523.
- Levy, J.I., Diez, D., Dou, Y., Barr, C.D. and Dominici, F. (2012). A meta-analysis and multisite time-series analysis of the differential toxicity of major fine particulate matter constituents. *Am. J. Epidemiol.* 175: 1091–1099.
- Ostro, B., Broadwin, R., Green, S., Feng, W.Y. and Lipsett, M. (2006). Fine particulate air pollution and mortality in nine California Counties: Results from calfine. *Environ. Health Perspect.* 114: 29–33.
- Ostro, B., Tobias, A., Querol, X., Alastuey, A., Amato, F., Pey, J., Pérez, N. and Sunyer, J. (2011). The effects of particulate matter sources on daily mortality: A case-crossover study of barcelona, Spain. *Environ. Health Perspect.* 119: 1781–1787.
- Ott, W.R. (1978). *Environ Indices: Theo and Prac*. Ann Arbor Science Publishers, Inc., Ann Arbor, MI.
- Pant, P., Shukla, A., Kohl, S.D., Chow, J.C., Watson, J.G. and Harrison, R.M. (2015). Characterization of ambient PM_{2.5} at a pollution hotspot in New Delhi, India and inference of sources. *Atmos. Environ.* 109: 178–189.
- Pope, C.A., 3rd, Burnett, R.T., Krewski, D., Jerrett, M., Shi, Y., Calle, E.E. and Thun, M.J. (2009). Cardiovascular mortality and exposure to airborne fine particulate matter and cigarette smoke: Shape of the exposure-response relationship. *Circulation* 120: 941–948.
- Rajarithnam, U., Sehgal, M., Nairy, S., Patnayak, R.C., Chhabra, S.K., Kilnani and Ragavan, K.V. (2011). Part 2. Time-series study on air pollution and mortality in Delhi. *Res. Rep. Health Eff. Inst.* 157: 47–74.
- Rao, M. and Rao, H. (1979). *Air Pollution*. McGraw Hill Education (India) Private Limited, New Delhi.
- Sadavarte, P. and Venkataraman, C. (2014). Trends in multi-pollutant emissions from a technology-linked inventory for India: I. Industry and transport sectors. *Atmos. Environ.* 99: 353–364.
- Sahu, L.K., Lal, S., Thouret, V. and Smit, H.G. (2009). Seasonality of tropospheric ozone and water vapor over Delhi, India: A study based on mosaic measurement data. *J. Atmos. Chem.* 62: 151–174.
- Sahu, S.K., Beig, G. and Parkhi, N.S. (2011). Emissions inventory of anthropogenic PM_{2.5} and PM₁₀ in Delhi during Commonwealth Games 2010. *Atmos. Environ.* 45: 6180–6190.
- Shang, Y., Sun, Z., Cao, J., Wang, X., Zhong, L., Bi, X., Li, H., Liu, W., Zhu, T. and Huang, W. (2013). Systematic review of chinese studies of short-term exposure to air pollution and daily mortality. *Environ. Int.* 54: 100–111.
- Sharma, M., Pandey, R., Maheshwari, M., Sengupta, B., Shukla, B.P., Gupta, N.K. and Johri, S. (2003). Interpretation of air quality data using an air quality index for the city of Kanpur, India. *J. Environ. Eng. Sci.* 2: 453–462.
- Simpson, R., Williams, G., Petroeschovsky, A., Best, T., Morgan, G., Denison, L., Hinwood, A., Neville, G. and Neller, A. (2005). The short-term effects of air pollution on daily mortality in four Australian Cities. *Aust. N. Z. J. Public Health* 29: 205–212.
- Tiwari, S., Bisht, D.S., Srivastava, A.K. and Gustafsson, O. (2015). Simultaneous measurements of black carbon and PM_{2.5}, CO, and NO_x variability at a locally polluted urban location in India. *Nat. Hazard.* 75: 813–829.
- Wang, Y., Zhuang, G.S., Tang, A.H., Yuan, H., Sun, Y.L., Chen, S.A. and Zheng, A.H. (2005). The ion chemistry and the source of PM_{2.5} aerosol in Beijing. *Atmos. Environ.* 39: 3771–3784.
- WHO (2005). *Who Air Quality Guidelines Global Update 2005*. Particulate Matter, Ozone, Nitrogen Dioxide and Sulphur Dioxide, World Health Organization, Copenhagen, Denmark.
- WHO (2014a). *World Health Statistics 2014*.
- WHO (2014b). *Air Quality Deteriorating in Many of the World's Cities*, <http://www.who.int/mediacentre/news/releases/2014/air-quality/en/>, Last Access: 26 September 2016.
- WU (2016). http://www.Wunderground.Com/History/Airport/Vidp/2011/1/10/Customhistory.Html?Req_City=Delhi&Req_StateName=India&Reqdb.Zip=00000&Reqdb.Magic=8&Reqdb.Wmo=42182, Last Access: 26 January 2016.
- Yang, C., Peng, X., Huang, W., Chen, R., Xu, Z., Chen, B. and Kan, H. (2012). A time-stratified case-crossover study of fine particulate matter air pollution and mortality in Guangzhou, China. *Int. Arch. Occup. Environ. Health* 85: 579–585.
- Zanobetti, A. and Schwartz, J. (2009). The effect of fine and coarse particulate air pollution on mortality: A national analysis. *Environ. Health Perspect.* 117: 898–903.

Received for review, June 26, 2016

Revised, September 29, 2016

Accepted, October 29, 2016