



## Characteristics of Heavy Metals Emitted from a Heavy Oil-Fueled Power Plant in Northern Taiwan

Ya-Fen Wang<sup>1\*</sup>, How-Ran Chao<sup>2</sup>, Lin-Chi Wang<sup>3</sup>, Guo-Ping Chang-Chien<sup>3</sup>, Tsui-Chun Tsou<sup>4</sup>

<sup>1</sup> Department of Bioenvironmental Engineering and R&D Center of Membrane Technology, Chung Yuan Christian University, Chung-Li, 320, Taiwan

<sup>2</sup> Department of Environmental Science and Engineering, National Pingtung University of Science and Technology, Pingtung 912, Taiwan

<sup>3</sup> Department of Chemical and Materials Engineering, Cheng-Shiu University, 833, Kaohsiung, Taiwan

<sup>4</sup> Division of Environmental Health and Occupational Medicine, National Health Research Institutes, Zhunan, Miaoli 350, Taiwan

---

### ABSTRACT

The characteristics and distribution of metal contents emitted from a power plant fueled by heavy oil and its impact to the ambient atmosphere near the power plant was investigated. The current investigation measured toxic (As, Cd, Cr, Hg, Ni and Pb), anthropogenic (Ba, Cu, Mn, Sb, Se, Sr, Ti, V and Zn) and crust (Al, Ca, Fe, K and Mg) elements from a 2,000 MW heavy oil-fired power plant. Results showed the emission concentration from the power plant contributed to 17,976 kg/yr annual emission of anthropogenic elements, which was significantly higher than those from some electrical arc furnaces and coke ovens in Taiwan. For toxic metals, As, Cd or Ni concentration do not exceed target values established by the European Council (2004/107/EC) for As (6 ng/m<sup>3</sup>), Cd (5 ng/m<sup>3</sup>) and Ni (20 ng/m<sup>3</sup>). This study also applies nonparametric statistical analyses for evaluating the relationship between metal concentrations and operational parameters (including emitted CO<sub>2</sub>, O<sub>2</sub>, flue gas emission temperature, flue gas velocity, moisture, heavy oil consumption rate, boiler steam temperature, boiler operational pressure, and electricity). Findings show negative correlations between most toxic metals (As, Cd, Cr and Hg) and operational parameters, though some pairs were not statistically significant. The current study provides only preliminary statistical results between metal concentrations and operational parameters due to small sample sizes. Further investigation requires larger sample sizes.

**Keywords:** Heavy oil power plant; metal; Emission factor; Ambient atmosphere; Operational parameters.

---

### INTRODUCTION

Scientists recognize certain trace metals, such as iron (Fe), zinc (Zn), copper (Cu), chromium (Cr), iodine (I), cobalt (Co), molybdenum (Mo), and selenium (Se), as essential elements for human health and plant nutrition. However, the toxic properties of many metals are a point of concern. (Sarkar, 2002; Saxena *et al.*, 2008). Findings show that exposure to transition metals associated with ambient particles originating from industrial sources elicit airway inflammation in healthy subjects (Ghio and Devlin, 2001; Schaumann *et al.*, 2004; Wang *et al.*, 2008). Highly concentrated airborne metals in industrial areas could play a role in regional prevalence of allergic conditions

(Heinrich *et al.*, 2002; Kuo *et al.*, 2007). This hypothesis is supported by evidence that sensitized mice exposed to PM<sub>2.5</sub> extracts rich in Zn, Mn, Pb, Cu and Cd, exhibited increased airway responsiveness to metacholine and lung inflammatory cells (Gavett *et al.*, 2003). Metals from industrial emissions or traffic have been associated with heart rate variability (Magari *et al.*, 2002), increase in hospital emergency visits for respiratory conditions, and increased risk of lung cancer after long term exposures (Pope Iii *et al.*, 2002). In urban and industrial environments, however, particles not only contain metals, but also include toxic organic compounds, such as PAHs, whose concentrations are greater in fine and ultrafine particles. This fact has caused people concern about various pollution sources around their habitations.

Coal and oil combustion facilities produce fine particles in the submicron size range (Jang *et al.*, 2007), enriched by heavy metals. Jang *et al.* (2007) indicated that heavy oil combustion resulted in relatively pronounced ultra-fine particles smaller than 1.0 μm, difficult to remove by

---


\* Corresponding author. Tel.: +886 3 2654912;  
Fax: +886 3 2654949  
E-mail address: yfwang@cycu.edu.tw.

existing air pollutant control devices (APCD). The diverse PM composition responsible for these health effects is still unknown and attracts researchers' attention, heightening awareness of pollutants emitted from power plants. Some literatures address heavy metal emissions from coal/oil power plants worldwide, such as Finland (Hatanpaa *et al.*, 1997), Greece (Petaloti *et al.*, 2006), Spain (Moreno *et al.*, 2007), Turkey (Yatkin and Bayram, 2008), India (Reddy *et al.*, 2005) and Korea (Jang *et al.*, 2007), however, few literatures refer to Taiwan. This study investigates the characteristics and emission factors of heavy metals from a power plant fueled by heavy oil. Besides the influence of power plant emission on the nearby atmosphere, the current work also evaluates the relationship between heavy metal and operational parameters to provide more information for further research on the effect of flue gas from the power plant.

## EXPERIMENTAL SECTION

The current investigation selects the largest power plant (2,000 MW) near the Pacific Ocean and located in northern Taiwan, for this study (Fig. 1). Sixteen samples were collected from the power plant stack fueled by low-sulfur heavy oil from March 2006 through January 2007. The power plant is equipped with four sets of boilers configured with lower-NO<sub>x</sub> burners and with electrostatic precipitators as air pollution control devices. The oil consumption ranged from 41.1 kL/hr to 94.5 kL/hr. All samples were collected from the flue gas stack according to Taiwan EPA NIEA A302.72C. To ensure contamination-free sampling and transportation, this research also took one blank trip and one blank field when conducting the field sampling. After completing the flue gas sampling, this work brought the samples back to the laboratory and placed them in a refrigerator at a temperature below 10°C.

Six ambient air samples were collected using a PS-1 sampler (Graseby Andersen, GA) according to the Taiwan EPA Reference Method NIEA A102.11A. Fig. 1 also shows the sampling site. The samples were collected separately in August and November. Three sampling sites A, B, and C were chosen as considering the upwind and downwind effect and available sampling location. The wind blew from the southwest 8% of the time on August and from the northeast around 30% of the time on November according to the wind rose. The sampling flow rate was specified at ~0.225 m<sup>3</sup>/min, and each sample was collected continuously on three consecutive days. The PS-1 sampler was equipped with a quartz fiber filter for sampling heavy metals. The filters were digested following the procedure of NIEA A301.11C, which is the EPA reference method in Taiwan, and metal elements were quantified by ICP-AES (Jobin-Yvon, JY38 Plus). Additionally, nonparametric correlations and the median test analysis for all samples were analyzed using Statistical Package for Social Sciences, version 15.0 (SPSS, Chicago, IL).



**Fig. 1.** The sampling sites were located in northern Taiwan.

## RESULTS AND DISCUSSION

### *Metal Concentrations in Flue Gas Stack of the Heavy Oil-Fueled Power Plant*

The current work classified metal elements into toxic (As, Cd, Cr, Hg, Ni and Pb), anthropogenic (Ba, Cu, Mn, Sb, Se, Sr, Ti, V and Zn) and crust (Al, Ca, Fe, K and Mg) groups (Wang *et al.*, 2003). Findings reveal higher mean concentrations in two crust elements of Al and K (1637 and 1153  $\mu\text{g}/\text{m}^3$ ), respectively, and two anthropogenic elements, Zn and Ba (1445 and 1128  $\mu\text{g}/\text{m}^3$ ). The four metals account for 82.2% of total metal element mass concentrations. Findings show Mercury (Hg) as having the lowest metal element mass concentrations of the twenty investigated, with a mean value of 0.15  $\mu\text{g}/\text{m}^3$ . The concentrations of toxic elements Cd, Cr, Ni and Pb, were higher than those in Reddy's study (Reddy *et al.*, 2005), however, As and Hg were lower than those in Reddy's study (Reddy *et al.*, 2005). The capacity in this study was 2,000MW, much higher than in Reddy's study (200 MW for a coal-fired power plant and 6 MW for a fuel-oil based power plant). Given a compatible power plant (1170 MW), the concentrations of As Cd, Cr, Hg, Ni and Pb were much lower than those in Aunela-Tapola's study (Aunela-Tapola *et al.*, 1998).

Table 1 lists metal concentrations in the flue gas stack of a power plant among different seasons, the first report among English literatures for a power plant fueled by heavy oil in Taiwan. Findings show most mean metal concentrations in fall as having the lowest concentration compared to those in other seasons except Fe, Cd, Cr, Mn, Ni, Sb, Se and Ti. Furthermore, findings show Cd and Hg as statistically significant with seasons. The result also indicates seasonal statistical significance for some operational parameters, flue gas temperature, flue gas velocity, fuel consumption, boiler steam temperature and electricity. Additionally, no significant differences exhibited in all variables among the four stacks via the nonparametric Kruskal-Wallis *H* tests (data not shown). These findings also qualified the analysis results.

**Table 1** Concentrations of metal in the flue gases of the power plant.

Elements ( $\mu\text{g}/\text{Nm}^3$ , dry)	Spring (n = 4)			Summer (n = 4)			Fall (n = 4)			Winter (n = 4)			$p^a$		
	Mean	SD	Range	Mean	SD	Range	Mean	SD	Range	Mean	SD	Range			
<i>Toxic metals</i>															
As	0.18	0.000	0.10-0.30	0.93	0.001	0.10-1.80	0.18	0.000	0.15	0.10-0.30	0.45	0.000	0.40	0.40-0.60	0.098
Cd	0.75	0.001	ND <sup>b</sup> -2.00	ND	ND	ND	5.68	0.005	3.85	1.40-13.6	1.58	0.002	1.35	ND-3.60	0.023*
Cr	10.1	0.012	3.00-27.9	7.43	0.005	1.50-13.3	3.93	0.001	3.65	2.60-5.80	1.90	0.001	2.00	0.50-3.10	0.113
Hg	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.60	0.001	0.25	ND-1.90	0.017*
Ni	32.2	0.027	13.0-72.3	10.4	0.009	2.0-22.4	12.3	0.008	12.4	4.80-19.6	15.8	0.012	11.4	7.10-33.2	0.195
Pb	7.76	0.004	3.00-13.0	7.20	0.005	2.20-12.9	4.93	0.001	4.65	3.80-6.60	5.23	0.003	5.35	2.30-7.90	0.599
Sum	51.1		25.9			27.0					25.5				
<i>Crust metals</i>															
Al	4776	8.351	465-17300	495	0.186	362-770	82.2	0.030	78.4	50.0-122	1193	1.355	996	40.4-2740	0.109
Ca	772	0.226	570-973	680	0.311	351-1061	262	0.130	256	142-395	1015	0.993	890	59.7-2220	0.207
Fe	420	0.205	245-710	241	0.219	122-569	243	0.169	169	141-494	193	0.133	190	53.4-337	0.300
K	1669	0.818	496-2400	1201	0.754	569-2290	106	0.015	104	92.5-124	1636	1.907	1343	17.0-3840	0.229
Mg	120	0.023	94.9-150	150	0.107	90.8-311	93.3	0.074	71.0	35.3-196	117	0.103	115	10.5-229	0.760
Sum	7757		2768			787					4154				
<i>Anthropogenic metals</i>															
Ba	1459	0.620	726-2220	810	0.806	5.20-1910	4.90	0.003	4.15	2.60-8.70	2236	2.687	1776	1.50-5390	0.210
Cu	5.38	0.002	2.80-6.90	5.63	0.006	0.80-15.1	1.25	0.002	0.65	0.10-3.60	2.28	0.002	1.95	0.40-4.80	0.086
Mn	35.8	0.010	28.1-50.7	26.1	0.010	17.0-39.2	32.3	0.009	34.8	20.0-39.8	18.5	0.017	15.8	1.70-40.7	0.446
Sb	0.71	0.000	0.30-1.20	0.28	0.000	0.10-0.50	0.38	0.000	0.15	0.10-1.10	0.35	0.000	0.35	0.30-0.40	0.207
Se	0.41	0.000	0.15-0.70	0.25	0.000	ND-0.80	0.45	0.000	0.40	0.20-0.80	0.06	0.000	0.07	ND-0.10	0.092
Str	7.25	0.003	4.39-10.6	5.93	0.002	4.10-8.50	1.28	0.001	1.25	0.60-2.0	6.60	0.008	4.50	0.30-17.1	0.135
Ti	1.38	0.000	0.80-1.70	1.53	0.000	0.80-1.90	2.90	0.002	1.85	1.50-6.40	3.33	0.003	3.20	0.60-6.30	0.390
V	12.7	0.009	5.97-25.4	6.08	0.004	2.30-9.70	5.60	0.006	3.65	1.30-13.8	8.6	0.010	4.85	1.90-22.7	0.388
Zn	1969	0.799	847-2630	1479	0.875	666-2660	87.0	0.047	82.6	35.0-148	2243	2.645	1812	10.4-5340	0.163
Sum	3492		2335			136					4519				
CO <sub>2</sub> (%)	12.4	0.33	12.0-12.8	12.5	0.31	12.1-12.8	12.5	0.10	12.4	12.4-12.6	12.6	0.14	12.7	12.4-12.7	0.503
O <sub>2</sub> (%)	5.48	0.38	5.00-5.90	5.35	0.34	5.00-5.80	5.10	0.34	4.95	4.90-5.60	5.28	0.38	5.20	4.90-5.80	0.377
FGT(°C)	152	5.29	147-158	137	13.5	127-157	130	4.53	128	127-137	126	2.84	125	124-130	0.021*
FGV(m/s)	21.0	3.62	16.6-25.3	12.1	6.01	8.53-21.1	13.5	1.35	13.4	12.1-15.3	14.9	1.28	15.2	13.0-15.9	0.038*
MT(%)	12.1	0.77	11.1-12.7	11.2	1.42	9.75-12.7	11.7	0.44	11.7	11.1-12.1	11.4	0.08	11.4	11.3-11.4	0.525
FC (kJ/h)	90.5	3.18	87.5-94.5	52.6	23.6	37.5-87.8	60.5	1.72	59.8	59.3-63.0	67.5	5.27	68.2	61.6-72.1	0.021*
BST (°C)	538	2.31	535-541	535	8.71	522-540	508	5.02	508	502-513	511	4.43	510	507-517	0.009*
BP(kg/cm <sup>2</sup> )	178	12.2	171-197	121	39.8	93.4-180	152	6.70	151	145-161	154	5.06	154	148-159	0.069
E (MWH)	399	3.33	396-404	226	115	164-398	266	7.89	267	255-275	299	12.5	301	284-312	0.026*

a: Kruskal-Wallis  $H$  tests, b: ND: not detectable,  $p < 0.05$ 


FGT: flue gas temperature, FGV: flue gas velocity, MT: moisture, FC: fuel consumption, BST: boiler steam temperature, BP: boiler pressure, E: electricity.

For toxic metals, Cr, Ni and Pb concentrations reached the highest in spring. However, Hg concentrations are only detected in winter and not detectable in other seasons. The sums of toxic metals and crust metals in spring were obviously much higher than those in other seasons. The sums of crust metals and anthropogenic metals were much lower in fall than those in other seasons. Dust storms always occur in the dry springtime (Fang *et al.*, 2009; Shen *et al.*, 2009), which will probably lead to more toxic metals from the power plant contributing to nearby ambient air. It warrants more concern in the future.

#### Metal Concentrations in Ambient Air around the Heavy Oil-Fueled Power Plant

As concentration ranged from 0.50–2.00 ng/m<sup>3</sup>, with an average of 1.30 ng/m<sup>3</sup>. Cd concentration ranged from 0.10–0.70 ng/m<sup>3</sup>, with an average of 0.45 ng/m<sup>3</sup>. Ni concentration ranged from 2.00–13.0 ng/m<sup>3</sup>, with an average of 5.35 ng/m<sup>3</sup>. Pb concentration ranged from 8.9–46.7 ng/m<sup>3</sup>, with an average of 26.2 ng/m<sup>3</sup>. Concentration was between Moreno's result (Moreno *et al.*, 2007) of PM<sub>10</sub> and PM<sub>2.5</sub> (a mean of 1.9 ng/m<sup>3</sup> and 1.0 ng/m<sup>3</sup>, respectively) in Puertollano, an important coal mining town including fossil fuel burning power plants as well as petrochemical and fertilizer complexes. Cd, Ni or Pb concentration was higher than that in Moreno's result (Moreno *et al.*, 2007) (Cd in PM<sub>10</sub> and PM<sub>2.5</sub> both were 0.1 ng/m<sup>3</sup>; Ni in PM<sub>10</sub> and PM<sub>2.5</sub> were 4.1 ng/m<sup>3</sup> and 3.0 ng/m<sup>3</sup>, respectively; Pb in PM<sub>10</sub> and PM<sub>2.5</sub> were 12 ng/m<sup>3</sup> and 9.3 ng/m<sup>3</sup>, respectively). However, Cd, Ni or Pb was lower than that in Yatkin's study (Yatkin and Bayram, 2007), whether in an urban or suburban sampling site (except for PM<sub>2.5</sub> sample in suburban winter). Lippmann *et al.* (2006) indicated Ni was a particularly influential component of ambient FPM (fine ambient particulate matter) in terms of cardiac responses to inhalation ambient air FPM. The Cd or Ni concentration did not exceed target values established by the European Council (2004/107/EC) for As (6 ng/m<sup>3</sup>), Cd (5 ng/m<sup>3</sup>) and Ni (20 ng/m<sup>3</sup>) (Petaloti *et al.*, 2006). The Pb concentration was also much lower than that (1.0 µg/m<sup>3</sup>) in the ambient air quality standard of the Taiwan EPA (2004/0060) (EPA). The Hg concentration was 0.10 ng/m<sup>3</sup> in all samples, the lowest concentration compared with other studies (Kim and Kim, 2001; Liu *et al.*, 2002; Sakata and Marumoto, 2002; Wang *et al.*, 2005).

Fig. 2a and 2b show the toxic metal profiles measured from ambient atmosphere of the heavy oil-fueled power plant in August and November, respectively. The Pb was the dominant metal among the six toxic metals for August and November at three sampling sites, except for November samples at the B site. Fig. 2c illustrates the averaged profiles of toxic metals in ambient atmosphere. The pattern was totally different from that in the power plant flue gas. The highest toxic metal concentration was Cd in flue gas samples; however, Pb was highest in ambient samples. The influence of metal concentrations from power plant flue gas to ambient atmosphere is seemingly not direct; other possible influential factors warrant further investigation.



**Fig. 2.** (a) Toxic metals profiles in the ambient atmosphere in August (b) Toxic metals profiles in the ambient atmosphere in November (c) Averaged toxic metals profiles in the ambient atmosphere

Al, Fe and Ca are the most common metal elements formed in the earth's crust (Taylor, 1964). The enrichment factor (EF) pattern is a reference value of contamination

status (Kim *et al.*, 2002). The EF provides information to judge enrichment (or depletion) of a given element relative to the reference element (Al, in this study). The Taylor's study defines EF as follows:  $EF = \{X/Al\}_{\text{sample}} / \{X/Al\}_{\text{crust}}$  (X denotes a metal of interest). The EF values were grouped into three categories in this study: (1) <10, Al, Ca, Fe, Mg, Mn, Sr and Ti, (2) 10–1000, As, Ba, Cr, Cu, K, Ni, Pb, Sb and V, (3) >1000, Cd, Se and Zn. Fig. 3 also shows the EF for the twenty analyzed metal elements for various studies.

For toxic metals, the values of EFs for As, Cd and Ni were highest in Reddy's study on the coal fired power plant (Reddy *et al.*, 2005). The 220 MWs power plant was located in western India with an independent bituminous coal generation unit and equipped with tangential burners. The EF for Cr was highest in a 6 MW-installed capacity oil-based power plant also located in western India (Reddy *et al.*, 2005). Lower EFs for the power plant in this study were probably due to higher combustion efficiency or higher APCD efficiency. Findings show the EF for Pb in Yatkin's study (Yatkin and Bayram, 2008) to be highest. The sampling site was in the city center near a motorway in Izmir, the third most populous city in Turkey. Heavy loading transportation obviously contributed more Pb to the atmosphere than other industries. Contrasted to toxic metals, the highest EFs found in this study were Al, Ba, K, Mg, Mn, Sr and Zn, perhaps because of the power plant's special geographic effect.

### Correlation of Metal Concentrations and Operational Parameters

This study applied nonparametric statistical analyses to evaluate the relationship between metal concentrations and operational parameters. The operational parameters included emitted CO<sub>2</sub>, O<sub>2</sub>, flue gas emission temperature,

flue gas velocity, moisture, heavy oil consumption rate, boiler steam temperature, boiler operational pressure, and electricity. The sampling size was not large enough to reach statistical significance; therefore, we chose the nonparametric Spearman correlation coefficient test. The results show no significant difference for all metal concentrations among four stack flue gases (data not shown). This finding indicates no obvious emission concentration variations for the mean annual values. Findings show significant Spearman's correlation coefficients among parts of the metals (Table 2), but As, Cd, Cr, Hg and Ti correlated less with other metals. The highest coefficient in nineteen metal pairs occurred between Ba and K, with  $r = 0.968$  and  $p < 0.001$ . All statistically related metals showed significant positive correlations, however, findings show only a negative correlation between As and Se ( $r = -0.564$ ,  $p = 0.023$ ). Seventeen pairs among sixty significant pairs have high coefficients with  $r > 0.800$  and  $p < 0.001$ , especially, the crust metals, Al, Ca and K related pairs, account for eleven pairs.

Statistically significant associations did not exhibit between metal concentrations and flue gas velocity. Significant positive correlations were found between Ni and fuel consumption ( $r = 0.753$ ,  $p = 0.001$ ), boiler pressure ( $r = 0.679$ ,  $p = 0.004$ ) and electricity ( $r = 0.662$ ,  $p = 0.005$ ). However, significant negative correlations were found between As and moisture ( $r = -0.606$ ,  $p = 0.013$ ), and boiler pressure ( $r = -0.625$ ,  $p = 0.01$ ). Ca, Cr, Cu, Mg, Mn, Ti, V and Zn showed no significant correlation with all operational parameters. Furthermore, the values of Cr and Pb were significantly related to decreasing CO<sub>2</sub> emission ( $r = -0.528$ ,  $p = 0.036$  and  $r = -0.553$ ,  $p = 0.026$ , respectively). Table 2 indicates negative correlations between most toxic metals (As, Cd, Cr and Hg) and




Fig. 3. The enrichment factor (EF) ratio for each metal element.



operational parameters, though some of the pairs were not statistically significant. Especially, for all operational parameters, Hg only negatively significantly correlated with flue gas temperature ( $r = -0.526$ ,  $p = 0.036$ ). Comparing the pairs of operational parameters, findings show significant positive correlations between electricity and flue gas velocity ( $r = 0.906$ ,  $p < 0.0001$ ), fuel consumption ( $r = 0.950$ ,  $p < 0.0001$ ) and boiler pressure ( $r = 0.909$ ,  $p < 0.0001$ ). This study provides only preliminary statistical results between metal concentrations and operational parameters because of small sample size. Further investigation requires larger sample sizes. Nevertheless, to our knowledge this is the first report to interpret correlations between metal concentrations and operational parameters.

## CONCLUSIONS

Comparing with other studies on power plants, metal emission concentration from the power plant was not the highest. However, comparing with local data in Taiwan, the value for anthropogenic element emission from the power plant (17,976 kg/yr) was much more significant (10.8, 104, and 4.81 fold higher than those emitted from the coal power plant, electrical arc furnace, and coke oven) compared with three reference emission sources (Wang et al., 2003). The value for crust element emission from the power plant (14,744 kg/yr) was not so significant (0.16, 7.16, and 0.24 fold higher than those emitted from the coal power plant, electrical arc furnace, and coke oven) compared with three reference emission sources (Wang et al., 2003). Metal concentration from the power plant was not so high; however, the total amount of metal emissions from power plants cannot be neglected owing to the high volume of flue gas. For proper environmental management of metals, especially toxic metals, establishing a complete source inventory of metal emission is necessary. The government should particularly pay more attention to power plants to address the information shortage.

## ACKNOWLEDGMENTS

We sincerely acknowledge Dr. Wen-Jhy Lee and Dr. Yong-Chien Ling for their insightful discussion and help in the sampling work and data analysis. Financial support by the Environmental Protection Bureau of Keelung City Government is gratefully acknowledged.

## REFERENCES

- Aunela-Tapola, L.A., Frandsen, F.J. and Hasanen, E.K. (1998). Trace Metal Emissions from the Estonian Oil Shale Fired Power plant. *Fuel Process. Technol.* 57: 1–24.
- EPA, <http://law.epa.gov.tw/en/laws/atmosph/air.html>.
- Fang, G.C., Wu, Y.S. and Chang, T.H. (2009). Comparison of Atmospheric Mercury (Hg) Among Korea, Japan, China and Taiwan during 2000–2008. *J. Hazard. Mater.* 162: 607–615.
- Gavett, S.H., Haykal-Coates, N., Copeland, L.B., Heinrich, J. and Gilmour, M.I. (2003). Metal Composition of Ambient PM<sub>2.5</sub> Influences Severity of Allergic Airways Disease in Mice. *Environ. Health Perspect.* 111: 1471–1477.
- Ghio, A.J. and Devlin, R.B. (2001). Inflammatory Lung Injury after Bronchial Instillation of Air Pollution Particles. *Am. J. Respir. Crit. Care Med.* 164: 704–708.
- Hatanpaa, E., Kajander, K., Laitinen, T., Piepponen, S. and Revitzer, H. (1997). A Study of Trace Element Behavior in Two Modern Coal-Fired Power Plants. I. Development and Optimization of Trace Element Analysis Using Reference Materials. *Fuel Process. Technol.* 51: 205–217.
- Heinrich, J., Hoelscher, B., Frye, C., Meyer, I., Wjst, M. and Wichmann, H.E. (2002). Trends in Prevalence of Atopic Diseases and Allergic Sensitization in Children in Eastern Germany. *Eur. Respir. J.* 19: 1040–1046.
- Jang, H.N., Seo, Y.C., Lee, J.H., Hwang, K.W., Yoo, J.I., Sok, C.H. and Kim, S.H. (2007). Formation of Fine Particles Enriched by V and Ni from Heavy Oil Combustion: Anthropogenic Sources and Drop-Tube Furnace Experiments. *Atmos. Environ.* 41: 1053–1063.
- Kim, K.H. and Kim, M.Y. (2001). The Temporal Distribution Characteristics of Total Gaseous Mercury at an Urban Monitoring Site in Seoul during 1999–2000. *Atmos. Environ.* 35: 4253–4263.
- Kim, K.H., Lee, J.H. and Jang, M.S. (2002). Metals in Airborne Particulate Matter from the First and Second Industrial Complex Area of Taejeon City, Korea. *Environ. Pollut.* 118: 41–51.
- Kuo, Y.M., Hung, H.F. and Yang, T.T. (2007). Chemical Compositions of PM<sub>2.5</sub> in Residential Homes of Southern Taiwan. *Aerosol Air Qual. Res.* 7: 403–416.
- Lippmann, M., Ito, K., Jing-Shiang, H., Polina, M. and Chen, L.C. (2006). Cardiovascular Effects of Nickel in Ambient Air. *Environ. Health Perspect.* 114: 1662–1669.
- Liu, S., Nadim, F., Perkins, C., Carley, R.J., Hoag, G.E., Lin, Y. and Chen, L. (2002). Atmospheric Mercury Monitoring Survey in Beijing, China. *Chemosphere.* 48: 97–107.
- Magari, S.R., Schwartz, J., Williams, P.L., Hauser, R., Smith, T.J. and Christiani, D.C. (2002). The Association of Particulate Air Metal Concentrations with Heart Rate Variability. *Environ. Health Perspect.* 110: 875–880.
- Moreno, T., Alastuey, A., Querol, X., Font, O. and Gibbons, W. (2007). The Identification of Metallic Elements in Airborne Particulate Matter Derived from Fossil Fuels at Puertollano. *Int. J. Coal Geol.* 71: 122–128.
- Petaloti, C., Triantafyllou, A., Kouimtzi, T. and Samara, C. (2006). Trace Elements in Atmospheric Particulate Matter over a Coal Burning Power Production Area of Western Macedonia, Greece. *Chemosphere.* 65: 2233–2243.
- Pope Iii, C.A., Burnett, R.T., Thun, M.J., Calle, E.E., Krewski, D., Ito, K. and Thurston, G.D. (2002). Lung Cancer, Cardiopulmonary Mortality, and Long-term Exposure to Fine Particulate Air Pollution. *J. Am. Med.*

- Assoc. 287: 1132–1141.
- Reddy, M.S., Basha, S., Joshi, H.V. and Jha, B. (2005). Evaluation of the Emission Characteristics of Trace Metals from Coal and Fuel Oil Fired Power Plants and Their Fate during Combustion. *J. Hazard. Mater.* 123: 242–249.
- Sakata, M. and Marumoto, K. (2002). Formation of Atmospheric Particulate Mercury in the Tokyo Metropolitan Area. *Atmos. Environ.* 36: 239–246.
- Sarkar, B. (2002). *Heavy Metals in the Environment*. Marcel Dekker, Inc., New York.
- Saxena, D.K., Singh, S. and Srivastava, K. (2008). Atmospheric Heavy Metal Deposition in Garhwal Hill Area (India): Estimation Based on Native Moss Analysis. *Aerosol Air Qual. Res.* 8: 94–111.
- Schaumann, F., Borm, P.J., Herbrich, A., Knoch, J., Pitz, M., Schins, R.P., Luettig, B., Hohlfeld, J.M., Heinrich, J. and Krug, N. (2004). Metal-Rich Ambient Particles (Particulate Matter 2.5) Cause Airway Inflammation in Healthy Subjects. *Am. J. Respir. Crit. Care Med.* 170: 898–903.
- Shen, Z.X., Cao, J.J., Arimoto, R., Han, Z.W., Zhang, R.J., Han, Y.M., Liu, S.X., Okuda, T., Nakao, S. and Tanaka, S. (2009). Ionic Composition of TSP and PM<sub>2.5</sub> during Dust Storms and Air Pollution Episodes at Xi'an, China. *Atmos. Environ.* 43: 2911–2918.
- Taylor, S.R. (1964). Abundance of Chemical Elements in the Continental Crust: A New Table. *Geochim. Cosmochim. Acta.* 28: 1273–1285.
- Wang, S., Feng, X., Qiu, G., Wei, Z. and Xiao, T. (2005). Mercury Emission to Atmosphere from Lanmuchang Hg-Tl Mining Area, Southwestern Guizhou, China. *Atmos. Environ.* 39: 7459–7473.
- Wang, W.C., Chen, K.S., Chen, S.J., Lin, C.C., Tsai, J.H., Lai, C.H. and Wang, S.K. (2008). Characteristics and Receptor Modeling of Atmospheric PM<sub>2.5</sub> at Urban and Rural Sites in Pingtung, Taiwan. *Aerosol Air Qual. Res.* 8: 112–129.
- Wang, Y.F., Huang, K.L., Li, C.T., Mi, H.H., Luo, J.H. and Tsai, P.J. (2003). Emissions of Fuel Metals Content from a Diesel Vehicle Engine. *Atmos. Environ.* 37: 4637–4643.
- Yatkin, S. and Bayram, A. (2007). Elemental Composition and Sources of Particulate Matter in the Ambient Air of a Metropolitan City. *Atmos. Res.* 85: 126–139.
- Yatkin, S. and Bayram, A. (2008). Determination of Major Natural and Anthropogenic Source Profiles for Particulate Matter and Trace Elements in Izmir, Turkey. *Chemosphere* 71: 685–696.

Received for review, September 14, 2009  
Accepted, November 18, 2009