



The Effects of Height on the Accumulation of *n*-Alkanes and Polycyclic Aromatic Hydrocarbons (PAHs) in Air-Conditioning Filter Dust from High-Rise Apartments

Ji-Zhong Wang^{1*}, Cheng Qian¹, Jia-Qin Liu², Xiu-Feng Ni¹, Shu-Chuan Peng^{1*}

¹ School of Resources and Environmental Engineering, Hefei University of Technology, Hefei 230009, China

² Institute of Industry & Equipment Technology, Hefei University of Technology, Hefei 230009, China

ABSTRACT

To assess the effect of floor levels of high-rise apartment buildings on the accumulation of contaminants in indoor environment, residential air-conditioner filter dust (ACFD) samples from the 1st, 10th, 20th and 30th floors of a high-rise apartment building were collected for the determination of *n*-alkanes and polycyclic aromatic hydrocarbons (PAHs). The results show that both *n*-alkanes and PAHs in the residential ACFD were ubiquitous but varied greatly in concentrations. The total concentrations of 27 *n*-alkanes ($\Sigma_{27}\text{AK}$) and 16 PAHs ($\Sigma_{16}\text{PAH}$) ranged from 1.35 to 9290 $\mu\text{g g}^{-1}$ and 278–34200 ng g^{-1} , respectively. Source apportionment revealed that *n*-alkanes were from mixed sources combining fossil fuel combustion, natural emission and solid biomass burning, but PAHs were mainly from indoor sources. Furthermore, the diagnostic ratio of paired low weight molecular PAH species may change during transportation and accumulation. Significantly higher concentrations of $\Sigma_{27}\text{AK}$ and $\Sigma_{16}\text{PAH}$ were observed in the samples from low floor levels (the 1st and 10th floors) compared to those from high floor levels (the 20th and 30th floors). The results of classification and regression tree analysis clearly suggested floor level is the most important factor influencing the accumulation of $\Sigma_{27}\text{AK}$ and $\Sigma_{16}\text{PAH}$ in the ACFD. Our findings imply that people living on lower floor levels have greater exposure risks to PAHs associated with indoor dust.

Keywords: High-rise apartment; Floor level; Air-conditioner filter dust; *n*-alkanes; PAHs.

INTRODUCTION

Exposure to indoor air contaminants is a vital health risk for urban populations because they generally spend more than 80% of time indoors (Mirhoseini *et al.*, 2016). However, with rapid urbanization, people living in countryside tend to migrate to the cities for better work opportunities and livings. It is reported more than 50% of the world's population resided in urban areas in 2013, compared to only 40% in 1980 (The World Bank, 2016). The United Nations Department of Economic and Social Affairs (DESA)'s Population Division also noted that there were 28 mega-cities worldwide in 2014, home to approximate 12% of global urban populations (United Nations Department of Economic and Social Affairs, 2014). As a result, the increase of urban population has a direct influence on the types of dwellings that people live in. High-rise apartment buildings are common in metropolitan regions worldwide, especially in China. Previous studies have demonstrated that the height of the dwelling in urban

areas is the key factor influencing the wind speed (Murakami *et al.*, 1979; Lu and Ip, 2009), microenvironment (Lee and Jo, 2006; Gao *et al.*, 2008) and inhabitant's health (Drennan *et al.*, 2016). More importantly, floor levels can influence indoor-outdoor exchange of contaminants, resulting in significant vertical variation of indoor and outdoor ambient pollution (Jo and Kim, 2002; Jo and Lee, 2006). The exposure to indoor airborne pollutants for the residents living in different floor levels may vary accordingly. Thus the information on how the floor level affects the indoor air quality lays groundwork for a comprehensive assessment of inhalation exposure risks for urban populations living in high-rise buildings. However, such information has not been well addressed to date.

The *n*-alkanes and polycyclic aromatic hydrocarbons (PAHs) are two groups of ubiquitous indoor environmental pollutants, both of which are commonly used to indicate the pollution significance because of their widespread sources including indoor and outdoor emissions (Huang *et al.*, 2015; Jung *et al.*, 2015; Mwangi *et al.*, 2015; Wang *et al.*, 2015). Furthermore, PAHs have obtained extensive attention because of their persistency and carcinogenicity nature (Zhang *et al.*, 2015). Thus numerous studies applied PAHs as model pollutants to assess the health risks (Bhargava *et al.*, 2004; Zhou and Zhao, 2012; Kavouras *et*

* Corresponding author.

E-mail address: wangjizh@hfut.edu.cn (J.Z. Wang); pengshuchuan@hfut.edu.cn (S.C. Peng)

et al., 2015; Lee *et al.*, 2016; Mohammed *et al.*, 2016). Due to the importance of indoor exposure, PAHs in indoor air (including gas phase and particle) and floor dust are commonly investigated to elucidate potential health risks (Essumang *et al.*, 2015; Tiwari *et al.*, 2015). However, most of these sampling strategies demand specialized equipment and are time-consuming and noise nuisance. Furthermore, there are many uncertainties with the results of health assessment based upon these monitoring data due to temporal and spatial variations of contaminant concentrations. For example, significantly higher concentrations of PAHs in both gaseous and particle phases were observed in winter season than in summer (Masih *et al.*, 2012). Muenhor and Harrad (2012) also demonstrated that floor dust from the same room and building varied greatly in the concentrations of polybrominated diphenyl ethers. Dust accumulated on the air condition filter (ACFD) may be a better candidate to elucidate indoor exposure risks because it can integrate contamination during a given time interval (Ni *et al.*, 2011; Chou *et al.*, 2016).

The main objectives of the present study were to investigate *n*-alkanes and PAHs in ACFD from a high-rise apartment building, and further to evaluate the potential exposure risk to indoor PAHs for the dwellers. Special attention was paid to the effect of floor levels on the accumulation of hydrocarbon compounds and associated human health risk.

MATERIALS AND METHODS

Sampling Location and Sampling

The City of Hefei is a capital city and the political, economic, and cultural center of Anhui Province in Eastern China. It is located in the middle altitude zone (30°57′–32°32′ North latitude and 116°41′–117°58′ East longitude) with an annual precipitation of ~1,000 mm. The city has a population of 7,790,000 inhabitants with over 70% of them living in urban areas. Undoubtedly, rapid population growth and urbanization has resulted in a concomitant increase on the high-rise apartment buildings in urban areas.

ACFD samples were collected from a new high-rise apartment building in Hefei. The apartment has 30 floors (about 90 m in height) and was built in 2006. Additional five buildings with the same height and floor levels are around the study building with the space between each other approximately 20 m. A total of 44 ACFD samples were collected in August 2014. Among these samples, seven of them (two in bedrooms and five in living rooms) were collected from seven homes on the 1st floor; thirteen samples (seven in bedrooms and six in living rooms) from nine homes on the 10th floor; eleven samples (eight in bedrooms and three in living rooms) from five homes on the 20th floor; thirteen samples (ten in bedrooms and three in living rooms) from five homes on the 30th floor. Less than 0.2 g to 1.3 g of dust particles on impervious surface of each air-conditioner filter were collected using plastic brush and dust pans by gentle sweeping motion. Before each sampling, brush and dust pans were cleaned with acetone and paper towels. All samples were wrapped with

solvent-rinsed aluminum foil and stored in polyethylene zip bags. After transported to the analytical laboratory, dust samples were stored at –20°C prior to chemical analysis.

A questionnaire survey was conducted to understand the potential indoor sources of hydrocarbons. All participating families returned with valid questionnaire. Most participants had ages ranging from 20 to 50 years old. The rooms had been interiorly decorated before the participating families moved in. There is no ventilation and central air-conditioner system in the apartment, and only independent air conditionings were used during summer seasons. All but one family had significant cooking activities (generally 2–3 times d⁻¹) and only four families were impacted by indoor smoking. Six filters of air-conditioners in three families had been cleaned before the summer.

Chemical and Reagents

n-hexane, dichloromethane, acetone and methanol supplied by Oceanpak Alexative Chemical Co., Ltd (Gothenburg, Sweden) were of pesticide grade. Neutral silica gel (80–100 mesh) was purchased from Guangzhou Xinshi Chemistry Experimental Equipment (Guangzhou, China) and was pre-cleaned by Soxhlet extraction for 48 hours using methanol and dichloromethane, respectively, prior to use. Sodium sulfate obtained from Sigma-Aldrich (St Louis, MO, USA) was baked at 450°C for 4 hours and stored in sealed containers. All glassware was cleaned with chromic acid and tap water, followed by rinsing with deionized water, and then baked at 450°C for 4 hours.

A mixture standard of *n*-alkanes including *n*C₁₀, *n*C₁₂, *n*C₁₄, *n*C₁₆, *n*C₁₈, *n*C₂₀, *n*C₂₂, *n*C₂₄, *n*C₂₆, *n*C₂₈, *n*C₃₀, *n*C₃₂, *n*C₃₄, *n*C₃₆ and *n*C₃₈ was purchased from AccuStandard (New Haven, CT, USA). The standard solutions of *n*C₂₄-*d*₅₀ and *n*C₃₀-*d*₆₂ at 1000 µg mL⁻¹ each were purchased from Cambridge Isotope Laboratory (Andover, MA, USA) as surrogate and internal standards of *n*-alkanes to monitor the performance of sample extraction and cleanup, respectively. A mixture solution containing sixteen PAHs identified as priority pollutants by the United States Environmental Protection Agency (USEPA), i.e., naphthalene (Nap), acenaphthylene (Ace), acenaphthene (Ac), fluorene (Flu), phenanthrene (Phe), anthracene (Ant), fluoranthene (Flu), pyrene (Pyr), benzo[a]anthracene (BaA), chrysene (Chr), benzo[b]fluoranthene (BbF), benzo[k]fluoranthene (BkF), benzo[a]pyrene (BaP), indeno[1,2,3-cd]pyrene (IcdP), dibenzo[a,h]anthracene (DahA), and benzo[g,h,i]perylene (BghiP), was purchased from AccuStandard. A deuterated standard mixture solution containing five PAHs (Nap-*d*₈, Ace-*d*₁₀, Phe-*d*₁₀, Chr-*d*₁₂, and perylene-*d*₁₂) at 100 µg mL⁻¹ each was purchased from Cambridge Isotope Laboratory (Massachusetts, USA) and applied as surrogate standards of PAHs. Reference standards of 2-fluoro-1,1-biphenyl, *p*-terphenyl-*d*₁₄, and DahA-*d*₁₄ at 100 µg mL⁻¹ each (Cambridge Isotope Laboratory, USA) were used as internal standards.

Extraction and Analysis of PAHs

The procedures of extraction and purification of hydrocarbon compounds were described elsewhere (Chuang *et al.*, 1995). Briefly, the dust samples were spiked with a

known amount of $nC_{24-d_{50}}$ and five deuterated PAHs as surrogate standards and then extracted with 15 mL of a mixture of dichloromethane and hexane (50:50 in volume) in an ultrasonic bath for 30 min. The extraction was repeated three times. The combined extract was filtered through a 7 μm glass fiber filter (Whatman International, Maidstone, UK) and then concentrated to 1 mL. The condensed extract was further solvent exchanged to hexane by adding 10 mL of hexane and then reduced to 1 mL again. The sample was purified by passing through a glass column packed with 3 g of silica gel. After the column was conditioned with 5 mL of hexane and loaded with the sample, the first fraction containing n -alkanes was eluted using 12 mL of hexane, and the second fraction containing PAHs was eluted with 15 mL of a mixture of hexane and dichloromethane (7:3 in volume). Both fractions were condensed to 500 μL and spiked with known amounts of internal standards.

Both n -alkanes and PAHs were instrumentally analyzed using an Agilent 7890A gas chromatograph coupled with a 5975C mass spectrometer (Agilent Technologies, Palo Alto, California, US) in electron impact (EI) ionization mode. A 60 m \times 0.25 mm \times 0.25 μm DB-5 capillary column (Agilent Technologies) was employed for chromatographic separation. Ultra high purity helium (99.999%) was used as carrier gas with a constant flow rate of 1 mL min^{-1} . Sample (1 μL) was injected using an automatic sampling system AOC-20 Series in the splitless mode, and the temperature of injector was held at 280°C. The temperatures of ion source and quadrupole were both set at 150°C, and the transfer line temperature was at 280°C. For n -alkanes, the oven temperature was programmed from 60 (held 2 min) to 180°C at a rate of 8 $^{\circ}\text{C min}^{-1}$, to 220 $^{\circ}\text{C}$ at a rate of 2 $^{\circ}\text{C min}^{-1}$, and then to 320°C at a rate of 8 $^{\circ}\text{C min}^{-1}$. For PAHs analysis, the GC oven was temperature-programmed from 60°C to 200°C at a rate of 10 $^{\circ}\text{C min}^{-1}$, and to 250°C at a rate of 2 $^{\circ}\text{C min}^{-1}$, then to 290°C at a rate of 20 $^{\circ}\text{C min}^{-1}$, finally held isothermally for 5 min. The quantification and confirmation of each analyte was achieved via selected ion monitoring (SIM) mode. All n -alkanes with even numbers of carbon atoms were directly quantified by the calibration curves built using their corresponding reference standards at eight calibration concentrations (0.05, 0.1, 0.2, 0.5, 1, 2, 5, and 10 $\mu\text{g mL}^{-1}$). Each calibration solution contained 1.0 $\mu\text{g mL}^{-1}$ of internal standards. Odd number carbon n -alkanes were qualified with the average response factor of two adjacent even-numbered carbon components, given that their reference standards were not available (Wang *et al.*, 2012).

Quality Assurance and Quality Control (QA/QC)

Quantification was performed using an internal calibration method based on eight-point calibration curves with regression coefficient (R^2) value higher than 0.99 for all target compounds. For every ten samples, a calibration solution containing 1.0 $\mu\text{g mL}^{-1}$ each of n -alkanes or 0.5 $\mu\text{g mL}^{-1}$ each of PAHs was analyzed to monitor the instrument performance. For each batch of 15 dust samples, one solvent blank was processed to monitor laboratory contamination. Solvent blanks contained n -alkanes and PAHs less than 10% of the mean concentrations from all

dust samples. The reporting limits (RLs) were calculated at 0.078 $\mu\text{g g}^{-1}$ for all n -alkanes and 7.76 ng g^{-1} for all PAHs after including the lowest concentrations of the calibration curves and average sample weight into consideration (Wang *et al.*, 2007). The average recovery efficiency of $nC_{24-d_{50}}$ was 100.6 \pm 25.9%, and five deuterated PAHs had the recoveries of 88.8 \pm 23.3% for Nap- d_8 , 87.2 \pm 21.1 for Ace- d_{10} , 88.8 \pm 21.8 for Phe- d_{10} , 104.4 \pm 20.4 for Chr- d_{12} , and 107.0 \pm 16.0 for perylene- d_{12} .

Data Analysis

Because nC_{37} and nC_{38} were not detected in any sample, 27 n -alkane homologues (from nC_{10} to nC_{36}) were discussed in the present study and their total concentration was referred to as $\Sigma_{27}\text{AK}$. The total concentration of 16 PAH compounds was defined as $\Sigma_{16}\text{PAH}$. For any measured concentration below RL, zero was used for concentration calculation. Data analyses were conducted using Excel 2007 (Microsoft Corporation, Silicon Valley, USA) and the statistical analyses including normality test and principal component analysis were performed using SPSS 16.0 (Chicago, IL, USA).

To assess the potential sources of n -alkanes in ACFD, carbon preference index (CPI) and plant wax alkanes (WaxCn(%)) were calculated according to the following equations:

$$\text{CPI}_1 = \frac{\sum \text{odd}(nC_{10} - nC_{36})}{\sum \text{envn}(nC_{10} - nC_{36})} \quad (1)$$

$$\text{CPI}_2 = \frac{nC_{27} + nC_{29} + nC_{31} + nC_{33}}{nC_{28} + nC_{30} + nC_{32} + nC_{34}} \quad (2)$$

$$\text{WaxC}_n(\%) = \frac{\sum \left[n-C_n - \frac{1}{2}(n-C_{n-1} - n-C_{n+1}) \right]}{\sum (n-C_{15} - n-C_{34})} \quad (3)$$

It should be noted that CPI_1 was usually used to discriminate between petrogenic and biogenic sources of total n -alkanes; in general, CPI_1 close to 1 suggests incomplete combustion of fossil fuel, and it indicates contemporary biological sources when CPI_1 is greater than 3 (Simoneit, 1984). CPI_2 is utilized to distinguish the fraction of n -alkanes from high plant wax, a high value usually indicative of high plant wax input (Reddy *et al.*, 2000). WaxCn(%) is close to zero for petroleum or crude oil residue, whereas it approaches 100 for high terrestrial plants or marine plants (Gogou *et al.*, 1996).

In the present study, the classification and regression tree analysis was employed to uncover the relationships and interactions between PAHs or n -alkanes contents in ACFD and floor levels or room type (bedroom or living room) variables (JMP software, Version 10.0, SAS institute, Cary, NC, USA). As a nonparametric modeling method, the classification and regression tree analysis can justify the relationship between variables from a set of independent variables and recursively separate the data to obtain

increasingly homogeneous subsets based upon the independent variable splitting criteria using variance minimizing algorithms (Zheng *et al.*, 2009).

To better parameterize ACFD carcinogenicity related to the whole PAH fraction, the concentrations of PAHs in ACFD were further transferred to BaP equivalent concentration (BaP_{eq}) using individual toxic equivalency factor (TEF) (Pongpiachan, 2015). TEF values are 0.001, 0.001, 0.001, 0.001, 0.001, 0.01, 0.001, 0.001, 0.1, 0.01, 0.1, 0.1, 1, 0.1, 1, and 0.01 for Nap, Ace, Ac, Fl, Phe, Ant, Flu, Pyr, BaA, Chr, BbF, BkF, BaP, IcdP, DahA and BghiP, respectively (Yang *et al.*, 2015).

RESULTS AND DISCUSSION

n-Alkanes in Residential ACFD

The total concentrations of $\Sigma_{27}AK$ in residential ACFD ranged from 1.35 to 9290 $\mu\text{g g}^{-1}\text{dw}$. Because logarithm-transformed the concentration of $\Sigma_{27}AK$ followed normal distribution ($p = 0.244$, with One - Sample Kolmogorov - Smirnov Test in SPSS), the geometric mean was reported at 321 $\mu\text{g g}^{-1}\text{dw}$. Despite all *n*-alkane homologues between nC_{10} to nC_{36} were detectable in each residential ACFD, the predominated congeners belong to nC_{24} – nC_{33} homologues, and the carbon maximum number (C_{max} , defined as the homologue with the highest concentration among all *n*-alkanes) was observed at nC_{27} , nC_{29} or nC_{31} in 6, 23 or 8 out of 44 samples, respectively. Interestingly, samples from the 10th floor generally contained C_{max} at nC_{27} , while the C_{max} was mainly at nC_{29} for samples from the 20th and 30th floors. The C_{max} in samples from the 1st floor was complex, i.e., one sample at nC_{27} , one at nC_{29} , three at nC_{31} and two at nC_{33} . These suggest that *n*-alkanes in most samples were

partly contributed by biological origins (Eglinton and Hamilton, 1967; Bi *et al.*, 2005b) or combustion of biomass (bin Abas *et al.*, 1995; Wang *et al.*, 2009). The left part of Fig. 1 illustrates the patterns of *n*-alkanes as a function of carbon number. The results show unimodal distribution without significant odd-carbon number preference. CPI_1 and CPI_2 values varied between 0.72–2.8 and 0.98–3.2 with arithmetic means of 1.4 ± 0.38 and 1.7 ± 0.44 , respectively, suggesting *n*-alkanes in residential ACFD were predominately from anthropogenic activities combining biological materials. Human activities related sources of *n*-alkanes contain CPI values close to unit, whereas natural sources from vegetation generally exhibit high CPI values (> 3.0) (Bi *et al.*, 2005a; Wu *et al.*, 2007). Significantly higher CPI_2 values than CPI_1 suggest that long-chain alkanes in residential ACFD were more dominated by biogenic sources when compared with short-chain homologues. $WaxC_n(\%)$ value was determined in a range of 4.4%–24.9% with an arithmetic mean of $13.2 \pm 4.3\%$, indicating a predominant contribution from anthropogenic sources. A similar contribution of *n*-alkanes in urban aerosol from biogenic origins was also investigated (Bi *et al.*, 2003; Lin and Lee, 2004; Feng *et al.*, 2006; Wu *et al.*, 2006). $WaxC_n(\%)$ significantly correlated against CPI_1 (Fig. 1(b)). Therefore, CPI_1 could be expected as a useful indicator for the prediction of biogenic sources derived *n*-alkanes.

PCA analysis shows three main factors which contributed the majority of the variance (87.0%). Factor 1 (accounting for 46.5% of the total variance) was mostly associated with short-chain alkanes (from nC_{10} to nC_{23}), suggesting fossil fuel combustion (Cincinelli *et al.*, 2007; Alves *et al.*, 2016). Factor 2 contributing to 32.7% of total variance was mostly explained by high loadings of long-chain *n*-alkanes,




Fig. 1. The profile of *n*-alkanes in the air-conditioner filter dust (ACFD) from a high-rise apartment building (a), and the correlation between carbon preference index (CPI_1) and plant wax alkanes ($WaxC_n(\%)$) (b). It is noted that CPI_1 and $WaxC_n(\%)$ were calculated based upon Eqs. (1) and (3), respectively.

indicating natural sources because long-chain alkanes are largely emitted from plant wax (Eglinton and Hamilton, 1967; Bi *et al.*, 2005b). The third factor (factor 3) could explain 7.8% of the total variance, and was dominated by nC_{21} – nC_{26} . Previous studies demonstrated that solid fuel and biomass burning would produce moderate n -chain alkanes (Oros and Simoneit, 2001; Bi *et al.*, 2008; Sun, 2012). Therefore, residual n -alkanes in residential ACFD were from mixed sources combining fossil fuel combustion, natural emission and solid biomass burning.

Occurrence of PAHs in Residential ACFD

Nap, Fl, Phe, Flu, Pyr, and Chr were detected in all ACFD samples with the concentration ranges of 14.2–2240, 10.0–376, 56.0–3290, 51.7–5030, 36.8–4090, and 36.4–3610 $ng\ g^{-1}$, respectively (Fig. 2). Only 15 samples contained Ace greater than the report limit, and Ace was observed with the lowest mean concentration ($5.1 \pm 37.4\ ng\ g^{-1}$) among all target PAHs. The detection frequencies of Ac, Ant, BaA, B(b+k)F, BaP, IcdP, DahA and BghiP were 63.7%, 93.1%, 84.1%, 97.7%, 76.3%, 93.2%, 50.0% and 90.9%, respectively. Therefore, the residential ACFD was dominated by high molecular weight PAHs (from 4 to 6 rings), accounting for $68.6 \pm 16.8\%$ of the total concentration of $\Sigma_{16}PAH$ in the present study. Low molecular weight PAHs with high volatility appeared to less accumulate in the ACFD (Zhou *et al.*, 2010). Because high molecular weight PAHs are more toxic and persistent in environment, TEF-adjusted concentrations of high molecular weight PAHs were 2–3 orders of magnitude higher than that of low molecular weight PAHs. The total concentration of $\Sigma_{16}PAH$ varied from 278 to 34200 $ng\ g^{-1}$ and appeared to have a lognormal distribution ($p = 0.84$, with One - Sample Kolmogorov - Smirnov Test in SPSS). The geometric mean of $\Sigma_{16}PAHs$ in the residential ACFD was determined at 2280 $ng\ g^{-1}$ and BaP_{eq} concentrations ranged from 2.27 to 5290 $ng\ g^{-1}$.

Zhou *et al.* (2010) reported the total concentrations of PAHs in ACFD of different indoor environments, i.e., 4895 $ng\ g^{-1}$ for household, 4231 $ng\ g^{-1}$ for office, 475 $ng\ g^{-1}$ for meeting room, 740 $ng\ g^{-1}$ for chemical laboratory, and 13090 $ng\ g^{-1}$ for restaurant. ACFD samples from the commercial office, secondary school, shopping mall, hospital, electronic factory and manufacturing plant were also found with the total concentration of PAHs at 5850 ± 3190 , 5610 ± 2330 , 5530 ± 2070 , 3720 ± 1820 , 4090 ± 2380 and $22200 \pm 6050\ ng\ g^{-1}$, respectively (Kang *et al.*, 2010). The concentrations of PAHs in ACFD were comparable to or slightly lower than those in door floor dust (Gevao *et al.*, 2007; Mannino and Orecchio, 2008; Peng *et al.*, 2012; Qi *et al.*, 2014). Therefore, our results, along with previous studies, indicate that dust retained on the AC filter is suitable for the evaluation of health risks from indoor exposure to indoor contaminants because they can reflect both the indoor dust and indoor air quality through air recirculation.

Paired isomers based diagnostic ratios of Ant/(Ant + Phe), Flu/(Flu + Pyr), BaA/(BaA + Chr) and IcdP/(IcdP + BghiP) were 0.07 ± 0.02 , 0.58 ± 0.05 , 0.16 ± 0.08 and 0.64 ± 0.11 , respectively. High ratio values for Flu/(Flu + Pyr) and IcdP/(IcdP + BghiP) indicate incomplete combustion of fossil fuels and biomasses (Yan *et al.*, 2005). Cooking would also be expected as an important source of residual PAHs in ACFD because of the high value of BaP/BghiP (2.26 ± 3.37) (Zhang *et al.*, 2009). However, low ratios of Ant/(Ant + Phe) and BaA/(BaA + Chr) suggest petroleum-related sources (Mwangi *et al.*, 2015). Therefore, the identification of the sources of residual PAHs in ACFD remained difficult, possibly due to the change in the diagnostic ratios during the transportation and accumulation. Previous studies indicated that subtle variation in the physicochemical properties of the paired PAH species would alter the ratios from sources to receptors (Schauer *et al.*, 1996; Zhang *et al.*, 2005). For example, Ant was more susceptible to degradation than Phe, resulting in a reduced




Fig. 2. The concentration distribution and detection frequency of individual PAHs in air-conditioner filter dust (ACFD) from a high-rise apartment building.

Ant/(Ant + Phe) ratio in pine needles (Hwang *et al.*, 2003). In the present study, very low ratios of Ant/(Ant + Phe) observed in ACFD would be explained as a result of the preferential decay or volatilization of Ant in the aged ACFD. However, ratios of high weight molecular PAH species, e.g., IcdP/(IcdP + BighP), were stable during the transportation (Wang *et al.*, 2011). High IcdP/(IcdP + BighP) ratios in the present study suggest sources from biomass produced by cigarettes burning (Du *et al.*, 2006) and cooking (Ye *et al.*, 2013). Therefore, PAHs accumulated in residential ACFD would be mainly from indoor sources.

The Effect of Floor Level on the Accumulation of PAHs and *n*-Alkanes in ACFD

The results of classification and regression tree analysis indicate that greater concentrations of $\Sigma_{27}\text{AK}$ and $\Sigma_{16}\text{PAH}$ were determined in the samples collected from lower floor levels (1st and 10th floor) compared with higher floor levels (20th and 30th floor) (Figs. 4(a) and 4(b)). The mean concentrations of $\Sigma_{27}\text{AK}$ and $\Sigma_{16}\text{PAH}$ in ACFD from lower floor levels were 4.6 and 5.2 times greater than the concentrations from higher floor levels. Samples from the 1st and 10th floors contained comparable $\Sigma_{27}\text{AK}$ concentrations ($p = 0.09$), and the concentrations in the dust from the 20th and 30th floors were also comparable ($p = 0.28$). Room type (bedroom and living room) did not significantly influence *n*-alkanes accumulation in the ACFD, but influenced PAHs accumulation. For example, greater concentrations of PAHs were found in ACFD from bedrooms versus living rooms located on the 10th floor ($p < 0.01$).

Obviously, the floor level is a critical factor influencing the accumulation of hydrocarbon contaminants in residential ACFD. Residents living on lower floor levels of high-rise

buildings are exposed to elevated concentrations of particle-associated PAHs. Previous studies demonstrated that both outdoor and indoor airborne contaminants such as carbon monoxide, fine particles, volatile organic compounds (VOCs) and PAHs were significantly higher at low floor levels than at high floor levels (Jo *et al.*, 2003; Jo and Lee, 2006; Jung *et al.*, 2011). Therefore, our findings strongly supported the conclusion on a decreasing trend in indoor nonvolatile PAHs concentration with increasing floor levels. Additionally, concentrations of $\Sigma_{16}\text{PAH}$ were significantly correlated with $\Sigma_{27}\text{AK}$ ($p \leq 0.01$) in the samples from low floor levels (1st and 10th floors), but they were not related in the ACFD collected from high floor levels (Fig. 3). These results imply that PAHs and *n*-alkanes had similar sources and accumulative processes in ACFD. A number of studies attributed some indoor air pollutants mainly to the input from outdoor anthropogenic sources (Cheruyot *et al.*, 2015). However, we were unable to determine the relative importance of outdoor and indoor sources in their contributions to *n*-alkanes and PAHs concentrations in residential ACFD, given that, PAHs in outdoor aerosol were not determined in the present study. It should be noted that the samples from the 20th and 30th floors generally contained C_{max} at nC_{29} , but those with C_{max} at nC_{31} were mostly from the 1st and 10th floors. The nC_{31} is usually considered as a chemical marker of grass plants, whereas nC_{29} is mainly from wax plants (Oros *et al.*, 2006). Thus lower and higher floors of the apartment building are more affected by grass and wax plants, respectively. These results imply significant outdoor sources.

Implication

Both concentrations of *n*-alkanes and $\Sigma_{16}\text{PAH}$ varied over




Fig. 3. Plot of $\Sigma_{27}\text{AK}$ and $\Sigma_{16}\text{PAH}$ in air-conditioner filter dust (ACFD) from the 1st and 10th floors (a) and from the 20th and 30th floors (b).




Fig. 4. Regression tree predicting $\Sigma_{27}\text{AK}$ (a) and $\Sigma_{16}\text{PAH}$ (b) from floor levels and room types. Each node (rectangle) is labeled with mean (standard deviation, with a unit of $\mu\text{g g}^{-1}$ for $\Sigma_{27}\text{AK}$ and ng g^{-1} for $\Sigma_{16}\text{PAH}$) and the number of samples. The model is read using a top-down approach until terminal nodes appear.

2–3 orders of magnitude in the present study and exhibited significant spatial variations within apartments or rooms. This was mainly due to large variations in contamination on different floor levels of the building. Additional factors, such as the dwellers' living habits (smoking and nonsmoking, cooking and seldom cooking) or additional emission sources, may also contribute to temporal and spatial variations in *n*-alkanes and PAHs concentrations in ACFD (Mannino and Orecchio, 2008; Harrad *et al.*, 2009; Jung *et al.*, 2011; Muenhor and Harrad, 2012; Qi *et al.*, 2014)). Particularly, indoor smoking and cooking are generally considered as the primary contributors of indoor dust bound PAHs (Ren *et al.*, 2006; Qi *et al.*, 2014). Nevertheless, there were no significant differences in concentrations between the samples from smoking and nonsmoking or between cooking and noncooking apartments (all $p > 0.05$), likely due to insufficient number of samples for statistical analysis.

Overall, *n*-alkanes and PAHs were investigated in the residential ACFD from different floor levels of a high-rise apartment. Results clearly suggest that elevated concentrations of *n*-alkanes and PAHs occurred in the samples from low floor levels (1st and 10th floor). Consequently, the cancer risk from exposure to dust bound

PAHs would be predicted to be greater for people living on lower floor levels than on higher floor levels. Previous studies have demonstrated that the vertical variations of ambient air pollution are influenced by emission sources and meteorological conditions. Several investigators also found significant declining trends on aerosol associated PAHs concentrations with height in urban areas (Li *et al.*, 2005; Tao *et al.*, 2007). Thus our results clearly demonstrated that elevated pollution by particle-associated contaminants occurred on lower floor levels compared with high floor levels of high-rise buildings. As mentioned above, the height of a building can greatly influence wind speed and microenvironment; thus it may further affect the vertical diffusion of different sizes of aerosols and the exchange of particle bounded pollutants between indoor and outdoor air. However, knowledge on these aspects remains limited. Additional efforts should also be taken to explore the change of contaminants in particles with different sizes between indoor and outdoor environment at different vertical heights.

CONFLICTS OF INTEREST

The authors declare no competing financial interest.

ACKNOWLEDGEMENTS

The present study was financially supported by the Fundamental Research Funds for the Central Universities (2015HGCH0002 and JZ2016HGTB0711) and the National Natural Science Foundation of China (41390240).

REFERENCE

- Alves, C.A., Oliveira, C., Martins, N., Mirante, F., Caseiro, A., Pio, C., Matos, M., Silva, H.F., Oliveira, C. and Camões, F. (2016). Road tunnel, roadside, and urban background measurements of aliphatic compounds in size-segregated particulate matter. *Atmos. Res.* 168: 139–148.
- Bhargava, A., Khanna, R.N., Bhargava, S.K. and Kumar, S. (2004). Exposure risk to carcinogenic PAHs in indoor-air during biomass combustion whilst cooking in rural India. *Atmos. Environ.* 38: 4761–4767.
- Bi, X.H., Sheng, G.Y., Peng, P.A., Chen, Y.J. and Fu, J.M. (2005a). Size distribution of n-alkanes and polycyclic aromatic hydrocarbons (PAHs) in urban and rural atmospheres of Guangzhou, China. *Atmos. Environ.* 39: 477–487.
- Bi, X., Sheng, G., Liu, X., Li, C. and Fu, J. (2005b). Molecular and carbon and hydrogen isotopic composition of n-alkanes in plant leaf waxes. *Org. Geochem.* 36: 1405–1417.
- Bi, X.H., Sheng, G.Y., Peng, P.A., Chen, Y.J., Zhang, Z.Q. and Fu, J.M. (2003). Distribution of particulate- and vapor-phase n-alkanes and polycyclic aromatic hydrocarbons in urban atmosphere of Guangzhou, China. *Atmos. Environ.* 37: 289–298.
- Bi, X.H., Simoneit, B.R.T., Sheng, G.Y. and Fu, J.M. (2008). Characterization of molecular markers in smoke from residential coal combustion in China. *Fuel* 87: 112–119.
- bin Abas, M.R., Simoneit, B.R.T., Elias, V., Cabral, J.A. and Cardoso, J.N. (1995). Composition of higher molecular weight organic matter in smoke aerosol from biomass combustion in Amazonia. *Chemosphere* 30: 995–1015.
- Cheruiyot, N.K., Lee, W.J., Mwangi, J.K., Wang, L.C., Lin, N.H., Lin, Y.C., Cao, J.J., Zhang, R.J. and Chang-Chien, G.P. (2015). An overview: Polycyclic aromatic hydrocarbon emissions from the stationary and mobile sources and in the ambient air. *Aerosol Air Qual. Res.* 15: 2730–2762.
- Chou, H.M., Kao, C.C., Chuang, K.P., Lin, C., Shy, C.G., Chen, R.F., Tsai, C.C., Chuang, C.Y., Cheng, Y.C., Chen, C.C. and Chao, H.R. (2016). Levels of polybrominated diphenyl ethers in air-conditioner filter dust used to assess health risks in clinic and electronic plant employees. *Aerosol Air Qual. Res.* 16: 184–194.
- Chuang, J.C., Callahan, P.J., Menton, R.G., Gordon, S.M., Lewis, R.G. and Wilson, N.K. (1995). Monitoring methods for polycyclic aromatic hydrocarbons and their distribution in house dust and track-in soil. *Environ. Sci. Technol.* 29: 494–500.
- Cincinelli, A., Bubba, M.D., Martellini, T., Gambaro, A. and Lepri, L. (2007). Gas-particle concentration and distribution of n-alkanes and polycyclic aromatic hydrocarbons in the atmosphere of Prato (Italy). *Chemosphere* 68: 472–478.
- Drennan, I.R., Strum, R.P., Byers, A., Buick, J.E., Lin, S., Cheskes, S., Hu, S. and Morrison, L.J. (2016). Out-of-hospital cardiac arrest in high-rise buildings: Delays to patient care and effect on survival. *Can. Med. Assoc. J.* 188: 413–419.
- Du, X.Q., Wang, X.M., Tang, J.H.D., Xiang, Lin, H.T., Zhong, L., Sheng, G.Y. and Fu, J.M. (2006). Polycyclic Aromatic Hydrocarbons in Mainstream Smokes from Major Cigarette Brands Sold in China. *Environ. Chem. (in Chinese)* 25: 785–788.
- Eglinton, G. and Hamilton, R.J. (1967). Leaf epicuticular waxes. *Science* 156: 1322–1335.
- Essumang, D.K., Ofori, J., Dodoo, D.K. and Adjei, J.K. (2015). Polycyclic aromatic hydrocarbons in settled dust particles in selected Ghanaian environments: Levels, source characterization, and assessment of inhalational health risks. *Indoor Built Environ.* 25: 242–253.
- Feng, J.L., Hu, M., Chan, C.K., Lau, P.S., Fang, M., He, L.Y. and Tang, X.Y. (2006). A comparative study of the organic matter in PM_{2.5} from three Chinese megacities in three different climatic zones. *Atmos. Environ.* 40: 3983–3994.
- Gao, N.P., Niu, J.L., Perino, M. and Heiselberg, P. (2008). The airborne transmission of infection between flats in high-rise residential buildings: Tracer gas simulation. *Build. Environ.* 43: 1805–1817.
- Gevao, B., Al-Bahloul, M., Zafar, J., Al-Matrouk, K. and Helaleh, M. (2007). Polycyclic aromatic hydrocarbons in indoor air and dust in Kuwait: Implications for sources and nondietary human exposure. *Arch. Environ. Contam. Toxicol.* 53: 503–512.
- Gogou, A., Stratigakis, N., Kanakidou, M. and Stephanou, E.G. (1996). Organic aerosols in Eastern Mediterranean: components source reconciliation by using molecular markers and atmospheric back trajectories. *Org. Geochem.* 25: 79–96.
- Harrad, S., Abdallah, M.A.E. and Covaci, A. (2009). Causes of variability in concentrations and diastereomer patterns of hexabromocyclododecanes in indoor dust. *Environ. Int.* 35: 573–579.
- Huang, K.L., Tsai, T.H., Tsai, J.H., Chen, S.J. and Lee, W.J. (2015). Emission of PAHs from a single hydrogen-oxygen PEM fuel cell: In relation to fuel cell carbon materials. *Aerosol Air Qual. Res.* 15: 2654–2667.
- Hwang, H.M., Wade, T.L. and Sericano, J.L. (2003). Concentrations and source characterization of polycyclic aromatic hydrocarbons in pine needles from Korea, Mexico, and United States. *Atmos. Environ.* 37: 2259–2267.
- Jo, W.K. and Kim, K.Y. (2002). Vertical variability of volatile organic compound (VOC) levels in ambient air of high-rise apartment buildings with and without occurrence of surface inversion. *Atmos. Environ.* 36: 5645–5652.
- Jo, W.K., Kim, K.Y., Park, K.H., Kim, Y.K., Lee, H.W. and Park, J.K. (2003). Comparison of outdoor and indoor

- mobile source-related volatile organic compounds between low- and high-floor apartments. *Environ. Res.* 92: 166–171.
- Jo, W.K. and Lee, J.Y. (2006). Indoor and outdoor levels of respirable particulates (PM₁₀) and Carbon Monoxide (CO) in high-rise apartment buildings. *Atmos. Environ.* 40: 6067–6076.
- Jung, D.B., Cho, Y.S., Kim, I.S., Lee, J.Y. and Kim, Y.P. (2015). Impact of energy consumption in northeast Asia to the particulate PAHs levels and composition at Seoul. *Aerosol Air Qual. Res.* 15: 2190–2199.
- Jung, K.H., Bernabe, K., Moors, K., Yan, B., Chillrud, S.N., Whyatt, R., Camann, D., Kinney, P.L., Perera, F.P. and Miller, R.L. (2011). Effects of floor level and building type on residential levels of outdoor and indoor polycyclic aromatic hydrocarbons, black carbon, and particulate matter in New York City. *Atmosphere* 2: 96–109.
- Kang, Y., Cheung, K.C. and Wong, M.H. (2010). Polycyclic aromatic hydrocarbons (PAHs) in different indoor dusts and their potential cytotoxicity based on two human cell lines. *Environ. Int.* 36: 542–547.
- Kavouras, I.G., DuBois, D.W., Nikolich, G. and Etyemezian, V. (2015). Monitoring, source identification and health risks of air toxics in Albuquerque, New Mexico, USA. *Aerosol Air Qual. Res.* 15: 556–571.
- Lee, C.L., Huang, H.C., Wang, C.C., Sheu, C.C., Wu, C.C., Leung, S.Y., Lai, R.S., Lin, C.C., Wei, Y.F., Lai, I.C., Jiang, H., Choug, W.L., Chung, W.Y., Huang, M.S. and Huang, S.K. (2016). A new grid-scale model simulating the spatiotemporal distribution of PM_{2.5}-PAHs for exposure assessment. *J. Hazard. Mater.* 314: 286–294.
- Lee, J.H. and Jo, W.K. (2006). Characteristics of indoor and outdoor bioaerosols at Korean high-rise apartment buildings. *Environ. Res.* 101: 11–17.
- Li, C.L., Fu, J.M., Sheng, G.Y., Bi, X.H., Hao, Y.M., Wang, X.M. and Mai, B.X. (2005). Vertical distribution of PAHs in the indoor and outdoor PM_{2.5} in Guangzhou, China. *Build. Environ.* 40: 329–341.
- Lin, J.J. and Lee, L.C. (2004). Characterization of *n*-alkanes in urban submicron aerosol particles (PM₁). *Atmos. Environ.* 38: 2983–2991.
- Lu, L. and Ip, K.Y. (2009). Investigation on the feasibility and enhancement methods of wind power utilization in high-rise buildings of Hong Kong. *Renewable Sustainable Energy Rev.* 13: 450–461.
- Mannino, M.R. and Orecchio, S. (2008). Polycyclic aromatic hydrocarbons (PAHs) in indoor dust matter of Palermo (Italy) area: Extraction, GC–MS analysis, distribution and sources. *Atmos. Environ.* 42: 1801–1817.
- Masih, J., Singhvi, R., Kumar, K., Jain, V.K. and Taneja, A. (2012). Seasonal variation and sources of polycyclic aromatic hydrocarbons (PAHs) in Indoor and outdoor air in a semi arid tract of northern India. *Aerosol Air Qual. Res.* 12: 515–525.
- Mirhoseini, S.H., Nikaeen, M., Satoh, K. and Makimura, K. (2016). Assessment of airborne particles in indoor environments: Applicability of particle counting for prediction of bioaerosol concentrations. *Aerosol Air Qual. Res.* 16: 1903–1910.
- Mohammed, M.O.A., Song, W.W., Ma, Y.L., Liu, L.Y., Ma, W.L., Li, W.L., Li, Y.F., Wang, F.Y., Qi, M.Y., Lv, N., Wang, D.Z. and Khan, A.U. (2016). Distribution patterns, infiltration and health risk assessment of PM_{2.5}-bound PAHs in indoor and outdoor air in cold zone. *Chemosphere* 155: 70–85.
- Muenhor, D. and Harrad, S. (2012). Within-room and within-building temporal and spatial variations in concentrations of polybrominated diphenyl ethers (PBDEs) in indoor dust. *Environ. Int.* 47: 23–27.
- Murakami, S., Uehara, K. and Komine, H. (1979). Amplification of wind speed at ground level due to construction of high-rise building in urban area. *J. Wind Eng. Ind. Aerodyn.* 4: 343–370.
- Mwangi, J.K., Lee, W.J., Tsai, J.H. and Wu, T.S. (2015). Emission reductions of nitrogen oxides, particulate matter and polycyclic aromatic hydrocarbons by using microalgae biodiesel, butanol and water in diesel engine. *Aerosol Air Qual. Res.* 15: 901–914.
- Ni, H.G., Cao, S.P., Chang, W.J. and Zeng, H. (2011). Incidence of polybrominated diphenyl ethers in central air conditioner filter dust from a new office building. *Environ. Pollut.* 159: 1957–1962.
- Oros, D.R. and Simoneit, B.R.T. (2001). Identification and emission factors of molecular tracers in organic aerosols from biomass burning. Part 2. Deciduous trees. *Appl. Geochem.* 16: 1545–1565.
- Oros, D.R., Abas, M.R.b., Omar, N.Y.M.J., Rahman, N.A. and Simoneit, B.R.T. (2006). Identification and emission factors of molecular tracers in organic aerosols from biomass burning: Part 3. Grasses. *Appl. Geochem.* 21: 919–940.
- Peng, H., Yang, Y., Liu, M. and Zhou, J.L. (2012). PAHs in indoor dust samples in Shanghai's universities: Levels, sources and human exposure. *Environ. Geochem. Health* 34: 587–596.
- Pongpiachan, S. (2015). Incremental lifetime cancer risk of PM_{2.5} bound polycyclic aromatic hydrocarbons (PAHs) before and after the wildland fire episode. *Aerosol Air Qual. Res.* 16: 2907–2919.
- Qi, H., Li, W.L., Zhu, N.Z., Ma, W.L., Liu, L.Y., Zhang, F. and Li, Y.F. (2014). Concentrations and sources of polycyclic aromatic hydrocarbons in indoor dust in China. *Sci. Total Environ.* 491–492: 100–107.
- Reddy, C.M., Eglinton, T.I., Palić, R., Benitez-Nelson, B.C., Stojanović, G., Palić, I., Djordjević, S. and Eglinton, G. (2000). Even carbon number predominance of plant wax *n*-alkanes: A correction. *Org. Geochem.* 31: 331–336.
- Ren, Y., Cheng, T.T. and Chen, J.M. (2006). Polycyclic aromatic hydrocarbons in dust from computers: One possible indoor source of human exposure. *Atmos. Environ.* 40: 6956–6965.
- Schauer, J.J., Rogge, W.F., Hildemann, L.M., Mazurek, M.A., Cass, G.R. and Simoneit, B.R.T. (1996). Source apportionment of airborne particulate matter using organic compounds as tracers. *Atmos. Environ.* 30: 3837–3855.

- Simoneit, B.R.T. (1984). Organic matter of the troposphere: III—characterization and sources of petroleum and pyrogenic residues in aerosols over the western United States. *Atmos. Environ.* 18: 51–67.
- Sun, L.N. (2012). Chemical composition of *n*-alkanes and *n*-alkanoic acids in three types of agricultural straw and the smoke, In *School of Environmental Science and Engineering*, Nanjing University of Information Science & Technology, Nanjing, pp. 37–44.
- Tao, S., Wang, Y., Wu, S.M., Liu, S.Z., Dou, H., Liu, Y.N., Lang, C., Hu, F. and Xing, B.S. (2007). Vertical distribution of polycyclic aromatic hydrocarbons in atmospheric boundary layer of Beijing in winter. *Atmos. Environ.* 41: 9594–9602.
- The World Bank (2016). *World Urbanization Prospects*, The World Bank Group.
- Tiwari, M., Sahu, S.K. and Pandit, G.G. (2015). Inhalation risk assessment of PAH exposure due to combustion aerosols generated from household fuels. *Aerosol Air Qual. Res.* 15: 582–590.
- United Nations Department of Economic and Social Affairs (2014). World's Population Increasingly Urban with More than Half Living in Urban Areas.
- Wang, G., Cheng, S.Y., Wei, W., Wen, W., Wang, X.Q. and Yao, S. (2015). Chemical characteristics of fine particles emitted from different Chinese cooking styles. *Aerosol Air Qual. Res.* 15: 2357–2366.
- Wang, J.Z., Guan, Y.F., Ni, H.G., Luo, X.L. and Zeng, E.Y. (2007). Polycyclic aromatic hydrocarbons in riverine runoff of the Pearl River Delta (China): Concentrations, fluxes, and fate. *Environ. Sci. Technol.* 41: 5614–5619.
- Wang, J.Z., Zhang, K., Liang, B. and Zeng, E.Y. (2011). Occurrence, source apportionment and toxicity assessment of polycyclic aromatic hydrocarbons in surface sediments of Chaohu, one of the most polluted lakes in China. *J. Environ. Monit.* 13: 3336–3342.
- Wang, J.Z., Yang, Z.Y. and Chen, T.H. (2012). Source apportionment of sediment-associated aliphatic hydrocarbon in a eutrophicated shallow lake, China. *Environ. Sci. Pollut. Res.* 19: 4006–4015.
- Wang, Z.Z., Bi, X.H., Sheng, G.Y. and Fu, J.M. (2009). Characterization of organic compounds and molecular tracers from biomass burning smoke in South China I: Broad-leaf trees and shrubs. *Atmos. Environ.* 43: 3096–3102.
- Wu, S.P., Tao, S. and Liu, W.X. (2006). Particle size distributions of polycyclic aromatic hydrocarbons in rural and urban atmosphere of Tianjin, China. *Chemosphere* 62: 357–367.
- Wu, S.P., Tao, S., Zhang, Z.H., Lan, T. and Zuo, Q. (2007). Characterization of TSP-bound *n*-alkanes and polycyclic aromatic hydrocarbons at rural and urban sites of Tianjin, China. *Environ. Pollut.* 147: 203–210.
- Yan, B., Abrajano, T.A., Bopp, R.F., Chaky, D.A., Benedict, L.A. and Chillrud, S.N. (2005). Molecular tracers of saturated and polycyclic aromatic hydrocarbon inputs into Central Park Lake, New York City. *Environ. Sci. Technol.* 39: 7012–7019.
- Yang, X.X., Ren, D., Sun, W.W., Li, X.M., Huang, B., Chen, R., Lin, C. and Pan, X.J. (2015). Polycyclic aromatic hydrocarbons associated with total suspended particles and surface soils in Kunming, China: distribution, possible sources, and cancer risks. *Environ. Sci. Pollut. Res.* 22: 6696–6712.
- Ye, S.F., Zhang, B., Fu, H.H., Tian, N., Shang, H.B., Chen, X.Q. and Wu, S.P. (2013). Emission of fine particles and fine particle-bound polycyclic aromatic hydrocarbons from simulated cooking fumes. *J. Xiamen Univ. (Natural Science)* 52: 824–829.
- Zhang, L.W., Bai, Z.P., You, Y., Wu, J.H., Feng, Y.C. and Zhu, T. (2009). Chemical and stable carbon isotopic characterization for PAHs in aerosol emitted from two indoor sources. *Chemosphere* 75: 453–461.
- Zhang, S., Peng, S.C., Chen, T.H. and Wang, J.Z. (2015). Evaluation of inhalation exposure to carcinogenic PM₁₀-bound PAHs of people at night markets of an urban area in a metropolis in eastern China. *Aerosol Air Qual. Res.* 15: 1944–1954.
- Zhang, X.L., Tao, S., Liu, W.X., Yang, Y., Zuo, Q. and Liu, Z.S. (2005). Source Diagnostics of polycyclic aromatic hydrocarbons based on species ratios: A multimedia approach. *Environ. Sci. Technol.* 39: 9109–9114.
- Zheng, H.F., Chen, L.D., Han, X.Z., Zhao, X.F. and Ma, Y. (2009). Classification and regression tree (CART) for analysis of soybean yield variability among fields in Northeast China: The importance of phosphorus application rates under drought conditions. *Agric. Ecosyst. Environ.* 132: 98–105.
- Zhou, B. and Zhao, B. (2012). Population inhalation exposure to polycyclic aromatic hydrocarbons and associated lung cancer risk in Beijing region: Contributions of indoor and outdoor sources and exposures. *Atmos. Environ.* 62: 472–480.
- Zhou, H.C., Zhang, C.C., Cai, H.X., Song, Y.Y. and Xue, H.B. (2010). The distribution and possible source of polycyclic aromatic hydrocarbons in dust from six air conditioner filters. *China Environ. Sci.* 30: 1303–1308.

Received for review, March 28, 2017

Revised, May 10, 2017

Accepted, May 13, 2017