



## Review

# Impacts and Effects Indicators of Atmospheric Deposition of Major Pollutants to Various Ecosystems - A Review

L. Paige Wright<sup>1</sup>, Leiming Zhang<sup>2\*</sup>, Irene Cheng<sup>2</sup>, Julian Aherne<sup>3</sup>, Gregory R. Wentworth<sup>4</sup>

<sup>1</sup> *Independent Researcher, Stratford, Prince Edward Island, Canada*

<sup>2</sup> *Air Quality Research Division, Science and Technology Branch, Environment and Climate Change Canada, Toronto, Ontario, Canada*

<sup>3</sup> *Environmental and Resource Studies, Trent University, Peterborough, Ontario, Canada*

<sup>4</sup> *Environmental Monitoring and Science Division, Alberta Environment and Parks, Edmonton, Alberta, Canada*

---

## ABSTRACT

In this paper, we review the current understanding on ecosystem and human health impacts from the atmospheric deposition of acidifying pollutants, eutrophying nitrogen (N), polycyclic aromatic hydrocarbons (PAHs), mercury (Hg), trace metals, and ozone (O<sub>3</sub>), as well as the biological indicators that have been used to assess the health of ecosystems following exposure to these pollutants. We provide overviews of the impacts of deposition for these pollutants and discuss the currently known biomonitors for each pollutant. The deposition of acidifying pollutants impacts terrestrial ecosystems by altering plant physiology and growth and by increasing plant susceptibility to stresses that can be indirectly damaging to the health of fish and birds. Indicators of the deposition of acidifying pollutants include soil base cation content and acid neutralizing capacity, among others. Eutrophying N deposition has been studied extensively; N enrichment directly impacts vegetative plant species cover, richness, growth rates, and susceptibility to other stressors. It indirectly impacts wildlife through changes in their habitats and food sources. Indicators for N deposition include changes in plant species and in tissue and litter N content. The deposition of PAHs has been found to cause significant damage to plant organisms and to be carcinogenic and mutagenic to humans and animals. Useful biomonitors of PAH deposition include lichens, mosses, and pine needles. Deposited Hg can undergo methylation (in the presence of sulphur reducing bacteria); bioaccumulation of methylmercury is highly toxic to animals. Effective biomonitors of Hg contamination of aquatic ecosystems are fish and marine birds. The impacts of O<sub>3</sub> are well understood, with well-established “flux” models being vast improvements on the previous AOT40 approaches. This review highlights the impacts that the above-mentioned pollutants have on terrestrial and aquatic organisms and the biomonitors that are currently being used to assess the deposition levels and effects of these pollutants.

**Keywords:** Air pollution; Atmospheric deposition; Ecosystem health; Ecological monitoring.

---

## INTRODUCTION

Atmospheric deposition is a major source of pollutants to terrestrial (e.g., forests, grasslands, ombrotrophic wetlands) and aquatic (rivers, lakes, oceans) environments (Lepori and Keck, 2012). Other non-depositional processes (e.g., effluent discharge, run-off, leaching) are also important pathways for pollutant input, especially to aquatic ecosystems (Vodyanitskii, 2013). Atmospheric deposition of pollutants can cause detrimental effects to both terrestrial and aquatic environments and to human and wildlife health (Driscoll *et*

*al.*, 2001; Baldigo *et al.*, 2009; Bobbink *et al.*, 2010; Bacon *et al.*, 2013; Vodyanitskii, 2013; de Vries *et al.*, 2014; Jones *et al.*, 2014; de Vries *et al.*, 2015; Duan *et al.*, 2016; Rodríguez-Estival and Smits, 2016). Once deposited onto land and water surfaces, pollutants are incorporated into soil or water where they can be taken up directly by biota, methylated by microorganisms in the case of mercury (Hg), or increase the leaching of soil base cations, metals, and plant nutrients in the case of acidifying pollutants. Moreover, pollutants can also be transported from where they were initially deposited via soil erosion, surface run-off, and groundwater flows. Physical, chemical and biological variables can affect pollutant uptake and toxicity in biota. Exposure to pollutants can lead to wide-ranging ecological consequences. Methylmercury is a neurotoxin capable of bio-accumulating in wildlife and biomagnifying in food

---

\* Corresponding author.

*E-mail address:* leiming.zhang@canada.ca

webs; humans are typically exposed via fish or other wildlife consumption (Wiener *et al.*, 2012). Depletion of plant nutrients and leaching of toxic inorganic aluminum due to acid rain can damage vegetation, increase stress to and reduce tolerance of plants, and inhibit tree growth. Acidified lakes are also detrimental to aquatic wildlife (Driscoll *et al.*, 2001). Polycyclic aromatic compounds (PACs) and its subgroup of polycyclic aromatic hydrocarbons (PAHs) comprise hundreds of hydrocarbons, some of which are cancer-causing agents or capable of causing damage to DNA or reproductive impairment (Kim *et al.*, 2013; Abdel-Shafy and Mansour, 2016).

Despite the knowledge that deposition occurs, monitoring of emissions, ambient air, and deposition is often conducted separately from the monitoring of ecosystem effects leading to gaps in the understanding of sources and fate of pollutants in the environment (Ochoa-Hueso *et al.*, 2017). In the former case, researchers are able to track the variability in the emissions, air concentrations and deposition but are uncertain about the post-deposition effects and how these changes will impact ecosystems. In the latter, researchers are able to assess the ecosystem effects, but are uncertain what the sources of contamination are and the role of atmospheric deposition versus the physical, chemical, and biological factors influencing pollutant uptake. These knowledge gaps may be filled by establishing stronger connections between atmospheric sources or deposition and ecological effects. This can be accomplished through the identification of biomonitors and their associated effects from pollutant exposure and in the determination of the critical loads and critical levels at which these effects occur. A biomonitor is defined as a change in biological responses, ranging from molecular through cellular and physiological responses to behavioral changes, which can be related to exposure to, or toxic effects of, environmental chemicals (Peakall and Walker, 1994). A biological indicator refers to a living organism that provides information on an ecosystem. A critical load refers to a concentration limit in any media (air, water, soil, tissue). In this review paper, we examine the ecosystem and human health impacts from atmospheric deposition of acidifying pollutants, eutrophying nitrogen (N), polycyclic aromatic hydrocarbons (PAH), Hg, trace metals, and ozone (O<sub>3</sub>), as well as the biological indicators that have been used to assess the health of ecosystems. Deposition studies on other groups of pollutants have also been made in literature (Lin *et al.*, 2010; Weissengruber *et al.*, 2018), but the information was not as rich as for those pollutants mentioned above, and thus were not included in this review.

## ACIDIC DEPOSITION

### Major Acidic Species

Major acidic species in the atmosphere include gases (SO<sub>2</sub>, HNO<sub>3</sub>, HCl, and organic acids) and particulate matters (sulphate, nitrate, chloride, organic acids), and associated aqueous-phase species (Seinfeld and Pandis, 2016). The components of acidic deposition vary based on the local and regional emissions but primarily consist of sulphur

species. The relative contribution of each pollutant to total acidic deposition also depends on meteorology (e.g., amount of precipitation, sunlight, wind speed), which can impact formation rates as well as deposition velocities. Typically, oxidized sulphur (i.e., SO<sub>2</sub> and SO<sub>4</sub><sup>2-</sup>) and nitrogen (primarily HNO<sub>3</sub> and NO<sub>3</sub><sup>-</sup>) dominate acidic deposition in polluted areas (Vet *et al.*, 2014). The deposition of reduced nitrogen (NH<sub>x</sub> = NH<sub>3</sub> + NH<sub>4</sub><sup>+</sup>) can also acidify terrestrial ecosystems (soils) through two indirect mechanisms: (i) NH<sub>4</sub><sup>+</sup> uptake into roots (which releases H<sup>+</sup>), or (ii) nitrification to NO<sub>3</sub><sup>-</sup> (which produces H<sup>+</sup>) (Krupa, 2003). Base cation species (e.g., Ca<sup>2+</sup>, Mg<sup>2+</sup>, Na<sup>+</sup>, K<sup>+</sup>) can mitigate acidic deposition by neutralizing acidic species, and should be taken into account when assessing acidification of receiving ecosystems (Watmough *et al.*, 2014). The characteristics of acidifying pollutants and base cations, as well as commonly used indicators, are summarized in Table 1.

### Impacts of Acidic Deposition

It is well established that acidic (sulphur and nitrogen) deposition can decrease soil pH, and increase soil solution concentrations of Al<sup>3+</sup>, resulting in the leaching of base cations (Ca<sup>2+</sup>, Mg<sup>2+</sup>, and K<sup>+</sup>). Direct effects from deposition on leaves are manifested in changes in leaf physiology, including an increase in leaf roundness (Bacon *et al.*, 2013), changes in the epicuticular and epistomatal waxes (Bartromo *et al.*, 2012; Elliott-Kingston *et al.*, 2014), smaller leaves, and changes to the cuticle and stomata (Haworth *et al.*, 2010; Haworth *et al.*, 2012; Elliott-Kingston *et al.*, 2014). Impacts on forests include nutrient deficiencies due to soil base cation leaching (Akselsson *et al.*, 2007), leading to decreased growth (Savva and Berninger, 2010), changes in tree ring width (Godek *et al.*, 2015), increased susceptibility to disease, destruction of root systems resulting in tree and crown dieback (Wilmot *et al.*, 1995; Godek *et al.*, 2015), and higher sensitivity to drought (Savva and Berninger, 2010). The effects of acid deposition have been found to have a geographical dependence (de Vries *et al.*, 1995). Regional properties of an ecosystem that have been found to relate to acid deposition include soil chemistry, such as texture, pH, and humidity, and forest properties, such as tree species and age (de Vries *et al.*, 1995; Økland *et al.*, 2004; de Vries *et al.*, 2007; Stevens *et al.*, 2009a; van Dobben and de Vries, 2010).

Impacts on aquatic ecosystems from acidic deposition are fairly well understood; the mobilisation of Al from soils and decreased water pH can lead to reduced fish species richness, changes in fish health, and degraded water quality. The reduction and extinction of Atlantic salmon in the Maritimes (Watt *et al.*, 2000) and fish populations elsewhere (Monteith *et al.*, 2005; Gray *et al.*, 2012) have been associated with increased water acidity.

Acidic deposition may impact wildlife by reducing the microbial biomass and shredder (macroinvertebrates that feed on plant and animal material and break it into smaller particles) biomass of gammarids (Meegan *et al.*, 1996) and caddisflies (Simon *et al.*, 2009), suppressing microbial respiration (Dangles and Guérol, 2001; Simon *et al.*, 2009), depressing aquatic fungi (Clivot *et al.*, 2014), damaging the gills of fish with gill lesions due to increases in Al and

**Table 1.** Characteristics of Acidifying Pollutants (N & S) and Base Cations ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^{+}$ ,  $\text{Na}^{+}$ ).

Element	Acidifying Pollutants		Base Cations (Alkaline)			
	Nitrogen (N)	Sulfur (S)	Calcium ( $\text{Ca}^{2+}$ )	Magnesium ( $\text{Mg}^{2+}$ )	Potassium ( $\text{K}^{+}$ )	Sodium ( $\text{Na}^{+}$ )
Atomic Number	7	16	20	12	19	11
Group	15	16	2	2	1	1
Period	2	3	4	3	4	3
Atomic Mass	14.00067 $\text{g mol}^{-1}$	32.06 $\text{g mol}^{-1}$	40.08 $\text{g mol}^{-1}$	24.305 $\text{g mol}^{-1}$	39.0983 $\text{g mol}^{-1}$	22.98977 $\text{g mol}^{-1}$
Series	Nonmetal	Nonmetal	Alkali Earth Metal	Alkali Earth Metal	Alkali Metal	Alkali Metal
Interesting	Inert in elemental form	Occurs naturally near volcanoes	<ul style="list-style-type: none"> <li>Occurs only in compounds (because it is so reactive)</li> </ul>			
Main Sources	On- and Off-Road vehicles, the oil and gas industry, and fuel for electricity generation and heating	Smelting and refining non-ferrous metals, coal-fired electricity generation, and the petroleum industry	<ul style="list-style-type: none"> <li>Dust associated with mining activities.</li> <li>Minerals like chalk, limestone, and marble.</li> <li>Atmospheric deposition and weathering.</li> </ul>	<ul style="list-style-type: none"> <li>Deposition of fertilizer</li> </ul>	<ul style="list-style-type: none"> <li>Weathering of primary materials</li> <li>Deposition of fertilizer</li> </ul>	
Common Compounds	<ul style="list-style-type: none"> <li>Organic Nitrogen Compounds (ONC)</li> <li>Nitric Oxide (NO)</li> <li>Nitrogen Oxides (<math>\text{NO}_x</math>)</li> <li>Nitrogen Dioxide (<math>\text{NO}_2</math>)</li> <li>Nitric Acid (<math>\text{HNO}_3</math>)</li> <li>Particulate Nitrate (<math>\text{NO}_3^-</math>)</li> <li>Particulate Nitrite (<math>\text{NO}_2^-</math>)</li> <li>Ammonia (<math>\text{NH}_3</math>)</li> <li>Ammonium (<math>\text{NH}_4^+</math>)</li> </ul>	<ul style="list-style-type: none"> <li>Sulfuric Acid (<math>\text{H}_2\text{SO}_4</math>)</li> <li>Sulphur Oxides (<math>\text{SO}_x</math>)</li> <li>Sulphur Dioxide (<math>\text{SO}_2</math>)</li> <li>Sulphate (<math>\text{SO}_4^{2-}</math>)</li> </ul>	<ul style="list-style-type: none"> <li>Limestone (<math>\text{CaCO}_3</math>)</li> <li>Gypsum (<math>\text{CaSO}_4 \cdot 2\text{H}_2\text{O}</math>)</li> <li>Fluorite (<math>\text{CaF}_2</math>)</li> <li>Calcium Chloride (<math>\text{CaCl}_2</math>)</li> </ul>	<ul style="list-style-type: none"> <li>Magnesium Chloride (<math>\text{MgCl}_2</math>)</li> </ul>	<ul style="list-style-type: none"> <li>Potassium Chloride (KCl)</li> </ul>	<ul style="list-style-type: none"> <li>Sodium Chloride (NaCl)</li> <li>Sodium Nitrate (<math>\text{NaNO}_3</math>)</li> </ul>
Indicators	<ul style="list-style-type: none"> <li>N concentration in plants</li> <li>Tissue N content</li> <li>Species richness</li> <li>Forb richness</li> <li>Graminoid:Forb ratio</li> <li>Shifts in plant communities</li> <li>Shifts in diatom communities</li> <li>Nitrate concentrations</li> <li>C:N ratio in soil</li> <li>Dissolved nitrogen</li> <li>Dissolved phosphorus</li> </ul>	<ul style="list-style-type: none"> <li>S content</li> <li>S:N ratio</li> <li>Tree leaves</li> <li>Tree rings</li> <li>Physiognomic leaf analysis</li> <li>Soil percent base saturation</li> <li>Acid neutralizing capacity</li> </ul>	<ul style="list-style-type: none"> <li>Base cation surplus in soil</li> <li>Ca:Al<sup>3+</sup> ratios in soil</li> <li>(Ca+Mg):S ratio in throughfall</li> <li>Foliar chemistry</li> <li>Soil water chemistry</li> <li>pH</li> <li>Inorganic Al<sup>3+</sup></li> <li>Calcium</li> </ul>			

decreases in calcium (Evans *et al.*, 1988), and thinning of bird egg shells (Hames *et al.*, 2002). Effects such as those mentioned above can both directly and indirectly affect macroinvertebrates (Baldigo *et al.*, 2009; Ferreira and Guérol, 2017). Tables 2, 3, and 4 summarize the key points of these studies that examine the impacts of N, S, and soil base cation deposition, respectively.

### **Effects Indicators of Acidic Deposition**

Effects indicators of acidification on terrestrial ecosystems include S and N tissue concentrations (Wieder *et al.*, 2016); tree ring widths (Lee *et al.*, 2011; Godek *et al.*, 2015; van der Sleen *et al.*, 2015); tree bark acidity (Grodzińska, 1977); and the molar ratio of Ca:Al<sup>3+</sup> (or base cation (BC):Al<sup>3+</sup>) in soil solution (de Vries, 1993; Vangelova *et al.*, 2007; Kleijn *et al.*, 2008; Jorge *et al.*, 2011; de Vries *et al.*, 2015; Irvine *et al.*, 2017), where a ratio < 1 implies a high risk and > 10 implies low risk; soil pH (Kleijn *et al.*, 2008; Tian and Niu, 2015); soil solution Al<sup>3+</sup>, exchangeable base cation (BC) content in soil (Lucas *et al.*, 2011; Tian and Niu, 2015); soil percent base saturation, with increased risk at < 10% and lower risk at > 30%; foliar chemistry, with limited sugar maple growth at levels of Ca < 5000 ppm and Mg < 700 ppm (Fenn *et al.*, 2011); leaf physiogenomic traits (Bacon *et al.*, 2013); leaf litter content (Meegan *et al.*, 1996; Simon *et al.*, 2009; Clivot *et al.*, 2014); cuticle micromorphology (Bartirolo *et al.*, 2012); and isotopic ratio fingerprinting (Hogan *et al.*, 2017).

Indicators for aquatic ecosystems include acid neutralizing capacity (ANC =  $\Sigma$  base cations –  $\Sigma$  strong acid anions) where values lower than a threshold of 20  $\mu\text{eq L}^{-1}$  result in established risks to aquatic life (Lien *et al.*, 1996; Bulger *et al.*, 2000; Fölster *et al.*, 2007; Futter *et al.*, 2014; Duan *et al.*, 2016; Sullivan *et al.*, 2017); pH, with a reduction in fish species below a pH of 5.0–6.0 (Jeffries and Lam, 1993; Rask *et al.*, 1995; Futter *et al.*, 2014); inorganic Al<sup>3+</sup>, with concentrations above 2  $\mu\text{mol L}^{-1}$  is deemed toxic to aquatic organisms (Sjostedt *et al.*, 2012); calcium, where concentrations below 1.5  $\text{mg L}^{-1}$  negatively affect crustaceans (Wærvågen *et al.*, 2002); leaf-associated fungi (Clivot *et al.*, 2014); base cation surplus in streams (Baldigo *et al.*, 2009); and macroinvertebrates diversity using the acid biological assessment profile (acidPAPD) index (Baldigo *et al.*, 2009). In acidified streams, leaf litter analysis is a useful bioassessment tool on stream health as leaf litter decomposition rates increase with increasing acidity (Dangles and Guérol, 2001; Simon *et al.*, 2009; Ferreira and Guérol, 2017).

## **NITROGEN DEPOSITION**

### **Major N-deposition Species**

Similar to acidic deposition, the major components of N-deposition vary from region to region and depend on local emissions and meteorology. Generally, N-deposition is dominated by inorganic reduced nitrogen (NH<sub>3</sub> and NH<sub>4</sub><sup>+</sup>) and oxidized nitrogen (NO<sub>x</sub>, HNO<sub>3</sub> and NO<sub>3</sub><sup>-</sup>) (e.g., Harrison *et al.*, 1999; Hayden *et al.*, 2003; Horii *et al.*, 2005; Zhang *et al.*, 2009; Benedict *et al.*, 2013; van den Elzen *et*

*al.*, 2018). Although organic N deposition is not typically measured, it has been shown to constitute a significant fraction (upwards of ~30%) at some field sites (Zhang *et al.*, 2009; Benedict *et al.*, 2013). Due to the variety of atmospheric N species, as well as technical challenges measuring certain N species (e.g., NH<sub>3</sub> (von Bobrutski *et al.*, 2010); organic N (Beem *et al.*, 2010); HONO (VandenBoer *et al.*, 2013)), complete accounting of total N deposition for a given ecosystem can be challenging.

### **Impacts of Nitrogen Deposition**

Exposure to N can cause a wide variety of well documented impacts on terrestrial and aquatic ecosystems. Below the critical level, N can be beneficial to an environment, such as causing an increase in biomass and CO<sub>2</sub> uptakes (Butterbach-Bahl and Dannenmann, 2011; Jones *et al.*, 2014; Carter *et al.*, 2017; Xu *et al.*, 2017). However, above the critical level, the effects of N tend to be negative. Impacts include direct toxicity from dry deposition to individual species (Marschner *et al.*, 1996; Carter *et al.*, 2017), eutrophication, acidification (as discussed above), and negative impacts on plant species diversity owing to changes in the form of available N, such as oxidized N (from e.g., vehicle emissions, industry, and domestic combustion) and reduced N (from e.g., agricultural sources, biomass burning) (Southon *et al.*, 2013; Field *et al.*, 2014; Jones *et al.*, 2014; van den Berg *et al.*, 2016). Long-term exposure to terrestrial ecosystems can cause significant declines in plant cover including lichens (Bobbink *et al.*, 2010; Stevens *et al.*, 2012; Field *et al.*, 2014; Bähring *et al.*, 2017; Carter *et al.*, 2017) and mosses and herbaceous plants (Dirkse and Martakis, 1992; Hallingbäck and Kellner, 1992; Mountford *et al.*, 1993; Morecroft *et al.*, 1994; Mountford *et al.*, 1994; Kirkham *et al.*, 1996; Carroll *et al.*, 2000; Strengbom *et al.*, 2001; Pitcairn *et al.*, 2006; Koranda *et al.*, 2007; Bobbink *et al.*, 2010; Caporn *et al.*, 2014; Field *et al.*, 2014; Bähring *et al.*, 2017). However, N addition has also been found to cause an increase in the cover of N tolerant (nitrophilic) plant species (Verhoeven *et al.*, 2011; Field *et al.*, 2014; Bähring *et al.*, 2017). Impacts of N enrichment to grasslands include loss of species richness (Stevens *et al.*, 2004; Stevens *et al.*, 2006; Shepherd *et al.*, 2009; Stevens *et al.*, 2009b; Bobbink *et al.*, 2010; Sheppard *et al.*, 2011; Stevens *et al.*, 2012; Isbell *et al.*, 2013; Henry and Aherne, 2014; Bähring *et al.*, 2017), an increase in growth rate (Power *et al.*, 1995; Power *et al.*, 2006), an increase in flower formation (Power *et al.*, 1995; Power *et al.*, 2006; Bähring *et al.*, 2017), a decrease in soil C:N ratio (Bähring *et al.*, 2017) and an increase in shoot increments (Bähring *et al.*, 2017). Other implications include increased susceptibility to secondary stresses and disturbances, such as pests, drought, frost, and disease, and indirectly to exotic earthworm activity (Vitousek *et al.*, 1997; Bobbink *et al.*, 1998; Gilliam, 2006; de Vries *et al.*, 2015; Simkin *et al.*, 2016; Soons *et al.*, 2016; Shao *et al.*, 2017).

In forest ecosystems, impacts include changes in plant species diversity and decomposition inhibition; increases in canopy size, water shortages, tree mortality, soil acidity, and Al mobility; and decreases in Ca availability, fine-root biomass, understory plant cover, and dwarf-shrub species

**Table 2.** Impacts of Acidifying Pollutant: Nitrogen (N).

Ecosystem Type/Species	Location	Approach	Ecological Impact	Effects Indicator	Comments	Reference
<b>Terrestrial Ecosystems</b>						
Calcareous dunes	Newborough Warren, North Wales, UK	Chronosequences and soil sampling Acidification	<ul style="list-style-type: none"> <li>• Accumulation of soil C and N</li> <li>• Suppression of N<sub>2</sub> fixation</li> <li>• Leaching of excess N (low retention capacity for N in dune soils)</li> </ul>	Chronosequences and soil sampling	Low N deposition. Regression modelling	Aggenbach et al., 2016
Calcareous dunes	Luchterduinen, the Netherlands	Chronosequences and soil sampling Acidification	<ul style="list-style-type: none"> <li>• Accelerated acidification</li> <li>• Decalcification of the topsoil</li> </ul>	Chronosequences and soil sampling	High N deposition. Regression modelling	Aggenbach et al., 2016
Acidic dunes	Luchterduinen, the Netherlands	Chronosequences and soil sampling Acidification	<ul style="list-style-type: none"> <li>• Accelerated acidification.</li> <li>• Accelerated N accumulation rates</li> </ul>	Chronosequences and soil sampling	High N deposition. Regression modelling	Aggenbach et al., 2016
<i>Calluna vulgaris</i> (Dwarf shrubs)	Fehmarn Island, Baltic Sea, North Germany	6 levels of N fertilization	<ul style="list-style-type: none"> <li>• Decrease of cover</li> <li>• Increase of shoot increments for low-N treatment</li> <li>• Increase in flower formation for high-N treatment</li> </ul>	<ul style="list-style-type: none"> <li>• Shoot increment most sensitive indicator</li> <li>• Soil C:N ratios deemed inappropriate indicator</li> </ul>	Current critical N loads confirmed in study (3 yr study)	Bähring et al., 2017
<i>Carex arenaria</i> and <i>Festuca ovina</i> (Graminoids)	Fehmarn Island, Baltic Sea, North Germany	6 levels of N fertilization	<ul style="list-style-type: none"> <li>• Increase of cover</li> <li>• Decrease in C:N ratio</li> </ul>		Current critical N loads confirmed in study (3 yr study)	Bähring et al., 2017
Bryophytes	Fehmarn Island, Baltic Sea, North Germany	6 levels of N fertilization	<ul style="list-style-type: none"> <li>• Decrease of cover</li> <li>• Decrease in C:N ratio</li> </ul>		Current critical N loads confirmed in study (3 yr study)	Bähring et al., 2017
Lichens	Fehmarn Island, Baltic Sea, North Germany	6 levels of N fertilization	<ul style="list-style-type: none"> <li>• Decrease of cover</li> <li>• Decrease in C:N ratio for low-N treatment</li> </ul>		Current critical N loads confirmed in study (3 yr study)	Bähring et al., 2017
Sand dunes			<ul style="list-style-type: none"> <li>• Sand dune binding by taller, more vigorous vegetation</li> <li>• Promotes higher dune building</li> <li>• Improves flood defences</li> </ul>			Barbier et al., 2008
<i>Elymus Canadensis</i>	Lake Michigan, USA	N-addition to soil (15 and 135 kg N ha <sup>-1</sup> yr <sup>-1</sup> NH <sub>4</sub> NO <sub>3</sub> )	<ul style="list-style-type: none"> <li>• No or poor emergence in field plots</li> <li>• Enhanced seed germination in greenhouse</li> </ul>			Bird and Choi, 2016
<i>Echinacea purpurea</i>	Lake Michigan, USA	N-addition to soil (15 and 135 kg N ha <sup>-1</sup> yr <sup>-1</sup> NH <sub>4</sub> NO <sub>3</sub> )	<ul style="list-style-type: none"> <li>• No emergence in field plots</li> <li>• Enhanced seed germination in greenhouse</li> </ul>			Bird and Choi, 2016

Table 2. (continued).

Ecosystem Type/Species	Location	Approach	Ecological Impact	Effects Indicator	Comments	Reference
<i>Dalea purpurea</i>	Lake Michigan, USA	N-addition to soil (15 and 135 kg N ha <sup>-1</sup> yr <sup>-1</sup> NH <sub>4</sub> NO <sub>3</sub> )	<ul style="list-style-type: none"> <li>Reduced seed germination in greenhouse</li> </ul>			Bird and Choi, 2016
<i>Elymus</i>	Lake Michigan, USA	N-addition to soil (15 and 135 kg N ha <sup>-1</sup> yr <sup>-1</sup> NH <sub>4</sub> NO <sub>3</sub> )	<ul style="list-style-type: none"> <li>Reduced seed germination in greenhouse and no emergence in field plots.</li> </ul>			Bird and Choi, 2016
<i>Panicum virgatum</i>	Lake Michigan, USA	N-addition to soil (15 and 135 kg N ha <sup>-1</sup> yr <sup>-1</sup> NH <sub>4</sub> NO <sub>3</sub> )	<ul style="list-style-type: none"> <li>Increased germination in greenhouse</li> <li>Increased germination in field plots in early years</li> </ul>			Bird and Choi, 2016
<i>Lupinus perennis</i>	Lake Michigan, USA	N-addition to soil (15 and 135 kg N ha <sup>-1</sup> yr <sup>-1</sup> NH <sub>4</sub> NO <sub>3</sub> )	<ul style="list-style-type: none"> <li>Low germination in greenhouse</li> <li>Substantial establishment in field plots</li> </ul>			Bird and Choi, 2016
<i>Monarda punctata</i>	Lake Michigan, USA	N-addition to soil (15 and 135 kg N ha <sup>-1</sup> yr <sup>-1</sup> NH <sub>4</sub> NO <sub>3</sub> )	<ul style="list-style-type: none"> <li>Low germination in greenhouse</li> <li>Substantial establishment in field plots</li> </ul>			Bird and Choi, 2016
<i>Schizachyrium scoparium</i>	Lake Michigan, USA	N-addition to soil (15 and 135 kg N ha <sup>-1</sup> yr <sup>-1</sup> NH <sub>4</sub> NO <sub>3</sub> )	<ul style="list-style-type: none"> <li>Low germination in greenhouse</li> <li>Substantial establishment in field plots</li> </ul>			Bird and Choi, 2016
<i>Rudbeckia hirta</i>	Lake Michigan, USA	N-addition to soil (15 and 135 kg N ha <sup>-1</sup> yr <sup>-1</sup> NH <sub>4</sub> NO <sub>3</sub> )	<ul style="list-style-type: none"> <li>No emergence in field plots</li> </ul>			Bird and Choi, 2016
Peat bogs	Europe	Peat litter decomposition	<ul style="list-style-type: none"> <li>Reduction in N limitations on microbial metabolism</li> <li>Higher dissolved organic carbon release</li> </ul>	<ul style="list-style-type: none"> <li>N:P ratio in litter</li> <li>C:N ratio in litter</li> </ul>		Bragazza <i>et al.</i> , 2006
Lichens and bryophytes	Europe (review previous studies)	Long-term NH <sub>3</sub> exposure			Annual critical level for lichens and bryophytes suggested to be 1 µg NH <sub>3</sub> m <sup>-3</sup> for long-term (several years)	Cape <i>et al.</i> , 2009
Herbaceous plants	Europe (review previous studies)	Long-term NH <sub>3</sub> exposure			Annual critical level for herbaceous species suggested to be 3 ± 1 µg NH <sub>3</sub> m <sup>-3</sup> for long-term (several years)	Cape <i>et al.</i> , 2009

Table 2. (continued).

Ecosystem Type/Species	Location	Approach	Ecological Impact	Effects Indicator	Comments	Reference
Calcifugous grasslands	UK	Long-term experimental N addition	Reduction in bryophyte cover and in cover of individual species			Carroll et al., 2000
Lichens		N uptake measured				Dahlman et al., 2004
<i>Holycomium splendens</i>	Europe	N eutrophication	Decline in species			Dirkse and Martakis, 1992; Hallingbäck and Kellner, 1992; Strengbom et al., 2001; Koranda et al., 2007
Forest floor vascular plants	Europe		Decrease in oligotrophic species with increasing CL exceedance			Dirnböck et al., 2014
Grasslands			Acidification of soil and metal availability			Falkengren-Grerup et al., 1987
Calcareous dunes (acid grassland bog, upland heath, lowland heath, and sand dune)	Great Britain	N deposition Acidification and eutrophication	<ul style="list-style-type: none"> <li>Decline in plant species richness</li> <li>Changed species composition</li> <li>Decline in diversity of mosses, lichens, and forbs</li> <li>Increase in graminoids cover</li> </ul>	<ul style="list-style-type: none"> <li>Decline in plant species richness</li> <li>Changed species composition</li> <li>Soil pH levels</li> </ul>	<ul style="list-style-type: none"> <li>135 locations surveyed (2 m × 2 m quadrats)</li> <li>Decline in forb diversity below pH of 3.8</li> <li>Decline in acid grassland richness below pH of 4.5</li> </ul>	Field et al., 2014
			Particulate Nitrate (NO <sub>3</sub> <sup>-</sup> ) Leaching causes acidification of soil			Fowler et al., 2013
	127 sampling sites in China		Higher leaf N concentrations correspond to high atmospheric N deposition.		(2094 observations in literature 1980–2005). Higher than critical load.	Gundersen and Rasmussen, 1990 Han et al., 2005
<i>Nardus stricta</i> and <i>Deschampsia flexuosa</i>		Testing if vascular plants are feasible bio-indicators of atmospheric N deposition				Hicks et al., 2000

Table 2. (continued).

Ecosystem Type/Species	Location	Approach	Ecological Impact	Effects Indicator	Comments	Reference
Mobile and semi-fixed coastal dunes	England and Wales, UK	N deposition gradient	<ul style="list-style-type: none"> <li>• Increase in height and cover of <i>Ammophila arenaria</i></li> </ul>	No common indicator common to all dune sites	Threshold around 15 kg N ha <sup>-1</sup> yr <sup>-1</sup> suggested	Jones et al., 2004
Fixed dunes	England and Wales, UK	N deposition gradient	<ul style="list-style-type: none"> <li>• Increase in biomass</li> <li>• Increase in the cover of certain species</li> <li>• Reduced N availability</li> </ul>	<ul style="list-style-type: none"> <li>• Above ground biomass</li> <li>• Soil C:N ratio</li> </ul>		Jones et al., 2004
Grasslands and heathland	Europe				Review paper includes management and N budgets for grasslands and heathland	Jones et al., 2017
Grasslands			Acidification of soil and metal availability			Jönsson et al., 2003
<i>Hippophae rhamnoides</i>		N-fixing experiments	N <sub>2</sub> fixation can be switched off when mineral N in soil is sufficient.			Kato et al., 2007
Temperate grasslands		Asymbiotic N <sub>2</sub> fixation				Keuter et al., 2014
Calcareous Grey dunes	NW Europe		<ul style="list-style-type: none"> <li>• Decline in plant species richness. Decalcification of top soil.</li> <li>• Grass encroachment not a major problem (1990–2014). Calcareous dunes less sensitive than acidic dunes.</li> </ul>		Extra N from ammonia emission from beaches. Critical load of 15 kg ha <sup>-1</sup> yr <sup>-1</sup> reached.	Kooijman et al., 2016
Acidic Grey dunes	NW Europe		<ul style="list-style-type: none"> <li>• Decline in plant species richness (functioning and species richness are very sensitive to high N-deposition). Grass encroachment. Net N-mineralization increased with N-deposition.</li> <li>• Grazing is needed to counteract grass encroachment</li> </ul>		Critical load of 10 kg ha <sup>-1</sup> yr <sup>-1</sup> exceeded considerably due to extra ammonium being deposited from the sea.	Kooijman et al., 2016



Table 2. (continued).

Ecosystem Type/Species	Location	Approach	Ecological Impact	Effects Indicator	Comments	Reference
Woody plant leaves	China		Higher leaf N concentrations correspond to high atmospheric N deposition.		981 observations in unfertilized and non-agricultural ecosystems	Liu et al., 2013
<i>Trifolium repens</i>		N-fixing experiments	N <sub>2</sub> fixation can be switched off when mineral N in soil is sufficient.			Macduff et al., 1996
Calcifugous grasslands	UK	Long-term experimental N addition	Decline in mosses after 3 years			Morecroft et al., 1994
Unimproved hay meadow	Tandham Moor, Somerset	Low N addition (25 kg N ha <sup>-1</sup> yr <sup>-1</sup> )	Significant decline in percent cover of <i>P. lanceolata</i>			Mountford et al., 1993; Mountford et al., 1994; Kirkham et al., 1996
National Parks	Greater Yellowstone Area	Estimation of total N critical loads and development of atmospheric N deposition maps	Lowest CL and highest exceedances were located in high elevation basins with limited vegetation, exposed bedrock, and steep slopes		<ul style="list-style-type: none"> <li>Wet and dry deposition</li> <li>CL for nutrient enrichment in aquatic ecosystems range from <math>&lt; 1.5 \pm 1.0</math> kg N ha<sup>-1</sup> yr<sup>-1</sup> to <math>&gt; 4.0 \pm 1.0</math> kg N ha<sup>-1</sup> yr<sup>-1</sup></li> </ul>	Nanus et al., 2017
Forest (30 stands: sugar maple, red maple, American beech, yellow birch, paper birch, red spruce, balsam fir, eastern hemlock)	White Mountain, New Hampshire				<ul style="list-style-type: none"> <li>Foliar chemistry and soil N status provided positive feedback</li> <li>Good soil C:N ratio prediction accuracy from canopy lignin:N ratio</li> </ul>	Ollinger et al., 2002
Acidic iron-poor dunes	Danish coast		High grass encroachment.			Ovesen, 2001
<i>Calluna vulgaris</i> and <i>Erica cinerea</i>		Testing if vascular plants are feasible bio-indicators of atmospheric N deposition	Indicators of elevated N deposition and CL exceedance described and evaluated	Tissue N content		Pitcairn et al., 1998; Pitcairn et al., 2001; Pitcairn et al., 2003
Ectohydric mosses	UK	N deposition	Loss of biodiversity	<ul style="list-style-type: none"> <li>Tissue N content of mosses excellent for sites where deposition is dominated by NH<sub>3</sub></li> <li>Tissue N content can be useful for identifying s</li> </ul>	5 km × 5 km spatial resolution Spatial variability in upland areas related to cloud cover	Pitcairn et al., 2006

Table 2. (continued).

Ecosystem Type/Species	Location	Approach	Ecological Impact	Effects Indicator	Comments	Reference
<i>Calluna vulgaris</i>		Testing if vascular plants are feasible bio-indicators of atmospheric N deposition		vegetation exposed to large concentrations of $\text{NH}_4^+$ or $\text{NO}_3^-$ such as hilltops and wind exposed woodland edges	Tissue N appears to be more closely linked to rainfall $\text{NH}_4^+$ or $\text{NO}_3^-$ concentrations than deposition. Urban-rural gradient	Power and Collins, 2010
<i>Calluna vulgaris</i>	UK	N addition ( $15.4 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ )	Increase in growth rates. Long-term increase in flower formation.		Lowland heathland	Power <i>et al.</i> , 1995; Power <i>et al.</i> , 2006
	UK	N deposition		<ul style="list-style-type: none"> <li>Mineral N leaching for high N deposition systems (midpoint indicator)</li> <li>Tissue N concentration for low N deposition systems (midpoint indicator)</li> </ul>	Metrics paper	Rowe <i>et al.</i> , 2016
Bog system ( <i>Sphagnum capillifolium</i> , <i>Cladonia</i> spp. and <i>Calluna vulgaris</i> )		Ammonia release experiment	Loss of species richness – all disappeared at higher ammonia concentrations			Shepherd <i>et al.</i> , 2009
<i>Calluna vulgaris</i> , bryophytes, and sedge	UK	Gaseous $\text{NH}_3$ deposition	<ul style="list-style-type: none"> <li>Reduction and almost total loss in <i>Calluna vulgaris</i>, <i>Sphagnum capillifolium</i>, and <i>Cladonia portentosa</i> cover after 3 yrs</li> <li>Increase in sedge cover</li> </ul>		<ul style="list-style-type: none"> <li>Effects due to direct foliar uptake and interaction with abiotic and biotic stresses not effects on soil.</li> <li>Effects on cover of sensitive species happen at lower <math>\text{NH}_3</math> concentrations for longer exposure times</li> </ul>	Sheppard <i>et al.</i> , 2011
<i>Calluna vulgaris</i> , bryophytes, and sedge	UK	Wet $\text{NH}_4$ deposition	Significant increase in <i>Calluna vulgaris</i> cover after 5 yrs			Sheppard <i>et al.</i> , 2011

Table 2. (continued).

Ecosystem Type/Species	Location	Approach	Ecological Impact	Effects Indicator	Comments	Reference
Grasslands			<ul style="list-style-type: none"> <li>Reduction in <i>Sphagnum capillifolium</i> and <i>Cladonia portentos</i> cover after 5 yrs</li> </ul>			Skiba et al., 1989
Herbaceous terrestrial and wetland ecosystems	China, Europe, the Americas	Long-term N enrichment experiments	<ul style="list-style-type: none"> <li>Acidification of soil and metal availability</li> <li>Reduced plant species richness by an average 16%</li> </ul>	<ul style="list-style-type: none"> <li>Presence or absence of indicator species found to not be useful as an indicator of N deposition impacts</li> <li>Percentage cover of individual species suggested to have more promise as an indicator: <i>L. corniculatus</i>, <i>C. rotundifolia</i>, <i>H. splendens</i>, <i>P. lanceolata</i> and <i>T. tamariscinum</i></li> </ul>	189 field experiments, 51 sites. More pronounced reduction in China than Europe and the Americas	Soons et al., 2016
Calcifugous Grasslands	England, Scotland, Wales (National gradient)		Acidification of soil and metal availability. Decline in forb and pleurocarpous moss richness.			Stevens et al., 2006; Stevens et al., 2009b
Lichen taxa	UK	N deposition	Declines in lichen taxa	Terricolous lichens as indicators	5 km × 5 km scale	Stevens et al., 2012
Grasslands, forests, heathlands, and wetlands		N reduction	Recovery from N deposition likely to be a slow process		Review paper	Stevens, 2016
Forests	Europe	Atmospheric N deposition	<ul style="list-style-type: none"> <li>Indicators of elevated N deposition and CL exceedance described and evaluated</li> <li>Slight shift towards nitrolytic species at high N deposition</li> </ul>			Sutton et al., 2001; Sutton et al., 2004
Saltmarshes			Increases vegetation growth			van Dobben and de Vries, 2010
Old sessile oak woodlands	Ireland	N deposition	<ul style="list-style-type: none"> <li>Increase in nitrophiles with higher N deposition</li> <li>Community change point of 13.2 kg N ha<sup>-1</sup> yr<sup>-1</sup> recommended</li> </ul>	Indicator species: <i>Quercus petraea</i> – <i>Luzula sylvatica</i>	1667 10 m × 10 m relevés	Van Wijnen and Bakker, 1999 Wilkins and Aherne, 2016
					Total N deposition not correlated with soil pH	

Table 2. (*continued*).

Ecosystem Type/Species	Location	Approach	Ecological Impact	Effects Indicator	Comments	Reference
Northern hardwood forests (Sugar Maple dominant)	Great Lakes, North America	Wet chemistry and Fourier transform infrared spectroscopy (FTIR)	Decomposition is inhibited by N enrichment	Concentration of acid-insoluble fraction (AIF)	3-yr study	Xia <i>et al.</i> , 2017
<i>Cinnamomum Camphora</i> and <i>Pinus massoniana</i>	Guiyang, SW China	N concentration and $\delta^{15}\text{N}$ values in soils and collected leaves to see feasibility of tall arbors as bio-indicators	$\delta^{15}\text{N}$ values in leaves found to be lower than those of soil. $\delta^{15}\text{N}$ values of Masson pine leaves lower than camphor leaves.		Linear increase in N in leaves with atmospheric N deposition. Observed differences between urban and rural areas showing different N inputs. Higher than critical load.	Xu <i>et al.</i> , 2017
<b>Water</b> Freshwaters		Acidity due to N and decrease in S	Increases dissolved organic carbon (DOC) concentrations (causing brown colouring)			Butterbach-Bahl and Dannemann, 2011
Freshwaters		Acidification of freshwater	Loss or damage to fish populations. Fish reproductive capability include effects on egg production, hatching success and on physiological parameters such as osmo-regulation and gill function			Donaghy and Verspoor, 1997; Kroglund <i>et al.</i> , 2007
	Norway		Health of Atlantic salmon in rivers. Health of brown trout in lakes.		Review paper on critical loads and dose-response relationships. Includes several tables on exceedance indications	de Vries <i>et al.</i> , 2010
						Lien <i>et al.</i> , 1996; Lydersen <i>et al.</i> , 2004; Kroglund <i>et al.</i> , 2007

Table 2. (continued).

Ecosystem Type/Species	Location	Approach	Ecological Impact	Effects Indicator	Comments	Reference
Freshwater		Acidification	Changes in changes in epilithic diatoms, aquatic macrophytes, macroinvertebrate communities, and fish and bird populations			Ormerod and Durance, 2009
**Fresh water and coastal regions			Noxious and toxic algal blooms, increased turbidity, and disruption of ecosystem functioning, shifts in food webs and loss of fish stocks		Review paper	Rabalais, 2002
Freshwaters		Eutrophication	Alteration of aquatic food-webs and fish populations			Smith and Schindler, 2009
		Winter nitrate concentration	Reduced macrophyte species richness			Van der Molen et al., 1998
N limited lakes and Freshwaters		N deposition	High algal growth			Van der Molen et al., 1998; Camargo and Alonso, 2006
<b>Biodiversity</b>						
Butterflies (grassland, heathland, woodland, and farmland)	The Netherlands	N deposition	<ul style="list-style-type: none"> <li>Induces excess early growth of plants</li> <li>Changes quality of food available for larvae</li> <li>Reduction in butterfly biodiversity quality</li> <li>Lack of management causes open bare ground unsuitable for habitat</li> <li>Management effects detrimental to lizards</li> </ul>	Butterflies as indicators	Index of butterfly sensitivity to N created (CNI) 17 yrs of data	Feest et al., 2014
Lizards and heathlands			Encroachment of <i>Calamagrotis</i> caused severe decline in carabids		Response to Maes et al., 2017	Jones et al., 2017
Lizards and heathlands			Encroachment of <i>Calamagrotis</i> caused severe decline in carabids		Comment on Jones et al. paper	Maes et al., 2017
Non-calcareous dune grasslands	Ameland and Terschelling, the Netherlands	Increased N deposition causing encroachment	Encroachment of <i>Calamagrotis</i> caused severe decline in carabids		Effect of grazed and non-grazed grassland on carabid assemblages	Nijssen et al., 2001
**Fauna		Increased N deposition	Direct and indirect effects on fauna		<ul style="list-style-type: none"> <li>Review paper on causal relationship between N deposition-fauna insufficient</li> </ul>	Nijssen et al., 2017

Table 2. (*continued*).

Ecosystem Type/Species	Location	Approach	Ecological Impact	Effects Indicator	Comments	Reference
Heathlands and invertebrates (Diptera and carabid beetles)	The Netherlands	Soil chemistry, plant chemistry, fauna sampling	<ul style="list-style-type: none"> <li>Vegetation N:P ratio negatively linked to invertebrate density and species richness</li> <li>Found to impact fauna more strongly than vegetation structure and plant species richness</li> <li>Soil acidity and P-availability also main causes for loss of habitat structure and plant species richness</li> </ul>	Vegetation N:P ratio	<ul style="list-style-type: none"> <li>Suggest importance of biogeochemistry and ecological stoichiometry</li> </ul>	Vogels <i>et al.</i> , 2016
Butterflies (56 resident species)	The Netherlands	N deposition	<ul style="list-style-type: none"> <li>Butterfly species traits strongly dependent on N availability</li> </ul>	Butterflies and CNI		WallisDeVries and van Swaay, 2016

(Bobbink *et al.*, 2010; Lu *et al.*, 2010; van Dobben and de Vries, 2010; Fenn *et al.*, 2011; Lu *et al.*, 2011; Lu *et al.*, 2014; de Vries *et al.*, 2015; Wilkins and Aherne, 2016; Du, 2017; Xia *et al.*, 2017). In sand dunes, effects include decalcification of topsoil, grass encroachment, changes in seed germination, and changes in species richness and composition (Ovesen, 2001; Jones *et al.*, 2004; Field *et al.*, 2014; Aggenbach *et al.*, 2016; Bird and Choi, 2016; Kooijman *et al.*, 2016).

Impacts from N enrichment on aquatic ecosystems include increased dissolved organic carbon (DOC) concentrations (Butterbach-Bahl and Dannenmann, 2011), loss or damage to fish populations due to changes in the capabilities in fish to reproduce (Donaghy and Verspoor, 1997; Kroglund *et al.*, 2008), and impacts to the health of fish in rivers (Atlantic salmon) and lakes (brown trout) (Lien *et al.*, 1996; Lydersen *et al.*, 2004; Kroglund *et al.*, 2008), changes in aquatic communities, such as epilithic diatoms, aquatic macrophytes, macroinvertebrates, and fish and bird populations (Van der Molen *et al.*, 1998; Ormerod and Durance, 2009; Smith and Schindler, 2009) and increase in algal growth (Van der Molen *et al.*, 1998; Camargo and Alonso, 2006; Hobbs *et al.*, 2010; Baron *et al.*, 2011; Fenn *et al.*, 2011).

Deposition of N on terrestrial and aquatic ecosystems has also resulted in direct and indirect effects on animal biodiversity due to changes in habitat structure and function (Nijssen *et al.*, 2001; Feest *et al.*, 2014; Jones *et al.*, 2016; Vogels *et al.*, 2016; Maes *et al.*, 2017), including declines in carabids and dipterans (Nijssen *et al.*, 2001; Vogels *et al.*, 2016), changes in habitats for lizards (Jones *et al.*, 2016), infestations of heather beetles (Berdowski and Zeilinga, 1983; Berdowski, 1993), and changes in butterfly species traits (Feest *et al.*, 2014; WallisDeVries and van Swaay, 2016). Other reported impacts on fauna due to N deposition include changes in food diversity, quality, and abundance, as well as decreased reproduction (Nijssen *et al.*, 2017).

#### Effects Indicators of Nitrogen Deposition

Impacts of nitrogen are widely assessed using the percent cover of individual plant species (Soons *et al.*, 2016) and the presence of an indicator species (Wilkins and Aherne, 2016). Useful plant indicators include the graminoid:forb ratio (Stevens *et al.*, 2006; Stevens *et al.*, 2009b) and epiphytic lichen and terricolous lichen taxa (Stevens *et al.*, 2012) with shifts in lichen communities occurring above a critical load of 5 kg N ha<sup>-1</sup> yr<sup>-1</sup> (Stevens *et al.*, 2012). Butterflies can be effective indicators for N deposition on faunal biodiversity, with an index of butterfly sensitivity to N having been created using 25 years of data based on changes in butterfly abundance (Feest *et al.*, 2014; WallisDeVries and van Swaay, 2016).

Tissue and litter N content and vegetation N:P ratio have been shown to be good biomonitors for shrubs and mosses and other organisms (Pitcairn *et al.*, 1998; Hicks *et al.*, 2000; Conti and Cecchetti, 2001; Pitcairn *et al.*, 2001; Pitcairn *et al.*, 2002; Pitcairn *et al.*, 2003; Mitchell *et al.*, 2004; Pitcairn *et al.*, 2006; McNeil *et al.*, 2007; Edmondson *et al.*, 2010; Caporn *et al.*, 2014; Harmens *et al.*, 2015; Rowe *et al.*, 2016; Vogels *et al.*, 2016; Du, 2017).

Table 3. Impacts of Acidifying Pollutant: Sulfur (S).

Ecosystem Type / Species	Location	Approach	Ecological Impact	Effects Indicators	Comments	Reference
Terrestrial ecosystems Ginkgoales, bennettites, and conifers	Astartekløft, East Greenland	Physiognomic leaf analysis on well-preserved fossil leaves	Modified leaf physiognomy (increase in leaf roundness)		Regular exposure to volcanic ash	Bacon et al., 2013
Fruticose lichens ( <i>Anaptychia ciliaris</i> , <i>Evernia prunastri</i> , and <i>Ramalina farinacea</i> ) <i>Pinus halepensis</i>	Campi Flegrei, Southern Italy	Cuticle micromorphology on epicuticular and epistomatal waxes	Degradation of epicuticular and epistomatal waxes as fusion of the wax structures Aquatic and terrestrial ecosystems acidified Smaller leaves that did not persist; distinct raised areas of cuticle surrounding stomata appeared; surface waxes altered; blistering of the cuticle occurred; and the stomatal complex became distorted		Influence of volcanic gases. Method shows promise	Balaguer and Manrique, 1991  Bartromo et al., 2012
Mesozoic gymnosperm and fern fossil floras	Growth chamber	SO <sub>2</sub> fumigation. Damage structures in leaf epidermis and cuticle			Volcanic SO <sub>2</sub> . Method effective	Doney et al., 2007 Elliott-Kingston et al., 2014
<i>Picea abies</i> and <i>Abies alba</i> tree cores <i>Picea Abies</i>	NE Italy  Sudetes, Southern Poland	Synchrotron radiation  Tree ring widths  SO <sub>2</sub> fumigation	Strongly negative dependence of tree ring widths on fog deposition rates. Acidification and tree dieback. Change in stomatal density and its ratio to the stomatal index		Study area was an ecological disaster site. Most trees >80 yrs old	Fairchild et al., 2009 Godek et al., 2015  Haworth et al., 2010; Haworth et al., 2012 Hogan et al., 2017
<i>Cladia retipora</i> (lichens)	Tasmania, Australia	Isotopic fingerprinting			Suitability for lichen bio-monitoring baseline determined ( $\delta^{34}\text{S}_{\text{mean}} = 0.03 \pm$ $0.05\%$ )	
Forests Grasses	Czech Republic  Belgium	Temporal and spatial trends S fertilization	Spatial pattern change: 18.1– 0.2 g m <sup>-2</sup> yr <sup>-1</sup> (1995–2011) Yield increases		Based on long-term monitoring	Hünová et al., 2014 Mathot et al.,

Table 3. (continued).

Ecosystem Type / Species	Location	Approach	Ecological Impact	Effects Indicators	Comments	Reference
Grasses		Sulfur nutritional diagnostic tool on S fertilization			Indicator for predicting S deficiency using linear relationships	Mathot <i>et al.</i> , 2009
Industrial sources	Fraser Valley and Saturna Island, Canada	Isotopic mass balance approach			Heavily reduced H <sub>2</sub> S from bacterial sulfate ( $\delta^{34}\text{S}$ range = -1.6 – 9‰)	Norman <i>et al.</i> , 2004
Forests	Nacetin, Czech Republic	Throughfall sampling, water and soil sampling	Reversal of nitrogen enrichment of ecosystems, cessation of nitrate leaching, and a major loss of accumulated organic soil carbon and nitrogen stocks.			Oulehle <i>et al.</i> , 2011
Scots pine	Eurasia		Large scale decrease in growth; higher sensitivity to drought			Savva and Berninger, 2010
Grassland sites	UK (68 sites) Europe	Measured surface and subsoil chemistry S fertilization	Acidification from S deposition		Summers of 2002–2003	Stevens <i>et al.</i> , 2006 Tarrasón <i>et al.</i> , 2006
Vegetation and soil	The Netherlands	Vegetation and soil sampling	Shift towards acidophytic species at high S-deposition			van Dobben and de Vries, 2010
Tree cores	NE Italy	Elemental analyser isotope ratio mass spectrometry (EA-IRMS) compared to tree rings	Yield increases			Wynn <i>et al.</i> , 2014
<i>Cinnamomum camphora</i> Camphor tree leaves	Shanghai, China	Synchrotron radiation X-ray fluorescence and X-ray absorption			Successfully used to study S concentration, speciation, and bioaccumulation	Zeng <i>et al.</i> , 2013



Table 4. Impacts of Base Cations.

Ecosystem Type/Species	Location	Approach	Ecological Impacts	Effects Indicator	Comments	Reference
<i>Picea abies</i> , <i>Fagus sylvatica</i> , and <i>Betula pendula</i>	South Sweden	Bulk deposition, throughfall, stemflow, litterfall, above-ground biomass increments and soil solution	<ul style="list-style-type: none"> <li>• Net foliage leaching of Ca, Mg, and K from canopies</li> <li>• Net loss of base cations from soils, especially spruce</li> <li>• Substantial leaching of base cations from foliage</li> </ul>		Soil acidification higher in spruce than in beach and birch	Bergkvist and Folkesson, 1995
Coniferous and deciduous forests	Northern Germany		Magnitude of inhibition of decomposition increases with increases in H <sup>+</sup> in Al concentration	Litter decomposition	Response stronger in coarse than in fine mesh bags	Bredemeier et al., 1990 Ferreira and Guérôld, 2017
Leaf litter ( <i>Alnus glutinosa</i> , <i>Acer pseudoplatanus</i> and <i>Fagus sylvatica</i> )	Vosges Mountains, NE France	Leaf litter decomposition response to acidification				
Temperate, boreal, coastal, and alpine forests	Ontario, British Columbia, and Nova Scotia, Canada	STA to estimate base cation weathering rate		Soil silt content suggested as a better predictor of soil weathering rates	General model agreement with data	Koseva et al., 2010
Forested catchments	SW China	Measurements and models	Simulations suggest decrease in soil base saturation and pH (soil degradation) over the last decades	Ca:S ratio		Larssen and Carmichael, 2000
Subtropical/tropical forest	Guangdong Province, China	Long-term N addition	<ul style="list-style-type: none"> <li>• Significant soil acidification</li> <li>• Depleted base cations</li> <li>• Increase in exchangeable H<sup>+</sup> proportion in soil cation pools</li> <li>• Increase in soil CEC</li> <li>• Greatly decreased soil solution pH at 20 cm depth but not at 40 cm depth</li> </ul>		No change in soil exchangeable Al <sup>3+</sup>	Lu et al., 2014
Northern forests	Quebec, Canada	Modelling soil chemical weathering rates			Changes in composition within soil profiles are mostly linear (degree of weathering, <i>k</i> , linear to calculated gain/loss of base cations)	Quimet et al., 2006; Quimet, 2008

Table 4. (continued).

Ecosystem Type/Species	Location	Approach	Ecological Impacts	Effects Indicator	Comments	Reference
Mixed boreal forest (dominated by Norway spruce)	Eastern Finland	Sulphate deposition	Ca <sup>2+</sup> and Mg <sup>2+</sup> leaching with greater leaching during growing season than dormant season	K <sup>+</sup> leaching higher than Ca <sup>2+</sup> and Mg <sup>2+</sup> leaching	3 yr study (Three 50 m × 50 m plots)	Pirainen <i>et al.</i> , 2002
Muskeg peatland, lakes, ponds, and uplands (jack pine and trembling aspen)	Oil sands region, Alberta, Canada	<ul style="list-style-type: none"> <li>• Base cation weathering rates estimated</li> <li>• Deposition in throughfall and open collectors measured</li> </ul>	Potential for soil acidification mitigated by high base cation deposition	(Ca + Mg):S ratio	63 sandy, acid-sensitive, upland forest sites	Whitfield <i>et al.</i> , 2011; Watmough <i>et al.</i> , 2014
Sugar maple	Northern Vermont	<ul style="list-style-type: none"> <li>• Base cation depletion</li> </ul>	Crown dieback in stands with low pH, low Ca and Mg, and high Al			Wilmot <i>et al.</i> , 1995

Empirical critical loads of N have been established for a range of terrestrial ecosystems, based on changes to ecosystem structure and function; a critical load of 15 kg N ha<sup>-1</sup> yr<sup>-1</sup> has been recommended for acidic and calcareous grassland (Bobbink and Hettelingh, 2011; Henry and Aherne, 2014), 15 kg N ha<sup>-1</sup> yr<sup>-1</sup> for calcareous grey dunes and 10 kg N ha<sup>-1</sup> yr<sup>-1</sup> for acidic grey dunes (Kooijman *et al.*, 2016). Similarly, critical levels for atmospheric N are 1 µg NH<sub>3</sub> m<sup>-3</sup> for lichens and bryophytes and 3 µg NH<sub>3</sub> m<sup>-3</sup> for herbaceous species (de Vries *et al.*, 2007; Cape *et al.*, 2009). de Vries *et al.* (2007) suggested critical N concentrations in soil solution for a variety of vegetation types to protect against N leaching, with concentrations ranging from 0.2 mg N L<sup>-1</sup> for lichens and cranberries to 6.5 mg N L<sup>-1</sup> for deciduous forests. Similarly, a general threshold for terrestrial soil C:N ratio of < 20–25:1 has been suggested as a threshold for nitrate leaching (Bähring *et al.*, 2017). Litter N content has also been found to be an effective indicator (White *et al.*, 1996; Pilkington *et al.*, 2005). Bähring *et al.* (2017) found shoot increment to be a more sensitive indicator than soil C:N ratio. Soil pH thresholds are suggested at 3.8 for forbs and 4.5 for acidic grasslands (Field *et al.*, 2014). Polyphenol oxidase (PPO) activity is also another biomarker of N enrichment with little difference above an application of 60 kg ha<sup>-1</sup> yr<sup>-1</sup> on peach trees (Edmondson *et al.*, 2010; Falguera *et al.*, 2012; Caporn *et al.*, 2014).

Indicators in aquatic ecosystems include shifts in the relative abundance of diatom communities, with a threshold for wet N deposition between 1.0 and 1.5 kg N ha<sup>-1</sup> yr<sup>-1</sup> (Baron, 2006; Saros *et al.*, 2011; Nanus *et al.*, 2012; Sheibley *et al.*, 2014); nitrate concentrations, where a threshold of 15 mg L<sup>-1</sup> has been determined for well water (Panno *et al.*, 2006); and the ratio of dissolved inorganic N to total phosphorus (DIN:TP), with a suggested range between 1.5 and 3.4 (Bergström, 2010; Fenn *et al.*, 2011; de Vries *et al.*, 2015).

## PAH DEPOSITION

### Major PAH Species

The major PAH species are generally considered to be the US EPA's 16 PAH priority pollutants, which includes naphthalene, acenaphthylene, acenaphthene, fluorene, phenanthrene, anthracene, fluoranthene, pyrene, chrysene, benz[a]anthracene, benzo[b]fluoranthene, benzo[k]fluoranthene, benzo[a]pyrene, indeno[1,2,3-cd]pyrene, dibenzo[a,h]anthracene, and benzo[ghi]perylene. These PAH species were selected as priority pollutants because they are found in the environment, analytical standards and toxicity information are readily available, and many scientific investigations have focused on these 16 PAHs (Andersson and Achten, 2015). However, recent advances in analytical methods have broadened the range of analyses to include polycyclic aromatic compounds (PACs). PACs consists of the 16 PAHs as well as alkylated PAHs, unsubstituted and alkylated dibenzothiophenes, heterocyclic aromatic compounds, and PAC transformation products, which together comprise thousands of compounds

(Manzano *et al.*, 2017; Wnorowski and Charland, 2017). Although these additional PAC species have been the subject of far fewer studies relative to unsubstituted PAHs, their impacts have been observed in air, water, and wildlife particularly near oil sands mining regions (Kelly *et al.*, 2009; Lundin *et al.*, 2015; Schuster *et al.*, 2015; Zhang *et al.*, 2015; Jariyasopit *et al.*, 2016). Toxicity information on these compounds is still very limited; therefore, the following sections will focus on the impacts and biomonitors of PAHs.

### **Impacts of PAH Deposition**

Impacts of PAH deposition on terrestrial ecosystems include visible leaf damage, chlorosis and necrosis, mesophyll collapse, leaf surface bronzing and silverying (Oguntimehin *et al.*, 2008; Oguntimehin *et al.*, 2010); oxidative stress response (Liu *et al.*, 2009); changes in biomass production (Wild *et al.*, 1992; Kummerová and Kmentová, 2004; Wiczorek and Wiczorek, 2007; Váňová *et al.*, 2009; Ahammed *et al.*, 2012a, b); changes in carbon allocation (Desalme *et al.*, 2011); changes to the soil-plant-rhizosphere system, such as enhancement by arbuscular mycorrhizal fungi and degradation by ectomycorrhiza (Koivula *et al.*, 2004; Joner *et al.*, 2006; Verdin *et al.*, 2006); and negative effects on photosynthesis (Wiczorek and Wiczorek, 2007; Oguntimehin *et al.*, 2008; Oguntimehin *et al.*, 2010; Desalme *et al.*, 2011; Ahammed *et al.*, 2012a, b).

Other effects of PAHs on living organisms include the enrichment of antibiotic resistance bacteria (ARB) in soils (Chen *et al.*, 2017) and liver tumours in aquatic organisms, for example, liver lesions in English sole, mortality of sensitive aquatic invertebrates, and skin ulcers and fin erosion in cod and corkscrew wrasse (Aas *et al.*, 2001). In both humans and animals, PAHs have been found to be carcinogenic, mutagenic, teratogenic, and immunosuppressive (Malmgren *et al.*, 1952; Yasuhira, 1964; Philips *et al.*, 1973; Grimmer *et al.*, 1983; Blanton *et al.*, 1986; Hahon and Booth, 1986; Blanton *et al.*, 1988; Zhao *et al.*, 1990; Dasgupta and Lahiri, 1992; Zhao *et al.*, 1992; ATSDR, 1993; Szczeklik *et al.*, 1994; Boström *et al.*, 2002; Yan *et al.*, 2004; Satya *et al.*, 2012; Rodríguez-Estival *et al.*, 2015; Mętrak *et al.*, 2016) given their potential to form carcinogenic, mutagenic diols and epoxides that react with DNA (Shukla and Upreti, 2009).

### **Effects Indicators of PAH Deposition**

Lichens have been shown to be effective indicators. Levels in lichens represent exposure (deposition) but there is no level associated with negative impacts of PAH deposition. Accumulation of PAH absorption in lichens results in PAH profiles that provide knowledge of both the origin and proximity of the source of the contamination (Migaszewski *et al.*, 2002; Blasco *et al.*, 2006, 2008; Augusto *et al.*, 2009; Shukla and Upreti, 2009; Augusto *et al.*, 2010; Satya *et al.*, 2012; Shukla *et al.*, 2012; Augusto *et al.*, 2013). It should be noted that an indicator refers to qualitative data about an organism (e.g., changes in presence/absence, physiology), whereas a biomonitor includes both quantitative and qualitative data on exposure (deposition). Two other highly used biomonitors are mosses (Ötvös *et al.*, 2004; Wang *et*

*al.*, 2009) and pine needles (Holoubek *et al.*, 2000; Conti and Cecchetti, 2001; Onianwa, 2001; Librando *et al.*, 2002; Migaszewski *et al.*, 2002; Piccardo *et al.*, 2005; Schröter-Kermani *et al.*, 2006; Yang *et al.*, 2007; Hwang and Wade, 2008; Lehndorff and Schwark, 2009; Augusto *et al.*, 2010; Ratola *et al.*, 2010; Sun *et al.*, 2010; Amigo *et al.*, 2011; Ratola *et al.*, 2011, 2012; Tomashuk *et al.*, 2012; Fernández-Varela *et al.*, 2015), with pine needles showing promise as better biomonitors for airborne PAH concentrations and mosses for atmospheric deposition (Oishi, 2013). The high PAH accumulation potential in wild rosemary leaves suggests this species to likely be a more valuable indicator than birch leaves or Scots pine needles, as found in a peat bog comparison study (Mętrak *et al.*, 2016). An effect, known as the forest filter effect, has been proposed to suggest that forests, such as deciduous forest canopies in Canada and Germany, intercept and transfer atmospheric PAHs to the surface (Su *et al.*, 2007). This effect shows the potential for using deciduous forests as biomonitors of PAH deposition. Other terrestrial ecosystem indicators include *Quercus ilex* L. Mediterranean evergreen oak (Librando *et al.*, 2002; De Nicola *et al.*, 2005; Orecchio, 2007; De Nicola *et al.*, 2008, 2011; Papa *et al.*, 2012; Baldantoni *et al.*, 2014), and to a lesser extent *Olea europaea* L. leaves (Baldantoni *et al.*, 2014); *Laurus nobilis* evergreen (Lodovici *et al.*, 1998); *Rhododendron oomurasaki* Azalea (Nakajima *et al.*, 1995); *Ficus benghalensis* leaves (Prajapati and Tripathi, 2008); *Calotropis gigantea* R. Br. leaves (Sharma and Tripathi, 2009); *Plantago* species (Bakker *et al.*, 1999; Bakker *et al.*, 2001); vascular plants (Desalme *et al.*, 2013); and the inner wood and rings of blue spruce trees (Rauert *et al.*, 2017). Biological indicators in animals include moose, grey wolf, and woodland caribou excrement (Lundin *et al.*, 2015) and non-lethal fecal sampling and muscle sampling of nestling tree swallows in the Athabasca Oil Sands (Cruz-Martinez *et al.*, 2015; Fernie *et al.*, 2018). Recent research has shown the potential of a passive dry deposition (PAS-DD) air sampling collector, composed of two parallel plates and a PUF disk, that can be used in the field as a replacement to biological indicators to monitor PAH deposition (Harner *et al.*, 2013; Eng *et al.*, 2014; Jariyasopit *et al.*, 2018).

In aquatic ecosystems, aquatic biomonitors include plants such as *Ceratophyllum demersum* L. and *Typha domingensis* pers (Hassan *et al.*, 2010); sentinel zooplankton, e.g., *Daphnia* (Kurek *et al.*, 2013); spider webs (Rybak and Olejniczak, 2014); fish and invertebrates (Rose *et al.*, 2012); and crabs, such as spider crabs, snow crabs (Hellou *et al.*, 1994), or fresh water crabs *Sesarma bouleengeri* (Salman *et al.*, 2014); and colonial waterbird eggs (Hebert *et al.*, 2011). Biliary fluorescence and DNA adducts have been found to be useful biomonitors for PAH contamination in cod and corkscrew wrasse (Aas *et al.*, 1998; Aas *et al.*, 2000; Aas *et al.*, 2001; Hanson and Larsson, 2008), as well as in perch, northern pike, European eels, plaice, blue mussels, flounder, jackfish, and other fish species (Lin *et al.*, 1996; Ericson *et al.*, 1998; Richardson *et al.*, 2001; Ruddock *et al.*, 2003; Richardson *et al.*, 2004; Schiedek *et al.*, 2006; Vuorinen *et al.*, 2006; Beyer *et al.*, 2010; Nkpa *et al.*, 2013; Ohiozebau *et al.*, 2017).

## MERCURY DEPOSITION

### *Major Hg Species*

There are three forms of atmospheric Hg including gaseous elemental Hg (GEM), gaseous oxidized Hg (GOM), and particulate-bound Hg (PBM). GEM is the dominant form of atmospheric Hg and capable of long range transport with an atmospheric lifetime of 0.5–1 year (Driscoll *et al.*, 2013). GOM and PBM typically have atmospheric lifetimes ranging from hours to weeks (Cole *et al.*, 2014). Hg is bi-directional in nature, with both emission and deposition occurring between the air and the surface (Wright *et al.*, 2016). All three forms of Hg can undergo dry deposition (Wright *et al.*, 2016); however, only GOM and PBM can also undergo wet deposition since oxidized Hg is more water soluble than GEM.

### *Impacts of Mercury Deposition*

Mercury deposited to the Earth can undergo methylation in the presence of sulphate-reducing bacteria (SRB). During this process, oxidized Hg is converted to methylmercury (MeHg), which is more often referred to as monomethylmercury (MMHg) (Gilmour *et al.*, 1992; King *et al.*, 2000). Methylmercury bioaccumulates in biota and is highly toxic. Hg deposition impacts humans indirectly through the consumption of contaminated fish, wildlife, and plants that are contaminated with Hg (Meili, 1997; Meili *et al.*, 2003). The resulting effects are risks to the neurological, immune, and reproductive systems (Harada, 1995; Takeuchi *et al.*, 1996; Myers *et al.*, 1998; Schoeman *et al.*, 2009; Bose-O'Reilly *et al.*, 2010; Fernandes Azevedo *et al.*, 2012; Hong *et al.*, 2012; Rice *et al.*, 2014). Wildlife at highest risk are large predatory fish, fish-eating mammals, and fish-eating birds. Impacts on wildlife include reduced reproduction, changes to egg incubation times, behavioural changes, and neurological problems (Myers *et al.*, 1998; Wolfe *et al.*, 1998; Evers *et al.*, 2005; Scheuhammer *et al.*, 2007; Richard Pilsner *et al.*, 2010; Penglase *et al.*, 2014). The consumption of fish with high levels of Hg is an indirect impact of Hg deposition through bioaccumulation of MeHg in the fish and biomagnification in the food chain. Aquatic ecosystems that are acidic, nutrient-deficient, or have higher levels of dissolved organic matter (DOM) have higher concentrations of MeHg than their counterparts (Gilmour *et al.*, 1992; Watras and Huckabee, 1994; Driscoll *et al.*, 2013; French *et al.*, 2014). The effects of Hg on terrestrial and wildlife ecosystems are fairly well understood (Meili *et al.*, 2003; Evers *et al.*, 2005; Evers *et al.*, 2008; Caldwell *et al.*, 2009). Mercury concentrations in Canada are higher in the eastern part of the country than the west owing to the legacy of high emission sources to the south, the preponderance of acidic lakes and soils (Depew *et al.*, 2013, and references therein), and the retention of Hg in organic soils in lake catchments in eastern Canada (Dennis *et al.*, 2005).

### *Effects Indicators of Mercury Deposition*

Ecosystem indicators that can exacerbate Hg bioaccumulation are total phosphorus (TP), DOC, acid

status, and pH. Critical loads for Hg (and persistent organic pollutants; POPs) are not as well developed as they are for N and S. The concentration threshold used as an indicator of Hg sensitivity for total phosphorus is 0.03 mg L<sup>-1</sup>, with fish having higher Hg below that threshold. DOC indicator values for Arctic tundra lakes range from 4 mg L<sup>-1</sup> to 8 mg C L<sup>-1</sup> (Driscoll *et al.*, 2007; French *et al.*, 2014; Stoken *et al.*, 2016). Ecosystem attributes which are associated with higher concentrations of bioavailable Hg include an ANC less than 100 µeq L<sup>-1</sup>, and a pH less than 6.0. The critical limit for Hg concentration in drinking water is 6 µg L<sup>-1</sup> (WHO, 2011). The Canadian water quality guidelines for the protection of aquatic life are 26 ng L<sup>-1</sup> for inorganic Hg in freshwater, 16 ng L<sup>-1</sup> in marine surface waters and 4 ng L<sup>-1</sup> for MeHg in freshwater (CCME, 2003).

Suitable aquatic Hg biomonitors are walleye and northern pike, with thresholds of 0.16 ppm in prey fish and 0.3 ppm in fish tissue (Simoneau *et al.*, 2005; Drevnick *et al.*, 2007; Kinghorn *et al.*, 2007; Rasmussen *et al.*, 2007; Iwu, 2010; Monson *et al.*, 2011; Åkerblom *et al.*, 2012; Gandhi *et al.*, 2015). A good wildlife Hg biomonitor species is the common loon, where the threshold in loon blood is 3.0 ppm (Evers *et al.*, 2003; Kenow *et al.*, 2003; Burgess *et al.*, 2005; Champoux *et al.*, 2006; Evers *et al.*, 2008). Omnivorous birds, such as the Greylag goose, have been found to have higher levels of Hg in their feathers than birds with other dietary habits (Ahmadpour *et al.*, 2016). Mercury levels in marine birds have been studied extensively (Monteiro and Furness, 1995; Bearhop *et al.*, 2000; Becker *et al.*, 2002; Goodale *et al.*, 2008; Burger *et al.*, 2009; Ackerman *et al.*, 2014; Mashroofeh *et al.*, 2015; Binkowski *et al.*, 2016) with indicator thresholds determined to be 1.3 ppm in the blood of songbirds, and 10.0 ppm in the hair of bats. Hg in colonial waterbird eggs and wood frogs has also been measured (Hebert *et al.*, 2011, 2013; Abbasi *et al.*, 2015). In Canada, the Hg limit is 0.2 ppm for safe fish consumption, whereas elsewhere it is 0.5 ppm (de Vries *et al.*, 2007; de Vries *et al.*, 2015). Hg levels in human urine, blood, and hair can also be used as indicators (Sherlock *et al.*, 1984; Apostoli *et al.*, 2002; Myers *et al.*, 2003; Schober *et al.*, 2003; McDowell *et al.*, 2004; Caldwell *et al.*, 2009).

## TRACE METAL DEPOSITION

### *Major Trace Metal Species*

The 13 major trace metal species considered as priority pollutants by US EPA are Antimony (Sb), Arsenic (As), Beryllium (Be), Cadmium (Cd), Chromium (Cr), Copper (Cu), Lead (Pb), Mercury (Hg), Nickel (Ni), Selenium (Se), Silver (Ag), Thallium (Tl) and Zinc (Zn). These metals are part of the priority pollutants list because of their toxic effects on biota and occurrence in the environment. They also have chemical standards and analytical methods available and are produced in significant quantities (US EPA, 2017).

### *Impacts of Trace Metal Deposition*

The impacts of trace metal deposition are well documented but especially for Pb. Much of the knowledge of the impacts

of trace metals such as aluminum, vanadium, and nickel to terrestrial ecosystems has come from acidification studies as decreased pH mobilises metals causing changes that include changes to the mycorrhiza and fine root systems of plants, chlorosis, dwarfing, and reduced root and shoot growth (Rosseland *et al.*, 1990; Efrogmson *et al.*, 1997; Ewais, 1997; Wang and Liu, 1999; Vachirapatama *et al.*, 2011).

Trace metals can accumulate in the suspended particulates and sediments in aquatic ecosystems causing serious potential risks to the health of ecosystems (Li *et al.*, 2013). Impacts of trace metals to aquatic ecosystems include effects on gill function by copper (Sola *et al.*, 1995; Rajkowska and Protasowicki, 2013), iron, manganese, and zinc (Rajkowska and Protasowicki, 2013); nervous systems by zinc, manganese, and iron (Baatrup, 1991; Takeda *et al.*, 2004); and growth and reproduction rates by lead, mercury, and cadmium (Mance, 1987; Ebrahimi and Taherianfard, 2011). Aluminum has been found to be toxic to fish and invertebrates and can interfere with the metabolic, reproductive, and breathing processes of mammals and birds, with potential links to tumours (Rosseland *et al.*, 1990; Exley *et al.*, 1991; Bast, 1993; Sparling and Lowe, 1996; Slaninova *et al.*, 2014). Cadmium has been linked to changes in growth and reproduction of biota (Eisler, 1985; Gallego *et al.*, 2012) and earthworms (Will and Suter, 1995; Chen *et al.*, 2017). Nickel accumulation has been linked to changes in metabolism, bone densities, growth, and survival in birds (Outridge and Scheuhammer, 1993; Eisler, 1998). Histopathological changes and other effects have been observed in carp due to trace metals (Vinodhini and Narayanan, 2009; Georgieva *et al.*, 2014). In humans, trace metal toxicities include damage to the lungs and nervous systems by Al; damage to the lungs, kidneys, and bones and renal dysfunction by Cd; and damage to the lungs, nasal passages, and skin by Ni (Williams and Burson, 1985; Denkhaus and Salnikow, 2002; Krewski *et al.*, 2007; Trzcinka-Ochocka *et al.*, 2010; Zambelli and Ciurli, 2013).

### **Effects Indicators of Trace Metal Deposition**

Terrestrial biomonitors of trace metals include the use of mycorrhizal fungi (Leyval *et al.*, 1997); *Quercus ilex* L. (Papa *et al.*, 2012); lichens (Garty, 2000; Augusto *et al.*, 2004; Augusto *et al.*, 2007; Augusto *et al.*, 2009); mosses and pine needles (Glooschenko, 1989; Bargagli *et al.*, 1995; Steinnes, 1995; Figueira *et al.*, 2002; Migaszewski *et al.*, 2002; Schilling and Lehman, 2002; Schintu *et al.*, 2005; Shotyk *et al.*, 2014; Singh *et al.*, 2017); bryophytes (Nimis *et al.*, 2002); *taraxacum officinale* dandelions (Kabata-Pendias and Dudka, 1991; Królak, 2003); *Tanacetum vulgare* tansy (Jasion *et al.*, 2013); and seagrass leaves (Govers *et al.*, 2014). Other biomonitors of metal contamination include metal enrichment factors and the geoaccumulation index (Abraham and Parker, 2008; Çevik *et al.*, 2009; Zhang *et al.*, 2009; Lourino-Cabana *et al.*, 2011). Large-scale biomonitoring surveys of trace metal concentrations in moss tissue are carried out across Europe (Eurasia) on a five-year cycle under ICP Vegetation (ICP Vegetation, 2017).

In aquatic ecosystems and wildlife, levels of trace metals

have been measured and used as indicators in the eggs, feathers, and tissues of Canadian geese (Tsipoura *et al.*, 2011); *Parus major* and *Ficedula hypoleuca* bird excrement (Berglund *et al.*, 2015); *Parus major* and *Carduelis sinica* birds (Deng *et al.*, 2007); harp seals (Agusa *et al.*, 2011); *Perna viridus* green-lipped mussels (Phillips, 1985); California mussels (Gordon *et al.*, 1980); *Perna perna* mussels (Bellotto and Miekeley, 2007); several fish species (Jaffar *et al.*, 1995; Atobatele and Olutona, 2015); earthworms (Morgan and Morgan, 1988; Stürzenbaum *et al.*, 2004; Hirano and Tamae, 2011; Chen *et al.*, 2017); and sentinel terrestrial animals, such as deer mice and meadow voles (Rodríguez-Estival and Smits, 2016). Critical limits for dissolved trace metal concentrations in drinking water include 3 µg L<sup>-1</sup> for Cd, 10 µg L<sup>-1</sup> for Pb, 2000 µg L<sup>-1</sup> for Cu, and 0.0078 µg L<sup>-1</sup> for Zn (de Vries *et al.*, 2007; WHO, 2011).

## **OZONE DEPOSITION**

### **Impacts of Ozone Deposition**

The impact of high ozone concentration on vegetation was traditionally considered as an exposure effect (concentration-based approach) (Fuhrer *et al.*, 1997). In recent decades, the accumulated stomatal flux (dose-based approach) was considered to be more suitable because ozone damage to vegetation is caused by injury to internal plant tissue and subsequent reduction in photosynthesis, plant growth, and productivity (Musselman *et al.*, 1994; WHO, 2000; Filella *et al.*, 2005; Ferretti *et al.*, 2007). An example of the comparison of the different approaches was shown in Zhang *et al.* (2006). This section focuses on effects of O<sub>3</sub> deposition on vegetation. Ozone is a highly reactive oxygen species and impacts human and animal health, as well as terrestrial ecosystems (Chappelka and Samuelson, 1998; Ashmore, 2005; Ashmore *et al.*, 2006; Emberson and Buker, 2011; Mills *et al.*, 2011; Fuhrer *et al.*, 2016; Bergmann *et al.*, 2017, and references therein). In mammals, significant impacts on respiratory health include reductions in lung function and changes in lung structure (McKee and Rodriguez, 1993; Chen *et al.*, 2007; Poursafa *et al.*, 2011; Bergmann *et al.*, 2017). Impacts to terrestrial ecosystems include decreased net photosynthesis in trees (Wittig *et al.*, 2009), decreased tree biomass (Karlsson *et al.*, 2003; Wittig *et al.*, 2009), decreased grassland productivity (Volk *et al.*, 2006), reduced flowering and bulb growth in woodland ground flora species (Keelan, 2007), plant cell death (Vainonen and Kangasjärvi, 2015), reduced cover of grass species (Thwaites *et al.*, 2006), decreased yields of rice and winter wheat (Feng *et al.*, 2003), agricultural yield losses (Van Dingenen *et al.*, 2009), changes to litter (Fuhrer *et al.*, 2016), changes in the decomposition rate of litter (Lindroth, 2010), among many others (Matyssek and Sandermann Jr., 2003). Rather than provide a table for O<sub>3</sub>, the reader is referred to an extensive review provided by Bergmann *et al.* (2017) that examines over 450 references of plant exposure to O<sub>3</sub> in literature. In addition, a review of impacts of O<sub>3</sub> on terrestrial biodiversity and their downstream effects highlights the aboveground and belowground effects of O<sub>3</sub> exposure (Fuhrer *et al.*, 2016).

### Effects Indicators of Ozone Deposition

Biomonitors for ozone have been studied intensively using field studies, such as the high use of white clover in ozone gardens (Villányi *et al.*, 2008; ICP Vegetation, 2017). There are, however, sometimes difficulties in distinguishing between the impacts being due to O<sub>3</sub> or other pollutants, such as reactive N (Führer *et al.*, 2016). Useful biomonitors for O<sub>3</sub> deposition include the highly-sensitive Myrtaceae and Salicaceae families (Bergmann *et al.*, 2017) and whereas lichens, on the other hand, are not effective biomonitors due to their high O<sub>3</sub> tolerance (Bertuzzi *et al.*, 2013). An interesting ranking of native herbaceous and woody plant species can be found in Bergmann *et al.* (2017). Visible indicators on plants have been found to be leaf browning (Panigada *et al.*, 2009) and necrotic lesions on leaf surfaces, such as on the upper leaf surfaces of tobacco cultivars (Ashmore *et al.*, 1978; Ribas *et al.*, 1998; Klumpp *et al.*, 2006; Kafiatulla *et al.*, 2012; Vainonen and Kangasjärvi, 2015). Other non-visible biomarkers for plants that have been found to show potential are stress transcripts, proteins, and metabolites (Sandermann, 2000). In humans, systemic responses such as inflammation and oxidative stress have been found to be the most useful (Goodman *et al.*, 2015). Recently, an atmospheric critical level concentration for O<sub>3</sub> of 80 ppb was suggested for two Mediterranean trees, although this was performed under controlled conditions and has yet to be tested in the field (Fusaro *et al.*, 2017). Interestingly, in this study, N deposition was observed to counter the effects of O<sub>3</sub> deposition (Fusaro *et al.*, 2017).

### CONCLUSIONS AND RECOMMENDATIONS

The ecosystem and human health impacts of the atmospheric deposition of various pollutants, and the indicators and biomonitors that have been found to be effective for these pollutants are summarized in this review paper. The impacts of the deposition of acidifying pollutants, eutrophying N, PAHs, Hg, Pb, and O<sub>3</sub> have been well-documented, whereas less is known about the impacts of alkylated and heterocyclic PAC and trace metals, such as iron and manganese. Further studies on possible biomonitors for PAC and O<sub>3</sub> are needed. Many studies have been tested under controlled conditions only. Future work should include testing suggested effects indicators, such as the critical load of O<sub>3</sub> on Mediterranean trees, in the field. The current understanding of effects and indicators is due to close collaboration between various research communities. This integration is essential to addressing environmental problems. For example, it would be interesting to explore the possibility of extending the use of the STressor-Ecological Production function-final ecosystem goods and Services (STEPS) Framework, which has been applied to aquatic eutrophication (Rhodes *et al.*, 2017), to other chemical species, such as PAHs, Hg, and trace metals. Recent success by Fernie *et al.* (2018) and Cruz-Martinez *et al.* (2015) on fecal and muscle sampling of PAHs and other contaminants in nesting tree swallows is being built upon with continuing research on toxicity in these birds. Perhaps expanding the list of chemical species, such as adding Hg,

in these fecal samples could be performed concurrently to expand our knowledge of these wild birds and the effects of the Alberta Oil Sands on them and to develop this species as a potential biomonitor for future studies.

### ACKNOWLEDGEMENTS

This work was supported by the Climate Change and Air Pollutants program of Environment and Climate Change Canada and by the Province of Alberta through the Oil Sands Monitoring Program.

### REFERENCES

- Aas, E., Baussant, T., Balk, L., Liewenborg, B. and Andersen, O.K. (2000). PAH metabolites in bile, cytochrome P4501A and DNA adducts as environmental risk parameters for chronic oil exposure: A laboratory experiment with Atlantic cod. *Aquat. Toxicol.* 51: 241–258.
- Aas, E., Beyer, J. and Goksøyr, A. (1998). PAH in fish bile detected by fixed wavelength fluorescence. *Mar. Environ. Res.* 46: 225–228.
- Aas, E., Beyer, J., Jonsson, G., Reichert, W. and Andersen, O. (2001). Evidence of uptake, biotransformation and DNA binding of polyaromatic hydrocarbons in Atlantic cod and corkwing wrasse caught in the vicinity of an aluminium works. *Mar. Environ. Res.* 52: 213–229.
- Abbasi, N.A., Jaspers, V.L.B., Chaudhry, M.J.I., Ali, S. and Malik, R.N. (2015). Influence of taxa, trophic level, and location on bioaccumulation of toxic metals in bird's feathers: A preliminary biomonitoring study using multiple bird species from Pakistan. *Chemosphere* 120: 527–537.
- Abdel-Shafy, H.I. and Mansour, M.S. (2016). A review on polycyclic aromatic hydrocarbons: Source, environmental impact, effect on human health and remediation. *Egypt. J. Pet.* 25: 107–123.
- Abraham, G. and Parker, R. (2008). Assessment of heavy metal enrichment factors and the degree of contamination in marine sediments from Tamaki Estuary, Auckland, New Zealand. *Environ. Monit. Assess.* 136: 227–238.
- Ackerman, J.T., Eagles-Smith, C.A., Heinz, G., De La Cruz, S.E., Takekawa, J.Y., Miles, A.K., Adelsbach, T.L., Herzog, M.P., Bluso-Demers, J.D. and Demers, S.A. (2014). Mercury in birds of san francisco bay-delta, California: Trophic pathways, bioaccumulation, and ecotoxicological risk to avian reproduction, US Geological Survey.
- Agency for Toxic Substances and Diseases Registry (ATSDR) (1993). *Toxicological profile for polycyclic aromatic hydrocarbons (PAHs)*. U.S. Department of Health and Human Services, Atlanta, USA.
- Aggenbach, C.J., Kooijman, A.M., Fujita, Y., van der Hagen, H., van Til, M., Cooper, D. and Jones, L. (2016). Does atmospheric nitrogen deposition lead to greater nitrogen and carbon accumulation in coastal sand dunes? *Biol. Conserv.* 212: 416–422.
- Agusa, T., Nomura, K., Kunito, T., Anan, Y., Iwata, H.

- and Tanabe, S. (2011). Accumulation of trace elements in harp seals (*Phoca groenlandica*) from Pangnirtung in the Baffin Island, Canada. *Mar. Pollut. Bull.* 63: 489–499.
- Ahamed, G.J., Choudhary, S.P., Chen, S., Xia, X., Shi, K., Zhou, Y. and Yu, J. (2012a). Role of brassinosteroids in alleviation of phenanthrene-cadmium co-contamination-induced photosynthetic inhibition and oxidative stress in tomato. *J. Exp. Bot.* 64: 199–213.
- Ahamed, G.J., Wang, M.M., Zhou, Y.H., Xia, X.J., Mao, W.H., Shi, K. and Yu, J.Q. (2012b). The growth, photosynthesis and antioxidant defense responses of five vegetable crops to phenanthrene stress. *Ecotoxicol. Environ. Saf.* 80: 132–139.
- Ahmadpour, M., Lan-Hai, L., Ahmadpour, M., Hoseini, S.H., Mashrofeh, A. and Binkowski, L.J. (2016). Mercury concentration in the feathers of birds from various trophic levels in Fereydunkenar International wetland (Iran). *Environ. Monit. Assess.* 188: 666.
- Åkerblom, S., Nilsson, M., Yu, J., Ranney, B. and Johansson, K. (2012). Temporal change estimation of mercury concentrations in northern pike (*Esox lucius* L.) in Swedish lakes. *Chemosphere* 86: 439–445.
- Akselsson, C., Westling, O., Sverdrup, H., Holmqvist, J., Thelin, G., Uggla, E. and Malm, G. (2007). Impact of harvest intensity on long term base cation budgets in Swedish forest soils. *Water Air Soil Pollut. Focus* 7: 201–210.
- Amigo, J.M., Ratola, N. and Alves, A. (2011). Study of geographical trends of polycyclic aromatic hydrocarbons using pine needles. *Atmos. Environ.* 45: 5988–5996.
- Andersson, J.T. and Achten, C. (2015). Time to say goodbye to the 16 EPA PAHs? Toward an up-to-date use of PACs for environmental purposes. *Polycyclic Aromat. Compd.* 35: 330–354.
- Apostoli, P., Cortesi, I., Mangili, A., Elia, G., Drago, I., Gagliardi, T., Soleo, L., Valente, T., Sciarra, G. and Aprea, C. (2002). Assessment of reference values for mercury in urine: The results of an Italian polycentric study. *Sci. Total Environ.* 289: 13–24.
- Ashmore, M. (2005). Assessing the future global impacts of ozone on vegetation. *Plant Cell Environ.* 28: 949–964.
- Ashmore, M., Toet, S. and Emberson, L. (2006). Ozone – a significant threat to future world food production? *New Phytol.* 170: 201–204.
- Ashmore, M.R., Bell, J.N.B., Reily, C.L. (1978). A survey of ozone levels in the British Isles using indicator plants. *Nature* 276: 813–815.
- Atobatele, O.E. and Olutona, G.O. (2015). Distribution of three non-essential trace metals (Cadmium, Mercury and Lead) in the organs of fish from Aiba Reservoir, Iwo, Nigeria. *Toxicol. Rep.* 2: 896–903.
- Augusto, S., Pinho, P., Branquinho, C., Pereira, M., Soares, A. and Catarino, F. (2004). Atmospheric dioxin and furan deposition in relation to land-use and other pollutants: A survey with lichens. *J. Atmos. Chem.* 49: 53–65.
- Augusto, S., Pereira, M.J., Soares, A. and Branquinho, C. (2007). The contribution of environmental biomonitoring with lichens to assess human exposure to dioxins. *Int. J. Hyg. Environ. Health* 210: 433–438.
- Augusto, S., Máguas, C. and Branquinho, C. (2009). Understanding the performance of different lichen species as biomonitors of atmospheric dioxins and furans: potential for intercalibration. *Ecotoxicology* 18: 1036–1042.
- Augusto, S., Máguas, C., Matos, J., Pereira, M.J. and Branquinho, C. (2010). Lichens as an integrating tool for monitoring PAH atmospheric deposition: A comparison with soil, air and pine needles. *Environ. Pollut.* 158: 483–489.
- Augusto, S., Máguas, C. and Branquinho, C. (2013). Guidelines for biomonitoring persistent organic pollutants (POPs), using lichens and aquatic mosses – A review. *Environ. Pollut.* 180: 330–338.
- Baattrup, E. (1991). Structural and functional effects of heavy metals on the nervous system, including sense organs, of fish. *Comp. Biochem. Physiol. C* 100: 253–257.
- Bacon, K.L., Belcher, C.M., Haworth, M. and McElwain, J.C. (2013). Increased atmospheric SO<sub>2</sub> detected from changes in leaf physiognomy across the Triassic-Jurassic boundary interval of East Greenland. *PLoS One* 8: e60614.
- Bähring, A., Fichtner, A., Ibe, K., Schütze, G., Temperton, V.M., von Oheimb, G. and Härdtle, W. (2017). Ecosystem functions as indicators for heathland responses to nitrogen fertilisation. *Ecol. Indic.* 72: 185–193.
- Bakker, M.I., Vorenhout, M., Sijm, D.T. and Kollöffel, C. (1999). Dry deposition of atmospheric polycyclic aromatic hydrocarbons in three *Plantago* species. *Environ. Toxicol. Chem.* 18: 2289–2294.
- Bakker, M.I., Koerselman, J.W., Tolls, J. and Kollöffel, C. (2001). Localization of deposited polycyclic aromatic hydrocarbons in leaves of *Plantago*. *Environ. Toxicol. Chem.* 20: 1112–1116.
- Balaguer, L. and Manrique, E. (1991). Interaction between sulfur dioxide and nitrate in some lichens. *Environ. Exp. Bot.* 31: 223–227.
- Baldantoni, D., De Nicola, F. and Alfani, A. (2014). Air biomonitoring of heavy metals and polycyclic aromatic hydrocarbons near a cement plant. *Atmos. Pollut. Res.* 5: 262–269.
- Baldigo, B.P., Lawrence, G.B., Bode, R.W., Simonin, H.A., Roy, K.M. and Smith, A.J. (2009). Impacts of acidification on macroinvertebrate communities in streams of the western Adirondack Mountains, New York, USA. *Ecol. Indic.* 9: 226–239.
- Barbier, E.B., Koch, E.W., Silliman, B.R., Hacker, S.D., Wolanski, E., Primavera, J., Granek, E.F., Polasky, S., Aswani, S. and Cramer, L.A. (2008). Coastal ecosystem-based management with nonlinear ecological functions and values. *Science* 319: 321–323.
- Bargagli, R., Brown, D. and Nelli, L. (1995). Metal biomonitoring with mosses: Procedures for correcting for soil contamination. *Environ. Pollut.* 89: 169–175.

- Baron, J.S. (2006). Hindcasting nitrogen deposition to determine an ecological critical load. *Ecol. Appl.* 16: 433–439.
- Baron, J.S., Driscoll, C.T., Stoddard, J.L. and Richer, E.E. (2011). Empirical critical loads of atmospheric nitrogen deposition for nutrient enrichment and acidification of sensitive US lakes. *Bioscience* 61: 602–613.
- Bartirromo, A., Guignard, G., Lumaga, M.R.B., Barattolo, F., Chiodini, G., Avino, R., Guerriero, G. and Barale, G. (2012). Influence of volcanic gases on the epidermis of *Pinus halepensis* Mill. in Campi Flegrei, Southern Italy: A possible tool for detecting volcanism in present and past floras. *J. Volcanol. Geotherm. Res.* 233: 1–17.
- Bast, C.B. (1993). *Toxicity summary for aluminum*. Oak Ridge Reservation Environmental Restoration Program; Oak Ridge, TN, USA.
- Bearhop, S., Waldron, S., Thompson, D. and Furness, R. (2000). Bioamplification of mercury in great skua *Catharacta skua* chicks: The influence of trophic status as determined by stable isotope signatures of blood and feathers. *Mar. Pollut. Bull.* 40: 181–185.
- Becker, P.H., González-Solís, J., Behrends, B. and Croxall, J. (2002). Feather mercury levels in seabirds at South Georgia: Influence of trophic position, sex and age. *Mar. Ecol. Prog. Ser.* 243: 261–269.
- Beem, K.B., Raja, S., Schwandner, F.M., Taylor, C., Lee, T., Sullivan, A.P., Carrico, C.M., McMeeking, G.R., Day, D., Levin, E., Hand, J., Kreidenweis, S.M., Schichtel, B., Malm, W.C. and Collett, J.L., Jr. (2010). Deposition of reactive nitrogen during the Rocky Mountain Airborne Nitrogen and Sulfur (RoMANS) study. *Environ. Pollut.* 158: 862–872.
- Bellotto, V. and Miekeley, N. (2007). Trace metals in mussel shells and corresponding soft tissue samples: A validation experiment for the use of *Perna perna* shells in pollution monitoring. *Anal. Bioanal. Chem.* 389: 769–776.
- Benedict, K.B., Carrico, C.M., Kreidenweis, S.M., Schichtel, B., Malm, W.C. and Collett, J.L., Jr. (2013). A seasonal nitrogen deposition budget for Rocky Mountain National Park. *Ecol. Appl.* 23: 1156–1169.
- Berdowski, J. and Zeilinga, R. (1983). The effect of the Heather Beetle (*Lochmaea suturalis* Thomson) on heather (*Calluna vulgaris* (L.) Hull) as a cause of mosaic patterns in heathlands. *Acta Bot. Need.* 32: 250–251.
- Berdowski, J. (1993). The effect of external stress and disturbance factors on calluna-dominated heathland vegetation. In *Heathlands*, Springer, pp. 85–124.
- Bergkvist, B. and Folkesson, L. (1995). The influence of tree species on acid deposition, proton budgets and element fluxes in south Swedish forest ecosystems. *Ecol. Bull.* 44: 90–99.
- Berglund, Å.M., Rainio, M.J. and Eeva, T. (2015). Temporal trends in metal pollution: Using bird excrement as indicator. *PLoS one* 10: e0117071.
- Bergmann, E., Bender, J. and Weigel, H.J. (2017). Impact of tropospheric ozone on terrestrial biodiversity: A literature analysis to identify ozone sensitive taxa. *J. Appl. Bot. Food Qual.* 90: 83–105.
- Bergström, A.K. (2010). The use of TN:TP and DIN:TP ratios as indicators for phytoplankton nutrient limitation in oligotrophic lakes affected by N deposition. *Aquat. Sci.* 72: 277–281.
- Bertuzzi, S., Davies, L., Power, S., and Tretiach, M. (2013) Why lichens are bad biomonitors of ozone pollution? *Ecol. Indic.* 34: 391–397.
- Beyer, J., Jonsson, G., Porte, C., Krahn, M.M. and Ariese, F. (2010). Analytical methods for determining metabolites of polycyclic aromatic hydrocarbon (PAH) pollutants in fish bile: A review. *Environ. Toxicol. Pharmacol.* 30: 224–244.
- Binkowski, Ł.J., Przystupińska, A. and Wojtaś, W. (2016). Levels of total mercury in tissues of Mallard Drakes from industrialized wetlands area. *Bull. Environ. Contam. Toxicol.* 96: 173–178.
- Bird, E.J. and Choi, Y.D. (2016). Response of native plants to elevated soil nitrogen in the sand dunes of Lake Michigan, USA. *Biol. Conserv.* 212: 398–405.
- Blanton, R.H., Lyte, M., Myers, M.J. and Bick, P.H. (1986). Immunomodulation by polyaromatic hydrocarbons in mice and murine cells. *Cancer Res.* 46: 2735–2739.
- Blanton, R.H., Myers, M.J. and Bick, P.H. (1988). Modulation of immunocompetent cell populations by benzo[*a*]pyrene. *Toxicol. Appl. Pharmacol.* 93: 267–274.
- Blasco, M., Domeño, C. and Nerin, C. (2006). Use of lichens as pollution biomonitors in remote areas: Comparison of PAHs extracted from lichens and atmospheric particles sampled in and around the Somport tunnel (Pyrenees). *Environ. Sci. Technol.* 40: 6384–6391.
- Blasco, M., Domeño, C. and Nerin, C. (2008). Lichens biomonitoring as feasible methodology to assess air pollution in natural ecosystems: Combined study of quantitative PAHs analyses and lichen biodiversity in the Pyrenees Mountains. *Anal. Bioanal. Chem.* 391: 759–771.
- Bobbink, R., Hornung, M. and Roelofs, J.G. (1998). The effects of air-borne nitrogen pollutants on species diversity in natural and semi-natural European vegetation. *J. Ecol.* 86: 717–738.
- Bobbink, R., Hicks, K., Galloway, J., Spranger, T., Alkemade, R., Ashmore, M., Bustamante, M., Cinderby, S., Davidson, E., Dentener, F., Emmett, B., Erisman, J.W., Fenn, M., Gilliam, F., Nordin, A., Pardo, L. and de Vries, W. (2010). Global assessment of nitrogen deposition effects on terrestrial plant diversity: A synthesis. *Ecol. Appl.* 20: 30–59.
- Bobbink, R. and Hettelingh, J.P. (2011). Review and revision of empirical critical loads and dose-response relationships: Proceedings of an Expert Workshop, Noordwijkerhout, 23-25 June 2010. Rijksinstituut voor Volksgezondheid en Milieu RIVM.
- Bose-O'Reilly, S., McCarty, K.M., Steckling, N. and Lettmeier, B. (2010). Mercury exposure and children's health. *Curr. Probl. Pediatr. Adolesc. Health Care* 40: 186–215.
- Boström, C.E., Gerde, P., Hanberg, A., Jernström, B., Johansson, C., Kyrklund, T., Rannug, A., Törnqvist, M., Victorin, K. and Westerholm, R. (2002). Cancer risk



- assessment, indicators, and guidelines for polycyclic aromatic hydrocarbons in the ambient air. *Environ. Health Perspect.* 110: 451.
- Bragazza, L., Freeman, C., Jones, T., Rydin, H., Limpens, J., Fenner, N., Ellis, T., Gerdol, R., Hájek, M. and Hájek, T. (2006). Atmospheric Nitrogen deposition promotes carbon loss from peat bogs. *Proc. Natl. Acad. Sci. U.S.A.* 103: 19386–19389.
- Bredemeier, M., Matzner, E. and Ulrich, B. (1990). Internal and external proton load to forest soils in northern Germany. *J. Environ. Qual.* 19: 469–477.
- Bulger, A.J., Cosby, B.J. and Webb, J.R. (2000). Current, reconstructed past, and projected future status of brook trout (*Salvelinus fontinalis*) streams in Virginia. *Can. J. Fish. Aquat. Sci.* 57: 1515–1523.
- Burger, J., Gochfeld, M., Jeitner, C., Burke, S., Volz, C.D., Snigaroff, R., Snigaroff, D., Shukla, T. and Shukla, S. (2009). Mercury and other metals in eggs and feathers of glaucous-winged gulls (*Larus glaucescens*) in the Aleutians. *Environ. Monit. Assess.* 152: 179–194.
- Burgess, N.M., Evers, D.C. and Kaplan, J.D. (2005). Mercury and other contaminants in common loons breeding in Atlantic Canada. *Ecotoxicology* 14: 241–252.
- Butterbach-Bahl, K. and Dannenmann, M. (2011). Denitrification and associated soil N<sub>2</sub>O emissions due to agricultural activities in a changing climate. *Curr. Opin. Environ. Sustainability* 3: 389–395.
- Caldwell, K.L., Mortensen, M.E., Jones, R.L., Caudill, S.P. and Osterloh, J.D. (2009). Total blood mercury concentrations in the U.S. population: 1999–2006. *Int. J. Hyg. Environ. Health* 212: 588–598.
- Camargo, J.A. and Alonso, A. (2006). Ecological and toxicological effects of inorganic nitrogen pollution in aquatic ecosystems: A global assessment. *Environ. Int.* 32: 831–849.
- Cape, J.N., van der Eerden, L.J., Sheppard, L.J., Leith, I.D. and Sutton, M.A. (2009). Evidence for changing the critical level for ammonia. *Environ. Pollut.* 157: 1033–1037.
- Caporn, S.J., Carroll, J.A., Dise, N.B. and Payne, R.J. (2014). Impacts and indicators of nitrogen deposition in moorlands: Results from a national pollution gradient study. *Ecol. Indic.* 45: 227–234.
- Carroll, J.A., Johnson, D., Morecroft, M., Taylor, A., Caporn, S.J. and Lee, J.A. (2000). The effect of long-term nitrogen additions on the bryophyte cover of upland acidic grasslands. *J. Bryology* 22: 83–89.
- Carter, T.S., Clark, C.M., Fenn, M.E., Jovan, S., Perakis, S.S., Riddell, J., Schaberg, P.G., Greaver, T.L. and Hastings, M.G. (2017). Mechanisms of nitrogen deposition effects on temperate forest lichens and trees. *Ecosphere* 8: e01717.
- Çevik, F., Göksu, M.Z.L., Dericci, O.B. and Fındık, Ö. (2009). An assessment of metal pollution in surface sediments of Seyhan dam by using enrichment factor, geoaccumulation index and statistical analyses. *Environ. Monit. Assess.* 152: 309–317.
- Champoux, L., Masse, D., Evers, D., Lane, O., Plante, M. and Timmermans, S. (2006). Assessment of mercury exposure and potential effects on common loons (*Gavia immer*) in Québec. *Hydrobiologia* 567: 263–274.
- Chappelka, A.H. and Samuelson, L.J. (1998). Ambient ozone effects on forest trees of the eastern United States: A review. *New Phytol.* 139: 91–108.
- Chen, B., He, R., Yuan, K., Chen, E., Lin, L., Chen, X., Sha, S., Zhong, J., Lin, L. and Yang, L. (2017). Polycyclic aromatic hydrocarbons (PAHs) enriching antibiotic resistance genes (Args) in the soils. *Environ. Pollut.* 220: 1005–1013.
- Chen, C., Arjomandi, M., Balmes, J., Tager, I. and Holland, N. (2007). Effects of chronic and acute ozone exposure on lipid peroxidation and antioxidant capacity in healthy young adults. *Environ. Health Perspect.* 115: 1732.
- Clivot, H., Cornut, J., Chauvet, E., Elger, A., Poupin, P., Guerold, F. and Pagnout, C. (2014). Leaf-associated fungal diversity in acidified streams: Insights from combining traditional and molecular approaches. *Environ. Microbiol.* 16: 2145–2156.
- Cole, A.S., Steffen, A., Eckley, C.S., Narayan, J., Pilote, M., Tordon, R., Graydon, J.A., St Louis, V.L., Xu, X. and Branfireun, B.A. (2014). A survey of mercury in air and precipitation across Canada: patterns and trends. *Atmosphere* 5: 635–668.
- Conti, M.E. and Cecchetti, G. (2001). Biological monitoring: Lichens as bioindicators of air pollution assessment—A review. *Environ. Pollut.* 114: 471–492.
- Cruz-Martinez, L., Fernie, K.J., Soos, C., Harner, T., Getachew, F. and Smits, J.E.G. (2015). Detoxification, endocrine, and immune responses of tree swallow nestlings naturally exposed to air contaminants from the Alberta oil sands. *Sci. Total Environ.* 502: 8–15.
- Dahlman, L., Persson, J., Palmqvist, K. and Näsholm, T. (2004). Organic and inorganic nitrogen uptake in lichens. *Planta* 219: 459–467.
- Dangles, O. and Guérol, F. (2001). Linking shredders and leaf litter processing: Insights from an acidic stream study. *Int. Rev. Hydrobiol.* 86: 395–406.
- Dasgupta, P. and Lahiri, T. (1992). Alteration of brain catecholamines during growth of benzo(a)pyrene induced murine fibrosarcoma. *Neoplasma* 39: 163–165.
- De Nicola, F., Maisto, G., Prati, M.V. and Alfani, A. (2005). Temporal variations in PAH concentrations in *Quercus ilex* L. (holm oak) leaves in an urban area. *Chemosphere* 61: 432–440.
- De Nicola, F., Maisto, G., Prati, M. and Alfani, A. (2008). Leaf accumulation of trace elements and polycyclic aromatic hydrocarbons (PAHs) in *Quercus ilex* L. *Environ. Pollut.* 153: 376–383.
- De Nicola, F., Claudia, L., MariaVittoria, P., Giulia, M. and Anna, A. (2011). Biomonitoring of PAHs by using *Quercus ilex* leaves: Source diagnostic and toxicity assessment. *Atmos. Environ.* 45: 1428–1433.
- de Vries, W. (1993). Average Critical Loads for Nitrogen and Sulfur and Its Use in Acidification Abatement Policy in the Netherlands. *Water Air Soil Pollut.* 68: 399–434.

- de Vries, W., van Grinsven, J.J.M., van Breemen, N., Leeters, E.E.J.M. and Jansen, P.C. (1995). Impacts of acid deposition on concentrations and fluxes of solutes in acid sandy soils in the Netherlands. *Geoderma* 67: 17–43.
- de Vries, W., Van der Salm, C., Reinds, G. and Erisman, J. (2007). Element fluxes through European forest ecosystems and their relationships with stand and site characteristics. *Environ. Pollut.* 148: 501–513.
- de Vries, W., Wamelink, G., Van Dobben, H., Kros, J., Reinds, G., Mol-Dijkstra, J., Smart, S., Evans, C., Rowe, E. and Belyazid, S. (2010). Use of dynamic soil–vegetation models to assess impacts of nitrogen deposition on plant species composition: an overview. *Ecol. Appl.* 20: 60–79.
- de Vries, W., Du, E. and Butterbach-Bahl, K. (2014). Short and long-term impacts of nitrogen deposition on carbon sequestration by forest ecosystems. *Curr. Opin. Environ. Sustainability* 9: 90–104.
- de Vries, W., Hettelingh, J.P. and Posch, M. (2015). *Critical loads and dynamic risk assessments: Nitrogen, acidity and metals in terrestrial and aquatic ecosystems*. Springer, Dordrecht, Netherlands.
- Deng, H., Zhang, Z., Chang, C. and Wang, Y. (2007). Trace metal concentration in Great Tit (*Parus major*) and Greenfinch (*Carduelis sinica*) at the Western Mountains of Beijing, China. *Environ. Pollut.* 148: 620–626.
- Denkhaus, E. and Salnikow, K. (2002). Nickel essentiality, toxicity, and carcinogenicity. *Crit. Rev. Oncol. Hematol.* 42: 35–56.
- Dennis, I.F., Clair, T.A., Driscoll, C.T., Kamman, N., Chalmers, A., Shanley, J., Norton, S.A. and Kahl, S. (2005). Distribution patterns of mercury in lakes and rivers of northeastern North America. *Ecotoxicology* 14: 113–123.
- Depew, D.C., Burgess, N.M., Anderson, M.R., Baker, R., Bhavsar, S.P., Bodaly, R., Eckley, C.S., Evans, M.S., Gantner, N. and Graydon, J.A. (2013). An overview of mercury concentrations in freshwater fish species: A national fish mercury dataset for Canada. *Can. J. Fish. Aquat. Sci.* 70: 436–451.
- Desalme, D., Binet, P., Bernard, N., Gilbert, D., Toussaint, M.L. and Chiapusio, G. (2011). Atmospheric phenanthrene transfer and effects on two grassland species and their root symbionts: A microcosm study. *Environ. Exp. Bot.* 71: 146–151.
- Desalme, D., Binet, P. and Chiapusio, G.v. (2013). Challenges in tracing the fate and effects of atmospheric polycyclic aromatic hydrocarbon deposition in vascular plants. *Environ. Sci. Technol.* 47: 3967–3981.
- Dirkse, G. and Martakis, G. (1992). Effects of fertilizer on bryophytes in Swedish experiments on forest fertilization. *Biol. Conserv.* 59: 155–161.
- Dirnböck, T., Grandin, U., Bernhardt-Römermann, M., Beudert, B., Canullo, R., Forsius, M., Grabner, M.T., Holmberg, M., Kleemola, S. and Lundin, L. (2014). Forest floor vegetation response to nitrogen deposition in Europe. *Global Change Biol.* 20: 429–440.
- Donaghy, M.J. and Verspoor, E. (1997). Egg survival and timing of hatch in two Scottish Atlantic salmon stocks. *J. Fish Biol.* 51: 211–214.
- Doney, S.C., Mahowald, N., Lima, I., Feely, R.A., Mackenzie, F.T., Lamarque, J.F. and Rasch, P.J. (2007). Impact of anthropogenic atmospheric nitrogen and sulfur deposition on ocean acidification and the inorganic carbon system. *Proc. Natl. Acad. Sci. U.S.A.* 104: 14580–14585.
- Drevnick, P.E., Canfield, D.E., Gorski, P.R., Shinneman, A.L., Engstrom, D.R., Muir, D.C., Smith, G.R., Garrison, P.J., Cleckner, L.B. and Hurley, J.P. (2007). Deposition and cycling of sulfur controls mercury accumulation in Isle Royale fish. *Environ. Sci. Technol.* 41: 7266–7272.
- Driscoll, C.T., Han, Y.J., Chen, C.Y., Evers, D.C., Lambert, K.F., Holsen, T.M., Kamman, N.C. and Munson, R.K. (2007). Mercury contamination in forest and freshwater ecosystems in the northeastern United States. *BioScience* 57: 17–28.
- Driscoll, C.T., Lawrence, G.B., Bulger, A.J., Butler, T.J., Cronan, C.S., Eagar, C., Lambert, K.F., Likens, G.E., Stoddard, J.L. and Weathers, K.C. (2001). Acidic Deposition in the Northeastern United States: Sources and Inputs, Ecosystem Effects, and Management Strategies: The effects of acidic deposition in the northeastern United States include the acidification of soil and water, which stresses terrestrial and aquatic biota. *BioScience* 51: 180–198.
- Driscoll, C.T., Mason, R.P., Chan, H.M., Jacob, D.J. and Pirrone, N. (2013). Mercury as a global pollutant: Sources, pathways, and effects. *Environ. Sci. Technol.* 47: 4967–4983.
- Du, E. (2017). Integrating species composition and leaf nitrogen content to indicate effects of nitrogen deposition. *Environ. Pollut.* 221: 392–397.
- Duan, L., Yu, Q., Zhang, Q., Wang, Z., Pan, Y., Larssen, T., Tang, J. and Mulder, J. (2016). Acid deposition in Asia: Emissions, deposition, and ecosystem effects. *Atmos. Environ.* 146: 55–69.
- Ebrahimi, M. and Taherianfard, M. (2011). Pathological and hormonal changes in freshwater fishes due to exposure to heavy metals pollutants. *Water Air Soil Pollut.* 217: 47–55.
- Edmondson, J.L., Carroll, J.A., Price, E.A. and Caporn, S.J. (2010). Bio-indicators of nitrogen pollution in heather moorland. *Sci. Total Environ.* 408: 6202–6209.
- Efroymson, R., Will, M., Suter, G. and Wooten, A. (1997). *Toxicological benchmarks for screening contaminants of potential concern for effects on terrestrial plants: 1997 revision*. US Department of Energy, Washington.
- Eisler, R. (1985). *Cadmium hazards to fish, wildlife, and invertebrates: A synoptic review*. Fish and Wildlife Service, US Department of the Interior.
- Eisler, R. (1998). *Nickel hazards to fish, wildlife, and invertebrates: A synoptic review*, Contaminant Hazard Reviews Report 34, Biological Science Report USGS/BRD/BSR--1998-0001, U.S. Geological Survey, Washington DC.
- Elliott-Kingston, C., Haworth, M. and McElwain, J. (2014).

- Damage structures in leaf epidermis and cuticle as an indicator of elevated atmospheric sulphur dioxide in early Mesozoic floras. *Rev. Palaeobot. Palynol.* 208: 25–42.
- Emberson, L. and Buker, P. (2011). *Current knowledge of the impacts of ozone on food crops in South Asia. Ozone Pollution: A hidden threat to food security.* ICP Vegetation Report, Center for Ecology, Bangor, UK, pp. 83–92.
- Eng, A., Harner, T. and Pozo, K. (2014). A Prototype passive air sampler for measuring dry deposition of polycyclic aromatic hydrocarbons. *Environ. Sci. Technol. Lett.* 1: 77–81.
- Ericson, G., Lindesjö, E. and Balk, L. (1998). DNA adducts and histopathological lesions in perch (*Perca fluviatilis*) and northern pike (*Esox lucius*) along a polycyclic aromatic hydrocarbon gradient on the Swedish coastline of the Baltic Sea. *Can. J. Fish. Aquat. Sci.* 55: 815–824.
- Evans, R.E., Brown, S.B. and Hara, T.J. (1988). The effects of aluminum and acid on the gill morphology in rainbow trout, *Salmo gairdneri*. *Environ. Biol. Fishes* 22: 299–311.
- Evers, D., Taylor, K., Major, A., Taylor, R., Poppenga, R. and Scheuhammer, A. (2003). Common loon eggs as indicators of methylmercury availability in North America. *Ecotoxicology* 12: 69–81.
- Evers, D.C., Burgess, N.M., Champoux, L., Hoskins, B., Major, A., Goodale, W.M., Taylor, R.J., Poppenga, R. and Daigle, T. (2005). Patterns and interpretation of mercury exposure in freshwater avian communities in northeastern north America. *Ecotoxicology* 14: 193–221.
- Evers, D.C., Savoy, L.J., DeSorbo, C.R., Yates, D.E., Hanson, W., Taylor, K.M., Siegel, L.S., Cooley, J.H., Bank, M.S. and Major, A. (2008). Adverse effects from environmental mercury loads on breeding common loons. *Ecotoxicology* 17: 69–81.
- Ewais, E. (1997). Effects of cadmium, nickel and lead on growth, chlorophyll content and proteins of weeds. *Biol. Plant.* 39: 403–410.
- Exley, C., Chappell, J. and Birchall, J. (1991). A mechanism for acute aluminium toxicity in fish. *J. Theor. Biol.* 151: 417–428.
- Fairchild, I.J., Loader, N.J., Wynn, P.M., Frisia, S., Thomas, P.A., Lagaard, J.G., Momi, A.d., Hartland, A., Borsato, A. and Porta, N.L. (2009). Sulfur fixation in wood mapped by synchrotron X-ray studies: Implications for environmental archives. *Environ. Sci. Technol.* 43: 1310–1315.
- Fakhraei, H., Driscoll, C.T., Renfro, J.R., Kulp, M.A., Blett, T.F., Brewer, P.F. and Schwartz, J.S. (2016). Critical loads and exceedances for nitrogen and sulfur atmospheric deposition in Great Smoky Mountains National Park, United States. *Ecosphere* 7:e 01466.
- Falguera, V., Sánchez-Riaño, A.M., Quintero-Cerón, J.P., Rivera-Barrero, C.A., Méndez-Arteaga, J.J. and Ibarz, A. (2012). Characterization of polyphenol oxidase activity in juices from 12 underutilized tropical fruits with high agroindustrial potential. *Food Bioprocess Technol.* 5: 2921–2927.
- Falkengren-Grerup, U., Linnermark, N. and Tyler, G. (1987). Changes in acidity and cation pools of south Swedish soils between 1949 and 1985. *Chemosphere* 16: 2239–2248.
- Feest, A., van Swaay, C. and van Hinsberg, A. (2014). Nitrogen deposition and the reduction of butterfly biodiversity quality in the Netherlands. *Ecol. Indic.* 39: 115–119.
- Feng, Z.W., Jin, M.H., Zhang, F.Z. and Huang, Y.Z. (2003). Effects of ground-level ozone (O<sub>3</sub>) pollution on the yields of rice and winter wheat in the Yangtze River Delta. *J. Environ. Sci. (China)* 15: 360–362.
- Fenn, M.E., Lambert, K.F., Blett, T.F., Burns, D.A., Pardo, L.H., Lovett, G.M., Haeuber, R.A., Evers, D.C., Driscoll, C.T. and Jeffries, D.S. (2011). Setting limits: Using air pollution thresholds to protect and restore U.S. ecosystems. *Issues in Ecology* No. 14. Ecological Society of America, Washington, DC.
- Fernandes Azevedo, B., Barros Furieri, L., Peçanha, F.M., Wiggers, G.A., Frizera Vassallo, P., Ronacher Simões, M., Fiorim, J., Rossi de Batista, P., Fioresi, M. and Rossoni, L. (2012). Toxic effects of mercury on the cardiovascular and central nervous systems. *J. Biomed. Biotechnol.* 2012: 949048.
- Fernández, J., Boquete, M., Carballeira, A. and Aboal, J. (2015). A critical review of protocols for moss biomonitoring of atmospheric deposition: Sampling and sample preparation. *Sci. Total Environ.* 517: 132–150.
- Fernández-Varela, R., Ratola, N., Alves, A. and Amigo, J.M. (2015). Relationship between levels of polycyclic aromatic hydrocarbons in pine needles and socio-geographic parameters. *J. Environ. Manage.* 156: 52–61.
- Fernie, K.J., Martinson, S.C., Chen, D., Eng, A., Harner, T., Smits, J.E.G., Soos, C. (2018). Elevated exposure, uptake and accumulation of polycyclic aromatic hydrocarbons by nestling tree swallows (*Tachycineta bicolor*) through multiple exposure routes in active mining-related areas of the Athabasca oil sands region. *Sci. Total Environ.* 624: 250–261.
- Ferreira, V. and Guérol, F. (2017). Leaf litter decomposition as a bioassessment tool of acidification effects in streams: Evidence from a field study and meta-analysis. *Ecol. Indic.* 79: 382–390.
- Field, C.D., Dise, N.B., Payne, R.J., Britton, A.J., Emmett, B.A., Helliwell, R.C., Hughes, S., Jones, L., Lees, S. and Leake, J.R. (2014). The role of nitrogen deposition in widespread plant community change across semi-natural habitats. *Ecosystems* 17: 864–877.
- Figueira, R., Sérgio, C. and Sousa, A. (2002). Distribution of trace metals in moss biomonitors and assessment of contamination sources in Portugal. *Environ. Pollut.* 118: 153–163.
- Filella, I., Peñuelas, J. and Ribas, A. (2005). Using plant biomonitors and flux modelling to develop O<sub>3</sub> dose-response relationships in Catalonia. *Environ. Pollut.* 134: 145–151.
- Fölster, J., Andrén, C., Bishop, K., Buffam, I., Cory, N.,

- Goedkoop, W., Holmgren, K., Johnson, R., Laudon, H. and Wilander, A. (2007). A novel environmental quality criterion for acidification in Swedish lakes—An application of studies on the relationship between biota and water chemistry. *Water Air Soil Pollut. Focus* 7: 331–338.
- Fowler, D., Coyle, M., Skiba, U., Sutton, M.A., Cape, J.N., Reis, S., Sheppard, L.J., Jenkins, A., Grizzetti, B. and Galloway, J.N. (2013). The global nitrogen cycle in the twenty-first century. *Phil. Trans. R. Soc. B* 368: 20130164.
- French, T.D., Houben, A.J., Desforges, J.P., Kimpe, L.E., Kokelj, S.V., Poulain, A.J., Smol, J.P., Wang, X. and Blais, J.M. (2014). Dissolved organic carbon thresholds affect mercury bioaccumulation in Arctic lakes. *Environ. Sci. Technol.* 48: 3162–3168.
- Fuhrer, J., Skarby, L. and Ashmore, M.R. (1997). Critical levels for ozone effects on vegetation in Europe. *Environ. Pollut.* 97: 91–106.
- Fuhrer, J., Val Martin, M., Mills, G., Heald, C.L., Harmens, H., Hayes, F., Sharps, K., Bender, J. and Ashmore, M.R. (2016). Current and future ozone risks to global terrestrial biodiversity and ecosystem processes. *Ecol. Evol.* 6: 8785–8799.
- Fusaro, L., Palma, A., Salvatori, E., Basile, A., Maresca, V., Karam, E.A. and Manes, F. (2017). Functional indicators of response mechanisms to nitrogen deposition, ozone, and their interaction in two Mediterranean tree species. *PLoS one* 12: e0185836.
- Futter, M.N., Valinia, S., Löfgren, S., Köhler, S.J. and Fölster, J. (2014). Long-term trends in water chemistry of acid-sensitive Swedish lakes show slow recovery from historic acidification. *Ambio* 43: 77–90.
- Gallego, S.M., Pena, L.B., Barcia, R.A., Azpilicueta, C.E., Iannone, M.F., Rosales, E.P., Zawoznik, M.S., Groppa, M.D. and Benavides, M.P. (2012). Unravelling cadmium toxicity and tolerance in plants: Insight into regulatory mechanisms. *Environ. Exp. Bot.* 83: 33–46.
- Gandhi, N., Bhavsar, S.P., Tang, R.W. and Arhonditsis, G.B. (2015). Projecting fish mercury levels in the province of Ontario, Canada and the implications for fish and human health. *Environ. Sci. Technol.* 49: 14494–14502.
- Garty, J. (2000). Environment and Elemental content of lichens. *Trace Met. Environ.* 4: 245–276.
- Georgieva, E., Velcheva, I., Yancheva, V. and Stoyanova, S. (2014). Trace metal effects on gill epithelium of common carp, *Cyprinus carpio* L. (Cyprinidae). *Acta Zool. Bulg.* 66: 277–282.
- Gilliam, F.S. (2006). Response of the herbaceous layer of forest ecosystems to excess nitrogen deposition. *J. Ecol.* 94: 1176–1191.
- Gilmour, C.C., Henry, E.A. and Mitchell, R. (1992). Sulfate stimulation of mercury methylation in freshwater sediments. *Environ. Sci. Technol.* 26: 2281–2287.
- Glooschenko, W.A. (1989). *Sphagnum fuscum* moss as an indicator of atmospheric cadmium deposition across Canada. *Environ. Pollut.* 57: 27–33.
- Godek, M., Sobik, M., Błaś, M., Polkowska, Ż., Owczarek, P. and Bokwa, A. (2015). Tree rings as an indicator of atmospheric pollutant deposition to subalpine spruce forests in the Sudetes (Southern Poland). *Atmos. Res.* 151: 259–268.
- Goodale, M.W., Evers, D.C., Mierzykowski, S.E., Bond, A.L., Burgess, N.M., Otorowski, C.I., Welch, L.J., Hall, C.S., Ellis, J.C. and Allen, R.B. (2008). Marine foraging birds as bioindicators of mercury in the Gulf of Maine. *EcoHealth* 5: 409–425.
- Goodman, J.E., Prueitt, R.L., Sax, S.N., Pizzurro, D.M., Lynch, H.N., Zu, K. and Venditti, F.J. (2015). Ozone exposure and systemic biomarkers: Evaluation of evidence for adverse cardiovascular health impacts. *Crit. Rev. Toxicol.* 45: 412–452.
- Gordon, M., Knauer, G.A. and Martin, J.H. (1980). *Mytilus californianus* as a bioindicator of trace metal pollution: Variability and statistical considerations. *Mar. Pollut. Bull.* 11: 195–198.
- Govers, L.L., Lamers, L.P., Bouma, T.J., Eygensteyn, J., de Brouwer, J.H., Hendriks, A.J., Huijbers, C.M. and van Katwijk, M.M. (2014). Seagrasses as indicators for coastal trace metal pollution: A global meta-analysis serving as a benchmark, and a Caribbean case study. *Environ. Pollut.* 195: 210–217.
- Gray, S.M., Chapman, L.J. and Mandrak, N.E. (2012). Turbidity reduces hatching success in Threatened Spotted Gar (*Lepisosteus oculatus*) *Environ. Biol. Fishes* 94: 689–694.
- Grimmer, G., Brune, H., Deutsch-Wenzel, R., Naujack, K.W., Misfeld, J. and Timm, J. (1983). On the contribution of polycyclic aromatic hydrocarbons to the carcinogenic impact of automobile exhaust condensate evaluated by local application onto mouse skin. *Cancer Lett.* 21: 105–113.
- Grodzińska, K. (1977). Acidity of tree bark as a bioindicator of forest pollution in Southern Poland. *Water Air Soil Pollut.* 8: 3–7.
- Gundersen, P. and Rasmussen, L. (1990). Nitrification in forest soils: Effects from nitrogen deposition on soil acidification and aluminum release. In *Reviews of environmental contamination and toxicology*, Ware, G.W. (Ed.), Springer New York, pp. 1–45.
- Hahon, N. and Booth, J.A. (1986). Coinhibition of viral interferon induction by benzo[a]pyrene and chrysotile asbestos. *Environ. Res.* 40: 103–109.
- Hallingbäck, T. and Kellner, O. (1992). Effects of simulated nitrogen rich and acid rain on the nitrogen-fixing lichen *Peltigera aphthosa* (L.) Willd. *New Phytol.* 120: 99–103.
- Hames, R.S., Rosenberg, K.V., Lowe, J.D., Barker, S.E. and Dhondt, A.A. (2002). Adverse effects of acid rain on the distribution of the Wood Thrush *Hylocichla ustulata* mustelina in North America. *Proc. Natl. Acad. Sci. U.S.A.* 99: 11235–11240.
- Han, W., Fang, J., Guo, D. and Zhang, Y. (2005). Leaf nitrogen and phosphorus stoichiometry across 753 terrestrial plant species in China. *New Phytol.* 168: 377–385.
- Hanson, N. and Larsson, Å. (2008). Fixed wavelength fluorescence to detect PAH metabolites in fish bile: Increased statistical power with an alternative dilution

- method. *Environ. Monit. Assess.* 144: 221–228.
- Harada, M. (1995). Minamata disease: Methylmercury poisoning in Japan caused by environmental pollution. *Crit. Rev. Toxicol.* 25: 1–24.
- Harmens, H., Norris, D., Sharps, K., Mills, G., Alber, R., Aleksiyenak, Y., Blum, O., Cucu-Man, S.M., Dam, M. and De Temmerman, L. (2015). Heavy metal and nitrogen concentrations in mosses are declining across Europe whilst some “hotspots” remain in 2010. *Environ. Pollut.* 200: 93–104.
- Harner, T., Su, K., Genualdi, S., Karpowicz, J., Ahrens, L., Mihele, C., Schuster, J., Charland, J.P. and Narayan, J. (2013). Calibration and application of PUF disk passive air samplers for tracking polycyclic aromatic compounds (PACs). *Atmos. Environ.* 75: 123–128.
- Harrison, R.M., Grenfell, J., Yamulki, S., Clemitshaw, K., Penkett, S., Cape, J. and McFadyen, G. (1999). Budget of NO<sub>y</sub> species measured at a coastal site. *Atmos. Environ.* 33: 4255–4272.
- Hassan, F.M., Al-Tae, M.M. and Mohammed, A.B. (2010). *Ceratophyllum demersum* L. and *Typha domingensis* Pers as bioindicator of some PAHs compounds in Euphrates river at AL-Hindiya city. *Basrah J. Sci.* 28: 288–298.
- Haworth, M., Elliott-Kingston, C., Gallagher, A., Fitzgerald, A. and McElwain, J.C. (2012). Sulphur dioxide fumigation effects on stomatal density and index of non-resistant plants: Implications for the stomatal palaeo-[CO<sub>2</sub>] proxy method. *Rev. Palaeobot. Palynol.* 182: 44–54.
- Haworth, M., Gallagher, A., Elliott-Kingston, C., Raschi, A., Marandola, D. and McElwain, J.C. (2010). Stomatal index responses of *Agrostis canina* to CO<sub>2</sub> and sulphur dioxide: Implications for palaeo-[CO<sub>2</sub>] using the stomatal proxy. *New Phytol.* 188: 845–855.
- Hayden, K., Anlauf, K., Hastie, D. and Bottenheim, J. (2003). Partitioning of reactive atmospheric nitrogen oxides at an elevated site in southern Quebec, Canada. *J. Geophys. Res.* 108: 4603.
- Hebert, C.E., Weseloh, D., MacMillan, S., Campbell, D. and Nordstrom, W. (2011). Metals and polycyclic aromatic hydrocarbons in colonial waterbird eggs from Lake Athabasca and the Peace-Athabasca Delta, Canada. *Environ. Toxicol. Chem.* 30: 1178–1183.
- Hebert, C.E., Campbell, D., Kindopp, R., MacMillan, S., Martin, P., Neugebauer, E., Patterson, L. and Shatford, J. (2013). Mercury trends in colonial waterbird eggs downstream of the oil sands region of Alberta, Canada. *Environ. Sci. Technol.* 47: 11785–11792.
- Hellou, J., Upshall, C., Taylor, D., O'Keefe, P., O'malley, V. and Abrajano, T. (1994). Unsaturated hydrocarbons in muscle and hepatopancreas of two crab species, *Chionoecetes opilio* and *Hyas coarctatus*. *Mar. Pollut. Bull.* 28: 482–488.
- Henry, J. and Aherne, J. (2014). Nitrogen deposition and exceedance of critical loads for nutrient nitrogen in Irish grasslands. *Sci. Total Environ.* 470–471: 216–223.
- Hicks, W., Leith, I., Woodin, S. and Fowler, D. (2000). Can the foliar nitrogen concentration of upland vegetation be used for predicting atmospheric nitrogen deposition? Evidence from field surveys. *Environ. Pollut.* 107: 367–376.
- Hirano, T. and Tamae, K. (2011). Earthworms and soil pollutants. *Sensors* 11: 11157–11167.
- Hobbs, W.O., Telford, R.J., Birks, H.J.B., Saros, J.E., Hazewinkel, R.R., Perren, B.B., Saulnier-Talbot, É. and Wolfe, A.P. (2010). Quantifying recent ecological changes in remote lakes of North America and Greenland using sediment diatom assemblages. *PLoS one* 5: e10026.
- Hogan, C.M., Proemse, B.C. and Barmuta, L.A. (2017). Isotopic fingerprinting of atmospheric nitrogen and sulfur using lichens (*Cladia retipora*) in Tasmania, Australia. *Appl. Geochem.* 84: 126–132.
- Holoubek, I., Kořínek, P., Šeda, Z., Schneiderová, E., Holoubková, I., Pacl, A., Tříška, J., Cudlín, P. and Čáslavský, J. (2000). The use of mosses and pine needles to detect persistent organic pollutants at local and regional scales. *Environ. Pollut.* 109: 283–292.
- Hong, Y.S., Kim, Y.M. and Lee, K.E. (2012). Methylmercury exposure and health effects. *J. Prev. Med. Public Health* 45: 353.
- Horii, C.V., Munger, J.W., Wofsy, S.C., Zahniser, M., Nelson, D. and McManus, J.B. (2005). Atmospheric reactive nitrogen concentration and flux budgets at a Northeastern U.S. forest site. *Agric. For. Meteorol.* 133: 210–225.
- Hůnová, I., Maznová, J. and Kurfürst, P. (2014). Trends in atmospheric deposition fluxes of sulphur and nitrogen in Czech forests. *Environ. Pollut.* 184: 668–675.
- Hwang, H.M. and Wade, T.L. (2008). Aerial distribution, temperature-dependent seasonal variation, and sources of polycyclic aromatic hydrocarbons in pine needles from the Houston metropolitan area, Texas, USA. *J. Environ. Sci. Health., Part A* 43: 1243–1251.
- ICP Vegetation (2017). The international cooperative programme on effects of air pollution on natural vegetation and crops, <https://icpvegetation.ceh.ac.uk/index.html>, Last Access: 30 January 2018.
- Irvine, I.C., Greaver, T., Phelan, J., Sabo, R.D. and Houtven, G.V. (2017). Terrestrial acidification and ecosystem services: Effects of acid rain bunnies, baseball, and Christmas trees. *Ecosphere* 8: e01857.
- Isbell, F., Tilman, D., Polasky, S., Binder, S. and Hawthorne, P. (2013). Low biodiversity state persists two decades after cessation of nutrient enrichment. *Ecol. Lett.* 16: 454–460.
- Iwu, N.W. (2010). Mercury accumulation in the northern pike (*Esox lucius*) and its intestinal parasite, *Triaenophorus nodulosus* (Cestoda). Master's thesis, Department of Biological and Environmental Science, University of Jyväskylä, Finland.
- Jaffar, M., Ashraf, M. and Tariq, J. (1995). Assessment of current trace metal pollution. status of the South East Arabian Sea Coast of Pakistan through fish. *J. Chem. Soc. Pakistan* 17: 204–207.
- Jariyasopit, N., Harner, T., Wu, D., Williams, A., Halappanavar, S. and Su, K. (2016). Mapping indicators of toxicity for polycyclic aromatic compounds in the

- atmosphere of the athabasca oil sands region. *Environ. Sci. Technol.* 50: 11282–11291.
- Jariyasopit, N., Zhang, Y., Martin, J. and Harner, T. (2018). Comparison of polycyclic aromatic compounds in air measured by conventional passive air samplers and passive dry deposition samplers and contributions from petcoke and oil sands ore. *Atmos. Chem. Phys.* 18: 9161–9171.
- Jasion, M., Samecka-Cymerman, A., Kolon, K. and Kempers, A.J. (2013). *Tanacetum vulgare* as a bioindicator of trace-metal contamination: A study of a naturally colonized open-pit lignite mine. *Arch. Environ. Contam. Toxicol.* 65: 442–448.
- Jeffries, D. and Lam, D. (1993). Assessment of the effect of acidic deposition on Canadian lakes: Determination of critical loads for sulphate deposition. *Water Sci. Technol.* 28: 183–187.
- Joner, E.J., Leyval, C. and Colpaert, J.V. (2006). Ectomycorrhizas impede phytoremediation of polycyclic aromatic hydrocarbons (PAHs) both within and beyond the rhizosphere. *Environ. Pollut.* 142: 34–38.
- Jones, L., Provins, A., Holland, M., Mills, G., Hayes, F., Emmett, B., Hall, J., Sheppard, L., Smith, R. and Sutton, M. (2014). A review and application of the evidence for nitrogen impacts on ecosystem services. *Ecosyst. Serv.* 7: 76–88.
- Jones, L., Stevens, C., Rowe, E., Payne, R., Caporn, S.J., Evans, C.D., Field, C. and Dale, S. (2016). Can on-site management mitigate nitrogen deposition impacts in non-wooded habitats? *Biol. Conserv.* 212: 464–475.
- Jones, M.L., Wallace, H.L., Norris, D., Brittain, S.A., Haria, S., Jones, R.E., Rhind, P.M., Reynolds, B.R. and Emmett, B.A. (2004). Changes in vegetation and soil characteristics in coastal sand dunes along a gradient of atmospheric nitrogen deposition. *Plant Biol. (Stuttg)* 6: 598–605.
- Jones, S.K., Helfter, C., Anderson, M., Coyle, M., Campbell, C., Famulari, D., Di Marco, C., van Dijk, N., Tang, Y.S. and Topp, C.F. (2017). The nitrogen, carbon and greenhouse gas budget of a grazed, cut and fertilised temperate grassland. *Biogeosciences* 14: 2069.
- Jönsson, U., Rosengren, U., Thelin, G. and Nihlgård, B. (2003). Acidification-induced chemical changes in coniferous forest soils in southern Sweden 1988–1999. *Environ. Pollut.* 123: 75–83.
- Jorge, M.G.C., Rita, A.L.M., Alexander, I.A. and Marianela, R.D.M. (2011). Al<sup>3+</sup>-Ca<sup>2+</sup> interaction in plants growing in acid soils: Al-phytotoxicity response to calcareous amendments. *J. Soil Sci. Plant Nutr.* 10: 217–243.
- Kabata-Pendias, A. and Dudka, S. (1991). Trace metal contents of *Taraxacum officinale* (dandelion) as a convenient environmental indicator. *Environ. Geochem. Health* 13: 108–113.
- Kafiatullah, Wahid, A., Ahmad, S.S. and Shamsi, S.R.A. (2012). Ozone biomonitoring in Pakistan using tobacco cultivar Bel-W3. *Pak. J. Bot.* 44:717–723.
- Karlsson, P.E., Uddling, J., Braun, S., Broadmeadow, M., Elvira, S., Gimeno, B.S., Le Thiec, D., Oksanen, E., Vandermeiren, K., Wilkinson, M. and Emberson, L. (2003). *New critical levels for ozone impact on trees based on AOT40 and leaf cumulated uptake of ozone, in establishing ozone critical levels II*, Karlsson, P.E., Selldén, G. and Pleijel, H. (Eds.), UNECE Workshop Report. IVL Report B 1523. IVL Swedish Environmental Research Institute, Gothenburg, Sweden.
- Kato, K., Yoshinori, K., Wataru, O. and Koki, K. (2007). Nitrogen fixation in seabuckthorn (*Hippophae rhamnoides* L.) root nodules and effect of nitrate on nitrogenase activity. *J. Jpn. Soc. Hortic. Sci.* 76: 185–190.
- Keelan, R.E. (2007). *The ecological effects of ozone exposure on upland semi-natural vegetation of the Yorkshire Dales National Park*, University of York.
- Kelly, E.N., Short, J.W., Schindler, D.W., Hodson, P.V., Ma, M., Kwan, A.K. and Fortin, B.L. (2009). Oil sands development contributes polycyclic aromatic compounds to the Athabasca River and its tributaries. *Proc. Natl. Acad. Sci. U.S.A.* 106: 22346–22351.
- Kenow, K.P., Gutreuter, S., Hines, R.K., Meyer, M.W., Fournier, F. and Karasov, W.H. (2003). Effects of methyl mercury exposure on the growth of juvenile common loons. *Ecotoxicology* 12: 171–181.
- Keuter, A., Veldkamp, E. and Corre, M.D. (2014). Asymbiotic biological nitrogen fixation in a temperate grassland as affected by management practices. *Soil Biol. Biochem.* 70: 38–46.
- Kim, K.H., Jahan, S.A., Kabir, E. and Brown, R.J. (2013). A review of airborne polycyclic aromatic hydrocarbons (PAHs) and their human health effects. *Environ. Int.* 60: 71–80.
- King, J.K., Kostka, J.E., Frischer, M.E. and Saunders, F.M. (2000). Sulfate-reducing bacteria methylate mercury at variable rates in pure culture and in marine sediments. *Appl. Environ. Microbiol.* 66: 2430–2437.
- Kinghorn, A., Solomon, P. and Chan, H.M. (2007). Temporal and spatial trends of mercury in fish collected in the English–Wabigoon river system in Ontario, Canada. *Sci. Total Environ.* 372: 615–623.
- Kirkham, F., Mountford, J. and Wilkins, R. (1996). The effects of nitrogen, potassium and phosphorus addition on the vegetation of a Somerset peat moor under cutting management. *J. Appl. Ecol.* 33: 1013–1029.
- Kleijn, D., Bekker, R.M., Bobbink, R., De Graaf, M.C. and Roelofs, J.G. (2008). In search for key biogeochemical factors affecting plant species persistence in heathland and acidic grasslands: A comparison of common and rare species. *J. Appl. Ecol.* 45: 680–687.
- Klumpp, A., Ansel, W., Klumpp, G., Vergne, P., Sifakis, N., Sanz, M.J., Rasmussen, S., Ro-Poulsen, H., Ribas, A., Peñuelas, J., Kambezidis, H., He, S., Garrec, J.P. and Calatayud, V. (2006). Ozone pollution and ozone biomonitoring in European cities Part II. Ozone-induced plant injury and its relationship with descriptors of ozone pollution. *Atmos. Environ.* 40: 7437–7448.
- Koivula, T.T., Salkinoja-Salonen, M., Peltola, R. and Romantschuk, M. (2004). Pyrene degradation in forest humus microcosms with or without pine and its mycorrhizal fungus. *J. Environ. Qual.* 33: 45–53.

- Kooijman, A., van Til, M., Noordijk, E., Remke, E. and Kalbitz, K. (2016). Nitrogen deposition and grass encroachment in calcareous and acidic Grey dunes (H2130) in NW-Europe. *Biol. Conserv.* 212: 406–415.
- Koranda, M., Kerschbaum, S., Wanek, W., Zechmeister, H. and Richter, A. (2007). Physiological responses of bryophytes *Thuidium tamariscinum* and *Hylocomium splendens* to increased nitrogen deposition. *Ann. Bot.* 99: 161–169.
- Koseva, I.S., Watmough, S.A. and Aherne, J. (2010). Estimating base cation weathering rates in Canadian forest soils using a simple texture-based model. *Biogeochemistry* 101: 183–196.
- Krewski, D., Yokel, R.A., Nieboer, E., Borchelt, D., Cohen, J., Harry, J., Kacew, S., Lindsay, J., Mahfouz, A.M. and Rondeau, V. (2007). Human health risk assessment for aluminium, aluminium oxide, and aluminium hydroxide. *J. Toxicol. Environ. Health B* 10: 1–269.
- Kroglund, F., Rosseland, B., Teien, H.C., Salbu, B., Kristensen, T. and Finstad, B. (2008). Water quality limits for Atlantic salmon (*Salmo salar* L.) exposed to short term reductions in pH and increased aluminum simulating episodes. *Hydrol. Earth Syst. Sci.* 12: 491–507.
- Królak, E. (2003). Accumulation of Zn, Cu, Pb and Cd by dandelion (*Taraxacum officinale* Web.) in environments with various degrees of metallic contamination. *Pol. J. Environ. Stud.* 12: 713–721.
- Krupa, S. (2003). Effects of atmospheric ammonia (NH<sub>3</sub>) on terrestrial vegetation: A review. *Environ. Pollut.* 124: 179–221.
- Kummerová, M. and Kmentová, E. (2004). Photoinduced toxicity of fluoranthene on germination and early development of plant seedling. *Chemosphere* 56: 387–393.
- Kurek, J., Kirk, J.L., Muir, D.C., Wang, X., Evans, M.S. and Smol, J.P. (2013). Legacy of a half century of Athabasca oil sands development recorded by lake ecosystems. *Proc. Natl. Acad. Sci. U.S.A.* 110: 1761–1766.
- Larsen, T. and Carmichael, G. (2000). Acid rain and acidification in China: the importance of base cation deposition. *Environ. Pollut.* 110: 89–102.
- Lee, K.S., Hung, D.V., Kwak, J.H., Lim, S.S., Lee, K.H. and Choi, W.J. (2011). Tree ring Ca/Al as an indicator of historical soil acidification of *Pinus densiflora* forest in southern Korea. *Korean J. Environ. Agric.* 30: 229–233.
- Lehndorff, E. and Schwark, L. (2009). Biomonitoring airborne parent and alkylated three-ring PAHs in the Greater Cologne Conurbation I: Temporal accumulation patterns. *Environ. Pollut.* 157: 1323–1331.
- Lepori, F. and Keck, F. (2012). Effects of atmospheric nitrogen deposition on remote freshwater ecosystems. *Ambio* 41: 235–246.
- Leyval, C., Turnau, K. and Haselwandter, K. (1997). Effect of heavy metal pollution on mycorrhizal colonization and function: Physiological, ecological and applied aspects. *Mycorrhiza* 7: 139–153.
- Li, G., Hu, B., Bi, J., Leng, Q., Xiao, C. and Yang, Z. (2013). Heavy metals distribution and contamination in surface sediments of the coastal Shandong Peninsula (Yellow Sea). *Mar. Pollut. Bull.* 76: 420–426.
- Librando, V., Perrini, G. and Tomasello, M. (2002). Biomonitoring of atmospheric PAHs by evergreen plants: correlations and applicability. *Polycyclic Aromat. Compd.* 22: 549–559.
- Lien, L., Raddum, G.G., Fjellheim, A. and Henriksen, A. (1996). A critical limit for acid neutralizing capacity in Norwegian surface waters, based on new analyses of fish and invertebrate responses. *Sci. Total Environ.* 177: 173–193.
- Lin, E.L., Cormier, S.M. and Torsella, J.A. (1996). Fish biliary polycyclic aromatic hydrocarbon metabolites estimated by fixed-wavelength fluorescence: comparison with HPLC-fluorescent detection. *Ecotoxicol. Environ. Saf.* 35: 16–23.
- Lin, L.F., Shih, S.I., Su, J.W., Shih, M., Lin, K.C., Wang, L.C. and Chang-Chien, G.P. (2010). Dry and wet deposition of polychlorinated dibenzo-*p*-dioxins and dibenzofurans on the drinking water treatment plant. *Aerosol Air Qual. Res.* 10: 231–234.
- Lindroth, R.L. (2010). Impacts of elevated atmospheric CO<sub>2</sub> and O<sub>3</sub> on forests: phytochemistry, trophic interactions, and ecosystem dynamics. *J. Chem. Ecol.* 36: 2–21.
- Liu, H., Weisman, D., Ye, Y.B., Cui, B., Huang, Y.H., Colón-Carmona, A. and Wang, Z.H. (2009). An oxidative stress response to polycyclic aromatic hydrocarbon exposure is rapid and complex in *Arabidopsis thaliana*. *Plant Sci.* 176: 375–382.
- Liu, X., Zhang, Y., Han, W., Tang, A., Shen, J., Cui, Z., Vitousek, P., Erisman, J.W., Goulding, K. and Christie, P. (2013). Enhanced nitrogen deposition over China. *Nature* 494: 459–462.
- Lodovici, M., Akpan, V., Casalini, C., Zappa, C. and Dolara, P. (1998). Polycyclic aromatic hydrocarbons in *Laurus nobilis* leaves as a measure of air pollution in urban and rural sites of Tuscany. *Chemosphere* 36: 1703–1712.
- Lourino-Cabana, B., Lesven, L., Charriau, A., Billon, G., Ouddane, B. and Boughriet, A. (2011). Potential risks of metal toxicity in contaminated sediments of Deûle river in northern France. *J. Hazard. Mater.* 186: 2129–2137.
- Lu, M., Yang, Y., Luo, Y., Fang, C., Zhou, X., Chen, J., Yang, X. and Li, B. (2011). Responses of ecosystem nitrogen cycle to nitrogen addition: A meta-analysis. *New Phytol.* 189: 1040–1050.
- Lu, X., Mo, J., Gilliam, F.S., Zhou, G. and Fang, Y. (2010). Effects of experimental nitrogen additions on plant diversity in an old-growth tropical forest. *Global Change Biol.* 16: 2688–2700.
- Lu, X., Mao, Q., Gilliam, F.S., Luo, Y. and Mo, J. (2014). Nitrogen deposition contributes to soil acidification in tropical ecosystems. *Global Change Biol.* 20: 3790–3801.
- Lucas, R., Klaminder, J., Fitter, M., Bishop, K.H., Egnell, G., Laudon, H. and Högberg, P. (2011). A meta-analysis of the effects of nitrogen additions on base cations:

- Implications for plants, soils, and streams. *For. Ecol. Manage.* 262: 95–104.
- Lundin, J.I., Riffell, J.A. and Wasser, S.K. (2015). Polycyclic aromatic hydrocarbons in caribou, moose, and wolf scat samples from three areas of the Alberta oil sands. *Environ. Pollut.* 206: 527–534.
- Lydersen, E., Larssen, T. and Fjeld, E. (2004). The influence of total organic carbon (TOC) on the relationship between acid neutralizing capacity (ANC) and fish status in Norwegian lakes. *Sci. Total Environ.* 326: 63–69.
- Macduff, J., Jarvis, S. and Davidson, I. (1996). Inhibition of N<sub>2</sub> fixation by white clover (*Trifolium repens* L.) at low concentrations of NO<sub>inf3</sub><sup>sup-</sup> in flowing solution culture. *Plant Soil* 180: 287–295.
- Maes, D., Decler, K., De Keersmaeker, L., Van Uytvanck, J. and Louette, G. (2017). Intensified habitat management to mitigate negative effects of nitrogen pollution can be detrimental for faunal diversity: A comment on Jones et al. (2017). *Biol. Conserv.* 212: 493–494.
- Malmgren, R., Bennison, B. and McKinley Jr, T. (1952). Reduced antibody titers in mice treated with carcinogenic and cancer chemotherapeutic agents. *Proc. Soc. Exp. Biol. Med.* 79: 484–488.
- Mance, G. (1987). Introduction, In *pollution threat of heavy metals in aquatic environments*, Springer, pp. 1–8.
- Manzano, C.A., Marvin, C., Muir, D., Harner, T., Martin, J. and Zhang, Y. (2017). Heterocyclic aromatics in petroleum coke, snow, lake sediments and air samples from the Athabasca oil sands region. *Environ. Sci. Technol.* 51: 5445–5453.
- Marschner, H., Kirkby, E. and Cakmak, I. (1996). Effect of mineral nutritional status on shoot-root partitioning of photoassimilates and cycling of mineral nutrients. *J. Exp. Bot.* 47: 1255–1263.
- Mashroofeh, A., Bakhtiari, A.R., Ghobeishavi, A., Ahmadpour, M., Asadi, A., Ahmadpour, M., Hosseini, S.H., Eskandari, T. and Burger, J. (2015). Mercury levels in avian feathers from different trophic levels of eight families collected from the northern region of Iran. *Environ. Monit. Assess.* 187: 275.
- Mathot, M., Mertens, J., Verlinden, G. and Lambert, R. (2008). Positive effects of sulphur fertilisation on grasslands yields and quality in Belgium. *Eur. J. Agron.* 28: 655–658.
- Mathot, M., Théliier-Huché, L. and Lambert, R. (2009). Sulphur and nitrogen content as sulphur deficiency indicator for grasses. *Eur. J. Agron.* 30: 172–176.
- Matyssek, R. and Sandermann, H. (2003). Impact of ozone on trees: An ecophysiological perspective, In *Progress in botany: Genetics physiology systematics ecology*, Esser, K., Lüttge, U., Beyschlag, W. and Hellwig, F. (Eds.), Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 349–404.
- McDowell, M.A., Dillon, C.F., Osterloh, J., Bolger, P.M., Pellizzari, E., Fernando, R., de Oca, R.M., Schober, S.E., Sinks, T. and Jones, R.L. (2004). Hair mercury levels in U.S. children and women of childbearing age: Reference range data from NHANES 1999–2000. *Environ. Health Perspect.* 112: 1165.
- McKee, D.J. and Rodriguez, R.M. (1993). Health effects associated with ozone and nitrogen dioxide exposure. *Water Air Soil Pollut.* 67: 11–35.
- McNeil, B.E., Read, J.M. and Driscoll, C.T. (2007). Foliar nitrogen responses to elevated atmospheric nitrogen deposition in nine temperate forest canopy species. *Environ. Sci. Technol.* 41: 5191–5197.
- Meegan, S.K., Perry, S.A. and Perry, W.B. (1996). Detrital processing in streams exposed to acidic precipitation in the Central Appalachian Mountains. *Hydrobiologia* 339: 101–110.
- Meili, M. (1997). Mercury in lakes and rivers. *Met. Ions Biol. Syst.* 34: 21–51.
- Meili, M., Bishop, K., Bringmark, L., Johansson, K., Munthe, J., Sverdrup, H. and de Vries, W. (2003). Critical levels of atmospheric pollution: Criteria and concepts for operational modelling of mercury in forest and lake ecosystems. *Sci. Total Environ.* 304: 83–106.
- Mętrak, M., Aneta, E., Wilkomirski, B., Staszewski, T. and Suska-Malawska, M. (2016). Interspecific differences in foliar 1 PAHs load between Scots pine, birch, and wild rosemary from three polish peat bogs. *Environ. Monit. Assess.* 188: 1–13.
- Migaszewski, Z.M., Gałuszka, A. and Paślawski, P. (2002). Polynuclear aromatic hydrocarbons, phenols, and trace metals in selected soil profiles and plant bioindicators in the Holy Cross Mountains, south-central Poland. *Environ. Int.* 28: 303–313.
- Mills, G., Hayes, F., Simpson, D., Emberson, L., Norris, D., Harmens, H. and Büker, P. (2011). Evidence of widespread effects of ozone on crops and (semi-)natural vegetation in Europe (1990–2006) in relation to AOT40- and flux-based risk maps. *Global Change Biol.* 17: 592–613.
- Mitchell, R., Sutton, M., Truscott, A.M., Leith, I., Cape, J., Pitcairn, C. and Van Dijk, N. (2004). Growth and tissue nitrogen of epiphytic atlantic bryophytes: Effects of increased and decreased atmospheric N deposition. *Funct. Ecol.* 18: 322–329.
- Monson, B.A., Staples, D.F., Bhavsar, S.P., Holsen, T.M., Schrank, C.S., Moses, S.K., McGoldrick, D.J., Backus, S.M. and Williams, K.A. (2011). Spatiotemporal trends of mercury in walleye and largemouth bass from the Laurentian Great Lakes region. *Ecotoxicology* 20: 1555–1567.
- Monteiro, L. and Furness, R. (1995). Seabirds as monitors of mercury in the marine environment. *Water Air Soil Pollut.* 80: 851–870.
- Monteith, D.T., Hildrew, A.G., Flower, R.J., Raven, P.J., Beaumont, W.R., Collen, P., Kreiser, A.M., Shilland, E.M. and Winterbottom, J.H. (2005). Biological responses to the chemical recovery of acidified fresh waters in the UK. *Environ. Pollut.* 137: 83–101.
- Morecroft, M., Sellers, E. and Lee, J. (1994). An experimental investigation into the effects of atmospheric nitrogen deposition on two semi-natural grasslands. *J. Ecol.* 82: 475–483.



- Morgan, J. and Morgan, A. (1988). Earthworms as biological monitors of cadmium, copper, lead and zinc in metalliferous soils. *Environ. Pollut.* 54: 123–138.
- Mountford, J., Lakhani, K. and Kirkham, F. (1993). Experimental assessment of the effects of nitrogen addition under hay-cutting and aftermath grazing on the vegetation of meadows on a Somerset peat moor. *J. Appl. Ecol.* 30: 321–332.
- Mountford, J.O., Lakhani, K. and Holland, R. (1994). *The effects of nitrogen on species diversity and agricultural production on the Somerset Moors, Phase II: A. after seven years of fertiliser application, b. after cessation of fertiliser input for three years*. Final report to the Institute of Grassland and Environmental Research. Huntingdon, Institute of Terrestrial Ecology, Abbots Ripton.
- Musselman, R.C., McCool, P.M. and Lefohn, A.S. (1994). Ozone descriptors for an air quality standard to protect vegetation. *J. Air Waste Manage. Assoc.* 44: 1383–1390.
- Myers, M.S., Johnson, L.L., Olson, O.P., Stehr, C.M., Horness, B.H., Collier, T.K. and McCain, B.B. (1998). Toxicopathic hepatic lesions as biomarkers of chemical contaminant exposure and effects in marine bottomfish species from the Northeast and Pacific Coasts, USA. *Mar. Pollut. Bull.* 37: 92–113.
- Myers, G.J., Davidson, P.W., Cox, C., Shamlaye, C.F., Palumbo, D., Cernichiari, E., Sloane-Reeves, J., Wilding, G.E., Kost, J. and Huang, L.S. (2003). Prenatal methylmercury exposure from ocean fish consumption in the Seychelles child development study. *Lancet* 361: 1686–1692.
- Nakajima, D., Yoshida, Y., Suzuki, J. and Suzuki, S. (1995). Seasonal changes in the concentration of polycyclic aromatic hydrocarbons in azalea leaves and relationship to atmospheric concentration. *Chemosphere* 30: 409–418.
- Nanus, L., Clow, D.W., Saros, J.E., Stephens, V.C. and Campbell, D.H. (2012). Mapping critical loads of nitrogen deposition for aquatic ecosystems in the Rocky Mountains, USA. *Environ. Pollut.* 166: 125–135.
- Nanus, L., McMurray, J., Clow, D., Saros, J., Blett, T. and Gurdak, J. (2017). Spatial variation of atmospheric nitrogen deposition and critical loads for aquatic ecosystems in the Greater Yellowstone Area. *Environ. Pollut.* 223: 644–656.
- Nijssen, M., Alders, K., van der Smissen, N. and Esselink, H. (2001). Effects of grass-encroachment and grazing management on carabid assemblages of dry dune grasslands. Proceedings of the section Experimental and Applied Entomology of the Netherlands Entomological Society, Volume 12, Amsterdam, pp. 113–120.
- Nijssen, M., WallisDeVries, M. and Siepel, H. (2017). Pathways for the effects of increased nitrogen deposition on fauna. *Biol. Conserv.* 212: 423–431.
- Nimis, P.L., Scheidegger, C. and Wolseley, P.A. (2002). *Monitoring with lichens—Monitoring lichens*, Springer Netherlands, pp. 1–4.
- Nkpaa, K., Basse, E. and Wegwu, M. (2013). Evaluation of polycyclic aromatic hydrocarbon (PAH) concentrations in crabs and shrimps from crude oil polluted waters of Ogoniland in rivers state, Nigeria. *IOSR J. Environ. Sci. Toxicol. Food Technol* 4: 73–80.
- Norman, A.L., Belzer, W. and Barrie, L. (2004). Insights into the biogenic contribution to total sulphate in aerosol and precipitation in the Fraser Valley afforded by isotopes of sulphur and oxygen. *J. Geophys. Res.* 109: D05311.
- Ochoa-Hueso, R., Munzi, S., Alonso, R., Arroniz-Crespo, M., Avila, A., Bermejo, V., Bobbink, R., Branquinho, C., Concostrina-Zubiri, L., Cruz, C., Cruz de Carvalho, R., De Marco, A., Dias, T., Elustondo, D., Elvira, S., Estebanez, B., Fusaro, L., Gerosa, G., Izquieta-Rojano, S., Lo Cascio, M., Marzuoli, R., Matos, P., Mereu, S., Merino, J., Morillas, L., Nunes, A., Paoletti, E., Paoli, L., Pinho, P., Rogers, I.B., Santos, A., Sicard, P., Stevens, C.J. and Theobald, M.R. (2017). Ecological impacts of atmospheric pollution and interactions with climate change in terrestrial ecosystems of the Mediterranean Basin: Current research and future directions. *Environ. Pollut.* 227: 194–206.
- Oguntimhin, I., Nakatani, N. and Sakugawa, H. (2008). Phytotoxicities of fluoranthene and phenanthrene deposited on needle surfaces of the evergreen conifer, Japanese red pine (*Pinus densiflora* Sieb. et Zucc.). *Environ. Pollut.* 154: 264–271.
- Oguntimhin, I., Eissa, F. and Sakugawa, H. (2010). Negative effects of fluoranthene on the ecophysiology of tomato plants (*Lycopersicon esculentum* Mill): Fluoranthene mists negatively affected tomato plants. *Chemosphere* 78: 877–884.
- Ohiozebau, E., Tendler, B., Codling, G., Kelly, E., Giesy, J.P. and Jones, P.D. (2017). Potential health risks posed by polycyclic aromatic hydrocarbons in muscle tissues of fishes from the Athabasca and Slave Rivers, Canada. *Environ. Geochem. Health* 39: 139–160.
- Oishi, Y. (2013). Comparison of pine needles and mosses as bio-indicators for polycyclic aromatic hydrocarbons. *J. Environ. Prot.* 4: 106.
- Økland, T., Bakkestuen, V., Økland, R.H. and Eilertsen, O. (2004). Changes in forest understorey vegetation in Norway related to long-term soil acidification and climatic change. *J. Veg. Sci.* 15: 437–448.
- Ollinger, S., Smith, M., Martin, M., Hallett, R., Goodale, C. and Aber, J. (2002). Regional variation in foliar chemistry and N cycling among forests of diverse history and composition. *Ecology* 83: 339–355.
- Onianwa, P. (2001). Monitoring atmospheric metal pollution: A review of the use of mosses as indicators. *Environ. Monit. Assess.* 71: 13–50.
- Orecchio, S. (2007). PAHs associated with the leaves of *Quercus ilex* L.: Extraction, GC-MS analysis, distribution and sources: Assessment of air quality in the Palermo (Italy) area. *Atmos. Environ.* 41: 8669–8680.
- Ormerod, S.J. and Durance, I. (2009). Restoration and recovery from acidification in upland Welsh streams over 25 years. *J. Appl. Ecol.* 46: 164–174.
- Ötvös, E., Kozák, I.O., Fekete, J., Sharma, V.K. and Tuba, Z. (2004). Atmospheric deposition of polycyclic aromatic

- hydrocarbons (PAHs) in mosses (*Hypnum cupressiforme*) in Hungary. *Sci. Total Environ.* 330: 89–99.
- Ouimet, R., Arp, P.A., Watmough, S.A., Aherne, J. and DeMerchant, I. (2006). Determination and mapping critical loads of acidity and exceedances for upland forest soils in eastern Canada. *Water Air Soil Pollut.* 172: 57–66.
- Ouimet, R. (2008). Using compositional change within soil profiles for modelling base cation transport and chemical weathering. *Geoderma* 145: 410–418.
- Oulehle, F., Evans, C.D., Hofmeister, J., Krejci, R., Tahovska, K., Persson, T., Cudlin, P. and Hruska, J. (2011). Major changes in forest carbon and nitrogen cycling caused by declining sulphur deposition. *Global Change Biol.* 17: 3115–3129.
- Outridge, P.M. and Scheuhammer, A.M. (1993). Bioaccumulation and toxicology of chromium: Implications for wildlife. In *Reviews of environmental contamination and toxicology*, Ware, G.W. (Ed.), Springer New York, pp. 31–77.
- Ovesen, C.H. (2001). Management of Danish dunes today: theory and practice. In *Coastal dune management: shared experience of European conservation practice*. Liverpool University Press, Liverpool Houston, J.A. and Rooney, P.A. (Eds.), Liverpool University Press, Liverpool, pp: 302–308.
- Panigada, C., Rossini, M., Meroni, M., Marzuoli, R., Gerosa, G. and Colombo, R. (2009). Indicators of ozone effects on *Fagus sylvatica* L. by means of spectroradiometric measurements. *Riv. Ital. Telerilevamento* 41: 3–20.
- Panno, S.V., Hackley, K.C., Hwang, H.H., Greenberg, S.E., Krapac, I.G., Landsberger, S. and O'Kelly, D.J. (2006). Characterization and identification of Na-Cl sources in ground water. *Ground Water* 44: 176–187.
- Papa, S., Bartoli, G., Nacca, F., D'Ambrosia, B., Cembrola, E., Pellegrino, A., Fiorentino, A., Fuggi, A. and Fioretto, A. (2012). Trace metals, peroxidase activity, PAHs contents and ecophysiological changes in *Quercus ilex* leaves in the urban area of Caserta (Italy). *J. Environ. Manage.* 113: 501–509.
- Peakall, D.W. and Walker, C.H. (1994). The role of biomarkers in environmental assessment. *Ecotoxicology* 3: 173–179.
- Penglase, S., Hamre, K. and Ellingsen, S. (2014). Selenium and mercury have a synergistic negative effect on fish reproduction. *Aquat. Toxicol.* 149: 16–24.
- Philips, F.S., Sirotnak, F.M., Sodergren, J.E. and Hutchison, D.J. (1973). Uptake of methotrexate, aminopterin, and methasquin and inhibition of dihydrofolate reductase and of DNA synthesis in mouse small intestine. *Cancer Res.* 33: 153–158.
- Phillips, D. (1985). Organochlorines and trace metals in green-lipped mussels *Perna viridis* from Hong Kong waters: A test of indicator ability. *Mar. Ecol. Prog. Ser.* 21: 251–258.
- Piccardo, M.T., Pala, M., Bonaccorso, B., Stella, A., Redaelli, A., Paola, G. and Valerio, F. (2005). *Pinus nigra* and *Pinus pinaster* needles as passive samplers of polycyclic aromatic hydrocarbons. *Environ. Pollut.* 133: 293–301.
- Piirainen, S., Finér, L. and Starr, M. (2002). Deposition and leaching of sulphate and base cations in a mixed boreal forest in Eastern Finland. *Water Air Soil Pollut.* 133: 185–204.
- Pilkington, M.G., Caporn, S.J., Carroll, J.A., Cresswell, N., Lee, J.A., Emmett, B.A. and Johnson, D. (2005). Effects of increased deposition of atmospheric nitrogen on an upland Calluna moor: N and P transformations. *Environ. Pollut.* 135: 469–480.
- Pitcairn, C., Leith, I., Sheppard, L., Sutton, M., Fowler, D., Munro, R., Tang, S. and Wilson, D. (1998). The relationship between nitrogen deposition, species composition and foliar nitrogen concentrations in woodland flora in the vicinity of livestock farms. *Environ. Pollut.* 102: 41–48.
- Pitcairn, C., Skiba, U., Sutton, M., Fowler, D., Munro, R. and Kennedy, V. (2002). Defining the spatial impacts of poultry farm ammonia emissions on species composition of adjacent woodland groundflora using Ellenberg Nitrogen Index, nitrous oxide and nitric oxide emissions and foliar nitrogen as marker variables. *Environ. Pollut.* 119: 9–21.
- Pitcairn, C., Fowler, D., Leith, I., Sheppard, L., Tang, S., Sutton, M. and Famulari, D. (2006). Diagnostic indicators of elevated nitrogen deposition. *Environ. Pollut.* 144: 941–950.
- Pitcairn, C.E., Leith, I.D., Fowler, D., Hargreaves, K.J., Moghaddam, M., Kennedy, V.H. and Granat, L. (2001). Foliar nitrogen as an indicator of nitrogen deposition and critical loads exceedance on a European scale. *Water Air Soil Pollut.* 130: 1037–1042.
- Pitcairn, C.E., Fowler, D., Leith, I.D., Sheppard, L.J., Sutton, M.A., Kennedy, V. and Okello, E. (2003). Bioindicators of enhanced nitrogen deposition. *Environ. Pollut.* 126: 353–361.
- Poursafa, P., Kelishadi, R., Lahijanzadeh, A., Modaresi, M., Javanmard, S.H., Assari, R., Amin, M.M., Moattar, F., Amini, A. and Sadeghian, B. (2011). The relationship of air pollution and surrogate markers of endothelial dysfunction in a population-based sample of children. *BMC Public Health* 11: 115.
- Power, S., Ashmore, M., Cousins, D. and Ainsworth, N. (1995). Long term effects of enhanced nitrogen deposition on a lowland dry heath in southern Britain. *Water Air Soil Pollut.* 85: 1701–1706.
- Power, S.A., Green, E.R., Barker, C.G., BELL, J.N.B. and Ashmore, M.R. (2006). Ecosystem recovery: Heathland response to a reduction in nitrogen deposition. *Global Change Biol.* 12: 1241–1252.
- Power, S.A. and Collins, C.M. (2010). Use of *Calluna vulgaris* to detect signals of nitrogen deposition across an urban-rural gradient. *Atmos. Environ.* 44: 1772–1780.
- Prajapati, S.K. and Tripathi, B. (2008). Biomonitoring seasonal variation of urban air polycyclic aromatic hydrocarbons (PAHs) using *Ficus benghalensis* leaves. *Environ. Pollut.* 151: 543–548.
- Rabalais, N.N. (2002). Nitrogen in aquatic ecosystems. *Ambio* 31: 102–112.

- Rajkowska, M. and Protasowicki, M. (2013). Distribution of metals (Fe, Mn, Zn, Cu) in fish tissues in two lakes of different trophy in Northwestern Poland. *Environ. Monit. Assess.* 185: 3493–3502.
- Rask, M., Mannio, J., Forsius, M., Posch, M. and Vuorinen, P.J. (1995). How many fish populations in Finland are affected by acid precipitation? *Environ. Biol. Fishes* 42: 51–63.
- Rasmussen, P.W., Schrank, C.S. and Campfield, P.A. (2007). Temporal trends of mercury concentrations in Wisconsin walleye (*Sander vitreus*), 1982–2005. *Ecotoxicology* 16: 541–550.
- Ratola, N., Amigo, J.M. and Alves, A. (2010). Comprehensive assessment of pine needles as bioindicators of PAHs using multivariate analysis. The importance of temporal trends. *Chemosphere* 81: 1517–1525.
- Ratola, N., Amigo, J.M., Oliveira, M.S., Araújo, R., Silva, J.A. and Alves, A. (2011). Differences between *Pinus pinea* and *Pinus pinaster* as bioindicators of polycyclic aromatic hydrocarbons. *Environ. Exp. Bot.* 72: 339–347.
- Ratola, N., Alves, A., Lacorte, S. and Barceló, D. (2012). Distribution and sources of PAHs using three pine species along the Ebro River. *Environ. Monit. Assess.* 184: 985–999.
- Rauert, C., Kananathalingam, A. and Harner, T. (2017). Characterization and modeling of polycyclic aromatic compound uptake into spruce tree wood. *Environ. Sci. Technol.* 51: 5287–5295.
- Rhodes, C., Bingham, A., Heard Andrea, M., Hewitt, J., Lynch, J., Waite, R. and Bell Michael, D. (2017). Diatoms to human uses: Linking nitrogen deposition, aquatic eutrophication, and ecosystem services. *Ecosphere* 8: e01858.
- Ribas, A., Filella, I., Gimeno, B.S. and Peñuelas, J. (1998). Evaluation of tobacco cultivars as bioindicators and biomonitors of ozone phytotoxic levels in Catalonia. *Water Air Soil Pollut.* 107: 347–365.
- Rice, K.M., Walker Jr, E.M., Wu, M., Gillette, C. and Blough, E.R. (2014). Environmental mercury and its toxic effects. *J. Preventative Med. Public Health* 47: 74.
- Richard Pilsner, J., Lazarus Alicia, L., Nam, D.H., Letcher Robert, J., Sonne, C., Dietz, R. and Basu, N. (2010). Mercury-associated DNA hypomethylation in polar bear brains via the LUMinometric Methylation Assay: A sensitive method to study epigenetics in wildlife. *Mol. Ecol.* 19: 307–314.
- Richardson, D., Davies, I., Moffat, C., Pollard, P. and Stagg, R. (2001). Biliary PAH metabolites and EROD activity in flounder (*Platichthys flesus*) from a contaminated estuarine environment. *J. Environ. Monit.* 3: 610–615.
- Richardson, D.M., Gubbins, M.J., Davies, I.M., Moffat, C.F. and Pollard, P.M. (2004). Effects of feeding status on biliary PAH metabolite and biliverdin concentrations in plaice (*Pleuronectes platessa*). *Environ. Toxicol. Pharmacol.* 17: 79–85.
- Rodríguez-Estival, J. and Smits, J.E. (2016). Small mammals as sentinels of oil sands related contaminants and health effects in northeastern Alberta, Canada. *Ecotoxicol. Environ. Saf.* 124: 285–295.
- Rose, M., Fernandes, A., Foxall, C. and Dowding, A. (2012). Transfer and uptake of polychlorinated dibenzo-p-dioxins and furans (PCDD/Fs) and polychlorinated biphenyls (PCBs) into meat and organs of indoor and outdoor reared pigs. *Food Addit. Contam. A* 29: 431–448.
- Rosseland, B., Eldhuset, T.D. and Staurnes, M. (1990). Environmental effects of aluminium. *Environ. Geochem. Health* 12: 17–27.
- Rowe, E., Jones, L., Dise, N., Evans, C., Mills, G., Hall, J., Stevens, C.J., Mitchell, R., Field, C. and Caporn, S. (2016). Metrics for evaluating the ecological benefits of decreased nitrogen deposition. *Biol. Conserv.* 212: 454–463.
- Ruddock, P., Bird, D.J., McEvoy, J. and Peters, L. (2003). Bile metabolites of polycyclic aromatic hydrocarbons (PAHs) in European eels *Anguilla anguilla* from United Kingdom estuaries. *Sci. Total Environ.* 301: 105–117.
- Rybak, J. and Olejniczak, T. (2014). Accumulation of polycyclic aromatic hydrocarbons (PAHs) on the spider webs in the vicinity of road traffic emissions. *Environ. Sci. Pollut. Res.* 21: 2313–2324.
- Salman, J.M., Al-Azawey, A.S. and Hassan, F.M. (2014). The study of pollution of PAHs (polycyclic aromatic hydrocarbons) in Al-Hilla River, Iraq by using bioindicator freshwater crab (*Sesarma Boulengeri* Calman). *J. Life Sci.* 8: 351–357.
- Sandermann, H. (2000). Ozone/biotic disease interactions: molecular biomarkers as a new experimental tool. *Environ. Pollut.* 108: 327–332.
- Saros, J.E., Clow, D.W., Blett, T. and Wolfe, A.P. (2011). Critical nitrogen deposition loads in high-elevation lakes of the western US inferred from paleolimnological records. *Water Air Soil Pollut.* 216: 193–202.
- Satya, Upreti, D.K. and Patel, D. (2012). Rinodina sophodes (Ach.) Massal.: A bioaccumulator of polycyclic aromatic hydrocarbons (PAHs) in Kanpur City, India. *Environ. Monit. Assess.* 184: 229–238.
- Savva, Y. and Berninger, F. (2010). Sulphur deposition causes a large-scale growth decline in boreal forests in Eurasia. *Global Biogeochem. Cycles* 24: GB3002.
- Scheuhammer, A.M., Meyer, M.W., Sandheinrich, M.B. and Murray, M.W. (2007). Effects of environmental methylmercury on the health of wild birds, mammals, and fish. *Ambio* 36: 12–19.
- Schiedek, D., Broeg, K., Baršienė, J., Lehtonen, K.K., Gercken, J., Pfeifer, S., Vuontisjärvi, H., Vuorinen, P.J., Dedonate, V. and Koehler, A. (2006). Biomarker responses as indication of contaminant effects in blue mussel (*Mytilus edulis*) and female eelpout (*Zoarces viviparus*) from the southwestern Baltic Sea. *Mar. Pollut. Bull.* 53: 387–405.
- Schilling, J.S. and Lehman, M.E. (2002). Bioindication of atmospheric heavy metal deposition in the Southeastern US using the moss *Thuidium delicatulum*. *Atmos. Environ.* 36: 1611–1618.
- Schintu, M., Cogoni, A., Durante, L., Cantaluppi, C. and

- Contu, A. (2005). Moss (*Bryum radiculosum*) as a bioindicator of trace metal deposition around an industrialised area in Sardinia (Italy). *Chemosphere* 60: 610–618.
- Schmeltz, D., Evers, D.C., Driscoll, C.T., Artz, R., Cohen, M., Gay, D., Haeuber, R., Krabbenhoft, D.P., Mason, R., Morris, K. and Wiener, J.G. (2011). MercNet: A national monitoring network to assess responses to changing mercury emissions in the United States. *Ecotoxicology* 20: 1713–1725.
- Schober, S.E., Sinks, T.H., Jones, R.L., Bolger, P.M., McDowell, M., Osterloh, J., Garrett, E.S., Canady, R.A., Dillon, C.F. and Sun, Y. (2003). Blood mercury levels in US children and women of childbearing age, 1999–2000. *JAMA* 289: 1667–1674.
- Schoeman, K., Bend, J.R., Hill, J., Nash, K. and Koren, G. (2009). Defining a lowest observable adverse effect hair concentrations of mercury for neurodevelopmental effects of prenatal methylmercury exposure through maternal fish consumption: A systematic review. *Ther. Drug Monit.* 31: 670–682.
- Schröter-Kermani, C., Kreft, D., Schilling, B., Herrchen, M. and Wagner, G. (2006). Polycyclic aromatic hydrocarbons in pine and spruce shoots—temporal trends and spatial distribution. *J. Environ. Monit.* 8: 806–811.
- Schuster, J.K., Harner, T., Su, K., Mihele, C. and Eng, A. (2015). First results from the oil sands passive air monitoring network for polycyclic aromatic compounds. *Environ. Sci. Technol.* 49: 2991–2998.
- Seinfeld, J.H. and Pandis, S.N. (2016). *Atmospheric chemistry and physics: From Air pollution to climate change*. John Wiley & Sons.
- Shao, Y., Zhang, W., Eisenhauer, N., Liu, T., Xiong, Y., Liang, C. and Fu, S. (2017). Nitrogen deposition cancels out exotic earthworm effects on plant-feeding nematode communities. *J. Anim. Ecol.* 86: 708–717.
- Sharma, A.P. and Tripathi, B. (2009). Assessment of atmospheric PAHs profile through *Calotropis gigantea* R.Br. leaves in the vicinity of an Indian coal-fired power plant. *Environ. Monit. Assess.* 149: 477–482.
- Sheibley, R.W., Enache, M., Swarzenski, P.W., Moran, P.W. and Foreman, J.R. (2014). Nitrogen deposition effects on diatom communities in lakes from three national parks in Washington state. *Water Air Soil Pollut.* 225: 1857.
- Shepherd, S.A., Watson, J.E., Womersley, H.B.S. and Carey, J.M. (2009). Long-term changes in macroalgal assemblages after increased sedimentation and turbidity in Western Port, Victoria, Australia. *Bot. Mar.* 52: 195–206.
- Sheppard, L.J., Leith, I.D., Mizunuma, T., Neil Cape, J., Crossley, A., Leeson, S., Sutton, M.A., Dijk, N. and Fowler, D. (2011). Dry deposition of ammonia gas drives species change faster than wet deposition of ammonium ions: Evidence from a long-term field manipulation. *Global Change Biol.* 17: 3589–3607.
- Sherlock, J., Hislop, J., Newton, D., Topping, G. and Whittle, K. (1984). Elevation of mercury in human blood from controlled chronic ingestion of methylmercury in fish. *Hum. Toxicol.* 3: 117–131.
- Shotyk, W., Belland, R., Duke, J., Kempster, H., Krachler, M., Noernberg, T., Pelletier, R., Vile, M.A., Wieder, K. and Zaccone, C. (2014). Sphagnum mosses from 21 ombrotrophic bogs in the athabasca bituminous sands region show no significant atmospheric contamination of "heavy metals". *Environ. Sci. Technol.* 48: 12603–12611.
- Shukla, V. and Upreti, D. (2009). Polycyclic aromatic hydrocarbon (PAH) accumulation in lichen, *Phaeophyscia hispidula* of DehraDun City, Garhwal Himalayas. *Environ. Monit. Assess.* 149: 1–7.
- Shukla, V., Patel, D., Upreti, D. and Yunus, M. (2012). Lichens to distinguish urban from industrial PAHs. *Environ. Chem. Lett.* 10: 159–164.
- Simkin, S.M., Allen, E.B., Bowman, W.D., Clark, C.M., Belnap, J., Brooks, M.L., Cade, B.S., Collins, S.L., Geiser, L.H., Gilliam, F.S., Jovan, S.E., Pardo, L.H., Schulz, B.K., Stevens, C.J., Suding, K.N., Throop, H.L. and Waller, D.M. (2016). Conditional vulnerability of plant diversity to atmospheric nitrogen deposition across the United States. *Proc. Natl. Acad. Sci. U.S.A.* 113: 4086–4091.
- Simon, K.S., Simon, M.A. and Benfield, E.F. (2009). Variation in ecosystem function in Appalachian streams along an acidity gradient. *Ecol. Appl.* 19: 1147–1160.
- Simoneau, M., Lucotte, M., Garceau, S. and Laliberté, D. (2005). Fish growth rates modulate mercury concentrations in walleye (*Sander vitreus*) from eastern Canadian lakes. *Environ. Res.* 98: 73–82.
- Singh, S., Srivastava, K., Gahtori, D. and Saxena, K.D. (2017). Bryomonitoring of atmospheric elements in *Rhodobryum giganteum* (Schwaegr.) Par., growing in Uttarakhand region of Indian Himalayas. *Aerosol Air Qual. Res.* 17: 810–820.
- Sjostedt, J., Koch-Schmidt, P., Pontarp, M., Canback, B., Tunlid, A., Lundberg, P., Hagstrom, A. and Riemann, L. (2012). Recruitment of members from the rare biosphere of marine bacterioplankton communities after an environmental disturbance. *Appl. Environ. Microbiol.* 78: 1361–1369.
- Skiba, U., Cresser, M., Derwent, R. and Futty, D. (1989). Peat acidification in Scotland. *Nature* 337: 68–69.
- Slaninova, A., Machova, J. and Svobodova, Z. (2014). Fish kill caused by aluminium and iron contamination in a natural pond used for fish rearing: A case report. *Vet. Med.* 59: 573–581.
- Smith, V.H. and Schindler, D.W. (2009). Eutrophication science: Where do we go from here? *Trends Ecol. Evol.* 24: 201–207.
- Sola, F., Isaia, J. and Masoni, A. (1995). Effects of copper on gill structure and transport function in the rainbow trout, *Oncorhynchus mykiss*. *J. Appl. Toxicol.* 15: 391–398.
- Soons, M.B., Hefting, M.M., Dorland, E., Lamers, L.P., Versteeg, C. and Bobbink, R. (2016). Nitrogen effects on plant species richness in herbaceous communities are more widespread and stronger than those of phosphorus. *Biol. Conserv.* 212: 390–397.

- Southon, G.E., Field, C., Caporn, S.J., Britton, A.J. and Power, S.A. (2013). Nitrogen deposition reduces plant diversity and alters ecosystem functioning: Field-scale evidence from a nationwide survey of UK heathlands. *PLoS one* 8: e59031.
- Sparling, D.W. and Lowe, T.P. (1996). Environmental hazards of aluminum to plants, invertebrates, fish, and wildlife. In *Reviews of environmental contamination and toxicology*, Ware, G.W. (Ed.), Springer, pp. 1–127.
- Steinnes, E. (1995). A critical evaluation of the use of naturally growing moss to monitor the deposition of atmospheric metals. *Sci. Total Environ.* 160: 243–249.
- Stevens, C., Dise, N.B. and Gowing, D.J. (2009a). Regional trends in soil acidification and exchangeable metal concentrations in relation to acid deposition rates. *Environ. Pollut.* 157: 313–319.
- Stevens, C., Maskell, L., Smart, S., Caporn, S., Dise, N. and Gowing, D. (2009b). Identifying indicators of atmospheric nitrogen deposition impacts in acid grasslands. *Biol. Conserv.* 142: 2069–2075.
- Stevens, C.J., Dise, N.B., Mountford, J.O. and Gowing, D.J. (2004). Impact of nitrogen deposition on the species richness of grasslands. *Science* 303: 1876–1879.
- Stevens, C.J., Dise, N.B., Gowing, D.J. and Mountford, J. (2006). Loss of forb diversity in relation to nitrogen deposition in the UK: Regional trends and potential controls. *Global Change Biol.* 12: 1823–1833.
- Stevens, C.J., Smart, S.M., Henrys, P.A., Maskell, L.C., Crowe, A., Simkin, J., Cheffings, C.M., Whitfield, C., Gowing, D.J. and Rowe, E.C. (2012). Terricolous lichens as indicators of nitrogen deposition: Evidence from national records. *Ecol. Indic.* 20: 196–203.
- Stevens, C.J. (2016). How long do ecosystems take to recover from atmospheric nitrogen deposition? *Biol. Conserv.* 200: 160–167.
- Stoken, O.M., Riscassi, A.L. and Scanlon, T.M. (2016). Association of dissolved mercury with dissolved organic carbon in U.S. rivers and streams: The role of watershed soil organic carbon. *Water Resour. Res.* 52: 3040–3051.
- Strengbom, J., Nordin, A., Näsholm, T. and Ericson, L. (2001). Slow recovery of boreal forest ecosystem following decreased nitrogen input. *Funct. Ecol.* 15: 451–457.
- Stürzenbaum, S.R., Georgiev, O., Morgan, A.J. and Kille, P. (2004). Cadmium detoxification in earthworms: From genes to cells. *Environ. Sci. Technol.* 38: 6283–6289.
- Su, Y., Wania, F., Harner, T., and Lei, Y.D. (2007). Deposition of polybrominated diphenyl ethers, polychlorinated biphenyls, and polycyclic aromatic hydrocarbons to a boreal deciduous forest. *Environ. Sci. Technol.* 41: 534–540.
- Sullivan, D.G., White, J.G. and Vepraskas, M.J. (2017). Using land-use change, soil characteristics, and a semi-automated on-line GIS database to inventory Carolina Bays. *Wetlands* 37: 89–98.
- Sun, F., Wen, D., Kuang, Y., Li, J., Li, J. and Zuo, W. (2010). Concentrations of heavy metals and polycyclic aromatic hydrocarbons in needles of Masson pine (*Pinus massoniana* L.) growing nearby different industrial sources. *J. Environ. Sci.* 22: 1006–1013.
- Sutton, M., Milford, C., Nemitz, E., Theobald, M., Hill, P., Fowler, D., Schjoerring, J., Mattsson, M., Nielsen, K.H. and Husted, S. (2001). Biosphere-atmosphere interactions of ammonia with grasslands: Experimental strategy and results from a new European initiative. *Plant Soil* 228: 131–145.
- Sutton, M., Pitcairn, C., Whitfield, C., Leith, I., Sheppard, L., van Dijk, N., Tang, S., Skiba, U., Smart, S. and Mitchell, R. (2004). Bioindicator and biomonitoring methods for assessing the effects of atmospheric nitrogen on statutory nature conservation sites. JNCC Report No: 356, Countryside Council for Wales, English Nature, Joint Nature Conservation Committee and Centre for Ecology and Hydrology, Peterborough.
- Szczeklik, A., Szczeklik, J., Galuszka, Z., Musial, J., Kolarzyk, E. and Targosz, D. (1994). Humoral immunosuppression in men exposed to polycyclic aromatic hydrocarbons and related carcinogens in polluted environments. *Environ. Health Perspect.* 102: 302.
- Takeda, A., Minami, A., Seki, Y. and Oku, N. (2004). Differential effects of zinc on glutamatergic and GABAergic neurotransmitter systems in the hippocampus. *J. Neurosci. Res.* 75: 225–229.
- Takeuchi, T., Nakajima, M. and Morimoto, K. (1996). ACCELERATED PAPER: Relationship between the intracellular reactive oxygen species and the induction of oxidative DNA damage in human neutrophil-like cells. *Carcinogenesis* 17: 1543–1548.
- Tarrasón, L., Fagerli, H., Klein, H., Simpson, D., Benedictow, A.C., Vestreng, V., Rigler, E., Emberson, L., Posch, M. and Spranger, T. (2006). *Transboundary Acidification, Eutrophication and Ground Level Ozone in Europe from 1990 to 2004 in support for the review of the Gothenburg Protocol*, EMEP Status Report 1/2006, Meteorological Institute, Oslo.
- Thwaites, R., Ashmore, M., Morton, A. and Pakeman, R. (2006). The effects of tropospheric ozone on the species dynamics of calcareous grassland. *Environ. Pollut.* 144: 500–509.
- Tian, D. and Niu, S. (2015). A global analysis of soil acidification caused by nitrogen addition. *Environ. Res. Lett.* 10: 024019.
- Tomashuk, T.A., Truong, T.M., Mantha, M. and McGowin, A.E. (2012). Atmospheric polycyclic aromatic hydrocarbon profiles and sources in pine needles and particulate matter in Dayton, Ohio, USA. *Atmos. Environ.* 51: 196–202.
- Trzcinka-Ochocka, M., Jakubowski, M., Szymczak, W., Janasik, B. and Brodzka, R. (2010). The effects of low environmental cadmium exposure on bone density. *Environ. Res.* 110: 286–293.
- Tsipoura, N., Burger, J., Newhouse, M., Jeitner, C., Gochfeld, M. and Mizrahi, D. (2011). Lead, mercury, cadmium, chromium, and arsenic levels in eggs, feathers, and tissues of Canada geese of the New Jersey Meadowlands. *Environ. Res.* 111: 775–784.
- US EPA (2017). Chemicals and Toxics Topics, <https://www.epa.gov/environmental-topics/chemicals-and-toxics-topics>, Last Access: 20 March 2018.

- Vachirapatama, N., Jirakiattiku, Y., Dicinoski, G., Townsend, A.T. and Haddad, P.R. (2011). Effect of vanadium on plant growth and its accumulation in plant tissues. *Songklanakarinn J. Sci. Technol.* 33: 255–261.
- Vainonen, J.P. and Kangasjärvi, J. (2015). Plant signalling in acute ozone exposure. *Plant Cell Environ.* 38: 240–252.
- van den Berg, L.J., Jones, L., Sheppard, L.J., Smart, S.M., Bobbink, R., Dise, N.B. and Ashmore, M.R. (2016). Evidence for differential effects of reduced and oxidised nitrogen deposition on vegetation independent of nitrogen load. *Environ. Pollut.* 208: 890–897.
- van den Elzen, E., van den Berg, L.J.L., van der Weijden, B., Fritz, C., Sheppard, L.J. and Lamers, L.P.M. (2018). Effects of airborne ammonium and nitrate pollution strongly differ in peat bogs, but symbiotic nitrogen fixation remains unaffected. *Sci. Total Environ.* 610–611: 732–740.
- Van der Molen, D., Portielje, R., De Nobel, W. and Boers, P. (1998). Nitrogen in dutch freshwater lakes: Trends and targets. *Environ. Pollut.* 102: 553–557.
- van der Sleen, P., Vlam, M., Groenendijk, P., Anten, N.P., Bongers, F., Bunyavejchewin, S., Hietz, P., Pons, T.L. and Zuidema, P.A. (2015). <sup>15</sup>N in tree rings as a bio-indicator of changing nitrogen cycling in tropical forests: An evaluation at three sites using two sampling methods. *Front. Plant Sci.* 6: 229.
- Van Dingenen, R., Dentener, F.J., Raes, F., Krol, M.C., Emberson, L. and Cofala, J. (2009). The global impact of ozone on agricultural crop yields under current and future air quality legislation. *Atmos. Environ.* 43: 604–618.
- van Dobben, H. and de Vries, W. (2010). Relation between forest vegetation, atmospheric deposition and site conditions at regional and European scales. *Environ. Pollut.* 158: 921–933.
- Van Wijnen, H. and Bakker, J. (1999). Nitrogen and phosphorus limitation in a coastal barrier salt marsh: the implications for vegetation succession. *J. Ecol.* 87: 265–272.
- VandenBoer, T.C., Brown, S.S., Murphy, J.G., Keene, W.C., Young, C.J., Pszenny, A., Kim, S., Warneke, C., Gouw, J.A. and Maben, J.R. (2013). Understanding the role of the ground surface in HONO vertical structure: High resolution vertical profiles during NACHTT-11. *J. Geophys. Res.* 118: 10155–10171.
- Vanguelova, E.I., Hirano, Y., Eldhuset, T.D., Sas-Paszt, L., Bakker, M.R., Püttsepp, Ü., Brunner, I., Löhmus, K. and Godbold, D. (2007). Tree fine root Ca/Al molar ratio – Indicator of Al and acidity stress. *Plant Biosyst.* 141: 460–480.
- Váňová, L., Kummerová, M., Klemš, M. and Zedulka, Š. (2009). Fluoranthene influences endogenous abscisic acid level and primary photosynthetic processes in pea (*Pisum sativum* L.) plants in vitro. *Plant Growth Regul.* 57: 39–47.
- Verdin, A., Sahraoui, A.L.H., Fontaine, J., Grandmougin-Ferjani, A. and Durand, R. (2006). Effects of anthracene on development of an arbuscular mycorrhizal fungus and contribution of the symbiotic association to pollutant dissipation. *Mycorrhiza* 16: 397–405.
- Verhoeven, J., Beltman, B., Dorland, E., Robat, S. and Bobbink, R. (2011). Differential effects of ammonium and nitrate deposition on fen phanerogams and bryophytes. *Appl. Veg. Sci.* 14: 149–157.
- Vet, R., Artz, R.S., Carou, S., Shaw, M., Ro, C.U., Aas, W., Baker, A., Bowersox, V.C., Dentener, F. and Galy-Lacaux, C. (2014). A global assessment of precipitation chemistry and deposition of sulfur, nitrogen, sea salt, base cations, organic acids, acidity and pH, and phosphorus. *Atmos. Environ.* 93: 3–100.
- Villányi, V., Ürmös, Z., Balogh, J., Horváth, L., Csintalan, Z. and Tuba, Z. (2008). Ozone biomonitoring at mountainous and lowland areas in Hungary. *Acta Biologica Szegediensis* 52: 209–212.
- Vinodhini, R. and Narayanan, M. (2009). The impact of toxic heavy metals on the hematological parameters in common carp (*Cyprinus Carpio* L.). *J. Environ. Health Sci. Eng.* 6: 23–28.
- Vitousek, P.M., Mooney, H.A., Lubchenco, J. and Melillo, J.M. (1997). Human domination of earth's ecosystems. *Science* 277: 494–499.
- Vodyanitskii, Y.N. (2013). Contamination of soils with heavy metals and metalloids and its ecological hazard (analytic review). *Eurasian Soil Sci.* 46: 793–801.
- Vogels, J., Verberk, W., Lamers, L. and Siepel, H. (2016). Can changes in soil biochemistry and plant stoichiometry explain loss of animal diversity of heathlands? *Biol. Conserv.* 212: 432–447.
- Volk, M., Bungener, P., Contat, F., Montani, M. and Fuhrer, J. (2006). Grassland yield declined by a quarter in 5 years of free-air ozone fumigation. *Global Change Biol.* 12: 74–83.
- von Bobrutski, K., Braban, C., Famulari, D., Jones, S., Blackall, T., Smith, T., Blom, M., Coe, H., Gallagher, M. and Ghalaieny, M. (2010). Field inter-comparison of eleven atmospheric ammonia measurement techniques. *Atmos. Meas. Tech* 3: 91–112.
- Vuorinen, P.J., Keinänen, M., Vuontisjärvi, H., Baršienė, J., Broeg, K., Förlin, L., Gercken, J., Kopecka, J., Köhler, A. and Parkkonen, J. (2006). Use of biliary PAH metabolites as a biomarker of pollution in fish from the Baltic Sea. *Mar. Pollut. Bull.* 53: 479–487.
- Wærvågen, S.B., A Rukke, N. and Hessen, D.O. (2002). Calcium content of crustacean zooplankton and its potential role in species distribution. *Freshwater Biol.* 47: 1866–1878.
- WallisDeVries, M.F. and van Swaay, C.A. (2016). A nitrogen index to track changes in butterfly species assemblages under nitrogen deposition. *Biol. Conserv.* 212: 448–453.
- Wang, J. and Liu, Z. (1999). Effect of vanadium on the growth of soybean seedlings. *Plant Soil* 216: 47–51.
- Wang, Z., Ma, X., Na, G., Lin, Z., Ding, Q. and Yao, Z. (2009). Correlations between physicochemical properties of PAHs and their distribution in soil, moss and reindeer dung at Ny-Alesund of the Arctic. *Environ. Pollut.* 157: 3132–3136.

- Watmough, S.A., Whitfield, C.J. and Fenn, M.E. (2014). The importance of atmospheric base cation deposition for preventing soil acidification in the Athabasca Oil Sands Region of Canada. *Sci. Total Environ.* 493: 1–11.
- Watras, C.J. and Huckabee, J.W. (1994). *Mercury pollution integration and synthesis*. CRC Press.
- Watt, W., Scott, C., Zamora, P. and White, W. (2000). Acid toxicity levels in Nova Scotian rivers have not declined in synchrony with the decline in sulfate levels. *Water Air Soil Pollut.* 118: 203–229.
- Weissengruber, L., Möller, K., Puschenreiter, M. and Friedel, J.K. (2018). Long-term soil accumulation of potentially toxic elements and selected organic pollutants through application of recycled phosphorus fertilizers for organic farming conditions. *Nutr. Cycling Agroecosyst.* 110: 427–449.
- White, D., Stair, J. and Ringelberg, D. (1996). Quantitative comparisons of in situ microbial biodiversity by signature biomarker analysis. *J. Ind. Microbiol. Biotechnol.* 17: 185–196.
- Whitfield, C.J., Watmough, S.A. and Aherne, J. (2011). Evaluation of elemental depletion weathering rate estimation methods on acid-sensitive soils of north-eastern Alberta, Canada. *Geoderma* 166: 189–197.
- WHO (2011). *Guidelines for drinking-water quality*, 4<sup>th</sup> Edition. World Health Organization, Geneva.
- WHO Regional Publications, European Series (2000). Effects of ozone on vegetation: Critical levels in Air Quality Guidelines Second Edition No. 91, [http://www.euro.who.int/\\_data/assets/pdf\\_file/0006/123099/AQG2ndEd\\_12ozonelevel.pdf?ua=1](http://www.euro.who.int/_data/assets/pdf_file/0006/123099/AQG2ndEd_12ozonelevel.pdf?ua=1), Last Access 30 January 2018.
- Wieczorek, J. and Wieczorek, Z. (2007). Phytotoxicity and accumulation of anthracene applied to the foliage and sandy substrate in lettuce and radish plants. *Ecotoxicol. Environ. Saf.* 66: 369–377.
- Wieder, R.K., Vile, M.A., Scott, K.D., Albright, C.M., McMillen, K.J., Vitt, D.H. and Fenn, M.E. (2016). Differential effects of high atmospheric N and S deposition on bog plant/lichen tissue and porewater chemistry across the Athabasca Oil Sands Region. *Environ. Sci. Technol.* 50: 12630–12640.
- Wiener, J.G., Evers, D.C., Gay, D.A., Morrison, H.A. and Williams, K.A. (2012). Mercury contamination in the Laurentian Great Lakes region: Introduction and overview. *Environ. Pollut.* 161: 243–251.
- Wild, S., Berrow, M., McGrath, S. and Jones, K. (1992). Polynuclear aromatic hydrocarbons in crops from long-term field experiments amended with sewage sludge. *Environ. Pollut.* 76: 25–32.
- Wilkins, K. and Aherne, J. (2016). Vegetation community change in Atlantic oak woodlands along a nitrogen deposition gradient. *Environ. Pollut.* 216: 115–124.
- Will, M. and Suter, G.W. (1995). Toxicological benchmarks for potential contaminants of concern for effects on soil and litter invertebrates and heterotrophic process. Report ES/ER/TM-126/R1, Oak Ridge National Lab., TN, USA.
- Williams, P.L. and Burson, J.L. (1985). *Industrial toxicology: Safety and health applications in the workplace*. Lifetime Learning Publications, New York.
- Wilmot, T.R., Ellsworth, D.S. and Tyree, M.T. (1995). Relationships among crown condition, growth, and stand nutrition in seven northern Vermont sugarbushes. *Can. J. For. Res.* 25: 386–397.
- Wittig, V.E., Ainsworth, E.A., Naidu, S.L., Karnosky, D.F. and Long, S.P. (2009). Quantifying the impact of current and future tropospheric ozone on tree biomass, growth, physiology and biochemistry: A quantitative meta-analysis. *Global Change Biol.* 15: 396–424.
- Wnorowski, A. and Charland, J.P. (2017). Profiling quinones in ambient air samples collected from the Athabasca region (Canada). *Chemosphere* 189: 55–66.
- Wolfe, M.F., Schwarzbach, S. and Sulaiman, R.A. (1998). Effects of mercury on wildlife: A comprehensive review. *Environ. Toxicol. Chem.* 17: 146–160.
- Wright, L.P., Zhang, L. and Marsik, F.J. (2016). Overview of mercury dry deposition, litterfall, and throughfall studies. *Atmos. Chem. Phys.* 16: 13399–13416.
- Wynn, P.M., Loader, N.J. and Fairchild, I. (2014). Interrogating trees for isotopic archives of atmospheric sulphur deposition and comparison to speleothem records. *Environ. Pollut.* 187: 98–105.
- Xia, M., Talhelm, A.F. and Pregitzer, K.S. (2017). Chronic nitrogen deposition influences the chemical dynamics of leaf litter and fine roots during decomposition. *Soil Biol. Biochem.* 112: 24–34.
- Xu, Y., Xiao, H. and Qu, L. (2017). Nitrogen concentrations and nitrogen isotopic compositions in leaves of *Cinnamomum Camphora* and *Pinus massoniana* (Lamb.) for indicating atmospheric nitrogen deposition in Guiyang (SW China). *Atmos. Environ.* 159: 1–10.
- Yan, J., Wang, L., Fu, P.P. and Yu, H. (2004). Photomutagenicity of 16 polycyclic aromatic hydrocarbons from the US EPA priority pollutant list. *Mutat. Res.* 557: 99–108.
- Yang, P., Chen, J., Wang, Z., Qiao, X., Cai, X., Tian, F. and Ge, L. (2007). Contributions of deposited particles to pine needle polycyclic aromatic hydrocarbons. *J. Environ. Monit.* 9: 1248–1253.
- Yasuhira, K. (1964). Damage to the thymus and other lymphoid tissues from 3-methylcholanthrene, and subsequent thymoma production, in mice. *Cancer Res.* 24: 558–569.
- Zambelli, B. and Ciurli, S. (2013). Nickel and human health. In *Interrelations between essential metal ions and human diseases*, Sigel, A., Sigel, H. and Sigel, R.K.O. (Eds.), Springer Netherlands, Dordrecht, pp. 321–357.
- Zeng, J., Zhang, G., Bao, L., Long, S., Tan, M., Li, Y., Ma, C. and Zhao, Y. (2013). Sulfur speciation and bioaccumulation in camphor tree leaves as atmospheric sulfur indicator analyzed by synchrotron radiation XRF and XANES. *J. Environ. Sci.* 25: 605–612.
- Zhang, L., Vet, R., Brook, J.R. and Leggae, A.H. (2006). Factors affecting stomatal uptake of ozone by different canopies and a comparison between dose and exposure. *Sci. Total Environ.* 370: 117–132.
- Zhang, L., Vet, R., O'Brien, J., Mihele, C., Liang, Z. and

- Wiebe, A. (2009). Dry deposition of individual nitrogen species at eight Canadian rural sites. *J. Geophys. Res.* 114.
- Zhang, L., Cheng, I., Wu, Z., Harner, T., Schuster, J., Charland, J.P., Muir, D. and Parnis J.M. (2015). Dry deposition of polycyclic aromatic compounds to various land covers in the Athabasca oil sands region. *J. Adv. Model. Earth Syst.* 7: 1339–1350.
- Zhao, Z.H., Quan, W.Y. and Tian, D.H. (1990). Urinary 1-hydroxypyrene as an indicator of human exposure to ambient polycyclic aromatic hydrocarbons in a coal-burning environment. *Sci. Total Environ.* 92: 145–154.
- Zhao, Z.H., Quan, W.Y. and Tian, D.H. (1992). Experiments on the effects of several factors on the 1-hydroxypyrene level in human urine as an indicator of exposure to polycyclic aromatic hydrocarbons. *Sci. Total Environ.* 113: 197–207.

*Received for review, March 25, 2018*

*Revised, June 20, 2018*

*Accepted, June 25, 2018*