



Characteristics of Individual Particles Emitted from an Experimental Burning Chamber with Coal from the Lung Cancer Area of Xuanwei, China

Wenhua Wang¹, Longyi Shao^{1*}, Jie Li¹, Lingli Chang¹, Daizhou Zhang², Chenchong Zhang³, Jingkun Jiang³

¹ State Key Laboratory of Coal Resources and Safe Mining & College of Geoscience and Surveying Engineering, China University of Mining and Technology (Beijing), Beijing 100083, China

² Faculty of Environmental and Symbiotic Sciences, Prefectural University of Kumamoto, Kumamoto 862-8502, Japan

³ State Key Joint Laboratory of Environmental Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing 100084, China

ABSTRACT

Pollutants emitted from household coal burning in Xuanwei, China, have been recognized as the reason for the high lung cancer mortality and morbidity rates in the area. To examine the characteristics of particles emitted from coal burning, a coal burning-dilution chamber was designed, and the individual particles emitted from the chamber at different burning stages were collected. The morphologies and elemental compositions of the individual particles were analyzed by high resolution transmission electron microscopy. Four types of particles, namely, organic particles, soot particles, S-rich particles, and mineral particles, were identified. The largest percentage of particles by number in the ignition stage, fierce-burning stage, and char-burning stage was composed of organic particles (66%), soot particles (71%), and mineral particles (73%), respectively. A distinctive characteristic was the remarkable abundance of Si- and Fe-rich particles during the char burning stage, compared with emissions from other types of coal. According to the elemental composition, 49% of the mineral particles were Si-rich, 25% were Ca-rich, 14% were Fe-rich, and 7% were Ti-rich. The Si-rich particles were partly identified as quartz (SiO₂), the Ca-rich particles were found to be CaSO₄ or CaCO₃, the Fe-rich particles were primarily Fe₂O₃ or Fe₃O₄, and the Ti-rich particles were mainly TiO₂. Notably, SiO₂ is a human carcinogen, and Fe-rich particles possess a high reactive potential with DNA-markers.

Keywords: Coal burning; Individual particle analysis; Si-rich particles; Fe-rich particles; Health risk.

INTRODUCTION

China both produced and consumed a large amount of coal every year. In 2016, about 2.6 billion tons of coal were consumed in China, which accounted for 62.0% of national primary energy source (National Bureau of Statistics of the People's Republic of China; <http://www.stats.gov.cn/tjsj/ndsj/>). Emissions from coal burning are important anthropogenic sources of particulate and gaseous pollutants in the atmosphere (Chen *et al.*, 2015; Li *et al.*, 2016a; Cai *et al.*, 2018). Coal contains many potentially harmful substances (Shao *et al.*, 2016; Finkelman and Tian, 2018). During the coal burning process, the harmful substances can be released into the air, causing adverse effect on environment and significantly influence human health (Zhang and Smith, 2007; Pian *et al.*, 2016).

Recently, industrial coal combustion shows relative low emission factors of particulate matter (PM) by installing air pollution control devices (Zhou *et al.*, 2016). However, the household coal burning shows distinct high PM emission factors due to the incomplete burning and the absence of dust control devices (Li *et al.*, 2016b). Emission factors of many pollutants from household stoves show two orders of magnitude higher than those from industrial boilers (Zhang *et al.*, 2008). Household coal burning for cooking and heating in Chinese rural areas is popular (Zhu *et al.*, 2012; Zhang *et al.*, 2014; Cai *et al.*, 2018), which causes high indoor PM pollution (Hu *et al.*, 2014), especially in wintertime (Li *et al.*, 2017). As a result, household coal burning can cause human health problems in some rural areas and has attracted attention in recent years (Chen *et al.*, 2015; Tiwari *et al.*, 2015; Li *et al.*, 2017; Lui *et al.*, 2017; Finkelman and Tian, 2018).

Xuanwei City, located in Yunnan Province, southwestern China, is rich in coal, iron, copper and other mines (Xiao *et al.*, 2012). Xuanwei has the highest lung cancer morbidity and mortality rates in China, especially in rural areas (He

* Corresponding author.

E-mail address: ShaoL@cumtb.edu.cn

et al., 1991; Xiao *et al.*, 2012; Kim *et al.*, 2014). Most of the important findings have suggested that the high lung cancer rates in Xuanwei are attributed to indoor smoky coal burning (Lan *et al.*, 2008; Barone-Adesi *et al.*, 2012; Hosgood *et al.*, 2013; Lui *et al.*, 2017). For example, according to a recent population-based case-control study, the lung cancer risk was significantly associated with the smoky coal use while the lung cancer association with cigarette was null in hazardous coal users (Kim *et al.*, 2014).

Knowledge of detailed physical and chemical characteristics of coal burning-derived fine particles has important significance in the field of explaining the mechanism of high lung cancer incidence in Xuanwei (Lu *et al.*, 2017). To our knowledge, there is few information available for revealing the evolution of individual fine particles throughout the different burning stages in Xuanwei. In this study, a set of chamber dilution measurement system was set up and the individual fine particles in different burning stages of Xuanwei coal were collected; characteristics of individual particles were analyzed by using transmission electron microscopy (TEM) with energy dispersive X-ray spectroscopy (EDX).

MATERIALS AND METHODS

Coal Burning-dilution Chamber

Fig. 1 shows the coal burning-dilution chamber measurement system used for generating coal-burning particles. The whole sealed room was $\sim 10 \text{ m}^3$ and at the center of the room installed a household stove sized $460 \times 410 \times 985 \text{ mm}$ (NS18-17, 18 kW; Laowan Company; Beijing, China). The stove has a thermal efficiency of more than 70%. The air was filtered and then pumped into the sealed room. The filtered PM mass concentration was less than $1 \mu\text{g m}^{-3}$, which was negligible compared with the high PM levels during coal burning. Emissions from

the stove are diluted with the filtered air in the sealed room and drawn into a circular pipe. The cold water entered the stove wall and then hot water flowed out to simulate a water heating process.

Raw coal samples used in this experiment were from Yantang coal mine in Xuanwei City (26.23055°N , 104.09751°E). Details of the coal information can be found in previous studies (Hao *et al.*, 2013; Shao *et al.*, 2015). The coal samples ($\sim 10 \text{ kg}$) were ignited by propane gas with a flow rate of 3 L min^{-1} and the ignition time was $\sim 10 \text{ min}$. Details of the burning-dilution system has been previously described (Li *et al.*, 2016b, c; Zhou *et al.*, 2016).

Sample Collection

A DKL-2 single-stage cascade impactor was used to collect particles on carbon coated Cooper (Cu) grids (300-mesh; Tianld Co.; Beijing, China). The sampler has a 0.5-mm (diameter) jet nozzle. The flow rate was 1 L min^{-1} . The collection efficiency of this sampler is $\sim 100\%$ at $0.5 \mu\text{m}$ if the particle density is 2 g cm^{-3} (Li *et al.*, 2016d). The collected samples were sealed and placed in an air dryer before analysis (Wang *et al.*, 2017).

To better characterize the individual fine particles throughout the burning process, we collected particles in different burning stages, including ignition stage, fierce burning stage and char burning stage. The ignition stage was characterized by low burning temperature and high PM concentration. We collected the samples at ~ 15 minutes after the propane ignition started and the sampling duration for one sample was $\sim 10 \text{ s}$. In fierce burning stages, the burning temperature was high and rapidly increased to its peak of more than 1000°C . We collected the samples after the peak temperature occurred at ~ 1 hour and the sampling duration for one sample was $\sim 25 \text{ s}$. In char burning stages, the burning temperature gradually decreased to $\sim 600^\circ\text{C}$ from its peak. The PM loading was low and the char burning




Fig. 1. Coal burning-dilution chamber measurement system used for generating coal-burning particles.

was dominant. We collected the samples at ~2.3 hours and the sampling duration for one sample was ~60 s. When the above sampling process was completed, we repeated the above sampling process with another 10 kg of coal samples. Therefore, two set of individual particle samples were collected.

TEM Analysis

Hitachi H-8100 TEM (Hitachi, Ltd.; Tokyo, Japan) was used to analyze the individual fine particles and the accelerate voltage was 200 kV. EDX was used to semi-quantitatively acquire the elemental composition with an acquisition time ~30 s and the elements with atomic number higher than 6 can be detected. Copper was not included in our analysis because the TEM grids were made of Cu (Wang *et al.*, 2018).

RESULTS

Particle Types

TEM-EDX can be adequately applied to identify

individual fine particles. Based on the morphology and elemental composition, individual fine particles were classified into two groups: carbonaceous and non-carbonaceous particles. Carbonaceous particles included soot particles and organic particles; non-carbonaceous particles included S-rich particles and mineral particles. Detailed characteristics of individual particles were shown in Table 1.

Soot particles were mainly composed of C and O. They showed distinct chain-like (Fig. 2(a)) or aggregate morphologies (Fig. 2(b)) with hundreds of C-rich spheres, which displayed an onion-like structure with disordered graphic layers under high resolution TEM images (Fig. 2(c)).

Organic particles also mainly consisted of C and O. Some of the organic particles showed inhomogeneous structure with darker and lighter areas under TEM images (Figs. 2(d) and 2(e)); these types of organic particles were near-spherical. The other organic particles showed homogeneous structure and they were spherical or irregular-shaped without any holes (Fig. 2(f)). The organic particles did

Table 1. Types and characteristics of individual particles emitted from the burning of Xuanwei coal.

Particle groups	Particle types	Major elements	Morphologies
Carbonaceous particles	Soot	C and O	Chain-like or aggregate morphologies with hundreds of C-rich spheres
	Organic	C and O	Partly showed inhomogeneous structure with darker and lighter areas; partly showed homogeneous structure with spherical or irregular shapes
Non-carbonaceous particles	Mineral	Si, Fe, Mg, Al, Ca, Ti, K and P	Irregular shaped and tended to have a larger diameter compared with other types of particles
	S-rich	S, O, N and K	Foam-like; beam sensitive and easily decomposed with high energy electron beam irradiation




Fig. 2. TEM images of carbonaceous particles. (a) chain-like soot, (b) aggregated soot, (c) onion-like structured high magnification soot, (d–e) inhomogeneous near-spherical organic particles, and (f) homogeneous irregular-shaped or spherical organic particles.

not show the graphic layers as seen in soot particles under high resolution TEM images.

S-rich particles were foam-like; they were beam-sensitive and easily decomposed with high energy electron beam irradiation (Figs. 3(a) and 3(b)). Some of S-rich particles were mainly composed of S and O (Fig. 3(e)); they were believed to be ammonium sulphates (Fu *et al.*, 2012). The other S-rich particles were mainly composed of S, O and K (Fig. 3(f)), which were believed to be K_2SO_4 (Li *et al.*, 2010).

Mineral particles were irregular-shaped and tended to have a larger diameter compared with other types of particles. They were mainly composed of crustal mineral elements (e.g., Si, Ca, Al, Fe, Na, K, Mg, P). According to their highest elemental composition (Okada *et al.*, 2005), the mineral particles were subdivided into Si-rich (Figs. 3(c) and 3(d)), Ca-rich (Figs. 3(i) and 3(j)), Fe-rich (Fig. 3(k)), Ti-rich (Fig. 3(l)) and other types. Some Si-rich particles only contained Si and O (Figs. 3(c) and 3(g)), and they were identified as SiO_2 ; the other Si-rich particles were mainly composed of Si, Al and O, and/or with minor Ca, Mg, Fe, Na and K (Figs. 3(d) and 3(h)), and were identified as aluminum silicate. Ca-rich particles mainly consisted of Ca, O, C and S, (Figs. 3(m) and 3(n)) and were identified as $CaCO_3$ or $CaSO_4$. Fe-rich particles mainly consisted of Fe and O, and they were mainly Fe_2O_3 or Fe_3O_4 (Figs. 3(k)

and 3(o)). Ti-rich particles mainly consisted of Ti and O, and were identified as TiO_2 (Figs. 3(l) and 3(p)).

Number Fractions in Different Burning Stages

Fig. 4 showed the low magnification TEM images of individual fine particles in different burning stages. Distinct characteristics of fine particles in different burning stages can be seen. The relative number percentage of different types of individual particle in different burning stages was calculated as shown in Fig. 5. We analyzed 873 individual particles in total among 6 TEM samples. In the ignition stage, carbonaceous particles were predominant, with organic particles 66% and soot particles 31% in number, respectively. In the fierce burning stage, carbonaceous particles were also predominant, but the soot particles (71%) were the highest among all analyzed particles, followed by organic particles (28%). In char burning stage, mineral particles were the highest in relative number percentage of all analyzed individual particles, at 73%, followed by organic particles (12%), soot particles (8%) and S-rich particles (7%). Among all analyzed mineral particles, Si-rich particles were predominant, at 49%, followed by Ca-rich (25%), Fe-rich (14%), Ti-rich (7%) and other types (Fig. 6). S-rich particles accounted for a small percentage throughout the whole burning process in this experiment.




Fig. 3. TEM images and elemental compositions of non-carbonaceous particles. (a–b) S-rich particles and (e–f) their elemental compositions, (c) quartz (SiO_2) and (g) its elemental composition, (d) aluminum silicate and (h) its elemental composition, (i) $CaCO_3$ and (m) its elemental composition, (j) $CaSO_4$ and (n) its elemental composition, (k) Fe-rich particles and (o) its elemental composition, (l) TiO_2 particles and (p) its elemental composition.




Fig. 4. Low magnification TEM images of individual particles in different burning stages. (a) organic particles dominated in ignition stage, (b) soot particles dominated in fierce burning stage, and (c) mineral particles dominated in char burning stage.




Fig. 5. Relative number percentage of individual particles in different burning stages. N represents the particle number analyzed.




Fig. 6. Relative number percentage of mineral particles in the char burning stage.

DISCUSSIONS

Xuanwei City has the highest lung cancer morbidity and mortality rates in China, which are attributed to local

household coal burning (Lan *et al.*, 2008; Barone-Adesi *et al.*, 2012; Hosgood *et al.*, 2013; Lui *et al.*, 2017). Detailed knowledge of physical and chemical characteristics of burning-derived individual particles might help to explain the toxicological effects. Considering that the PM emissions in different burning stages show different characteristics (Fig. 5), we hence discuss the PM emission characteristics in different burning stages.

Formation of Carbonaceous Particles in Ignition and Fierce Burning Stages

Household coal burning in China has attracted much attention in recent years because it can emit various kinds of important pollutants (Chen *et al.*, 2015). Household coal burning has low burning efficiency because of the low burning temperature, as a result, household coal burning can often emit a large fraction of carbonaceous particles (Li *et al.*, 2016b).

In the ignition and fierce burning stages, coal chunks undergo pyrolysis and generate a high number of organic volatiles and most of them are burned completely, forming H_2O and CO_2 (Zhou *et al.*, 2016; Li *et al.*, 2017). However, a small number of the volatile matters are not oxidized or

partially oxidized, forming organic aerosols (Wang *et al.*, 2015). Some of the gas phase high-molecular-weight hydrocarbons, such as PAHs can undergo nucleation, condensation or polymerization reactions in very short time periods of ~5 ms to compete with oxidation in the absence of oxygen, forming soot particles (Ma *et al.*, 1996; Fletcher *et al.*, 1997; Richter and Howard, 2000; Mansurov, 2005; Apicella *et al.*, 2017; Xiao *et al.*, 2017). Therefore, the coal pyrolysis product escaped from the burning region and formed either organic or soot particles in the exhaust.

Both the ignition and fierce burning stages were dominated by carbonaceous particles. However, the organic particles were predominant in relative number percentage in the ignition stage while the soot particles were predominant in the fierce burning stage. It should be mentioned that PM emissions from the ignition stage were much higher than in the fierce burning stage and the absolute value of soot particles was much higher in the ignition stage than the fierce burning stage. The higher relative number percentage of soot particles in the fierce burning stage might be related to the burning temperature, concentration of oxygen and the variation of pyrolysis product (Ma *et al.*, 1996; Fletcher *et al.*, 1997; Richter and Howard, 2000; Mansurov, 2005; Apicella *et al.*, 2017; Xiao *et al.*, 2017), and further research is needed in the future.

Only a few of mineral particles were found in the first two burning stages. They may come from the soil dust coated on the surface of the coal when stored and transported or emit from the inner coal chunks along with the emission of pyrolysis product.

Non-carbonaceous Particles Influenced by the Composition of Coal

The volatile materials of coal have been mostly consumed in the first two burning stages, and the char burning became dominant in the following stage (Zhou *et al.*, 2016).

During the coal burning process, the minerals in coal were partially deposited as bottom ash and partially emitted into atmosphere (Lu *et al.*, 2016). With the decreasing of volatile content, mineral particles were dominant in the char burning stage and the type of mineral particles were related with the mineral composition of coal chunks.

The elemental composition of individual particles emitted in char burning stage was complicated. Result from the EDX showed that over 17 elements have been detected in individual non-carbonaceous particles, including O, Si, Fe, Mg, Al, Ca, Ti, S, K, P, Na, Cl, Sr, Ni, V, Mn and Zn, as shown in Fig. 7. O occurred on all analyzed particles. Si, Fe, Mg, Al, Ca, Ti occurred in more than half of all analyzed particles. The result was consistent with the content of major elements in coals from Xuanwei City (Hao *et al.*, 2013; Shao *et al.*, 2015).

Minerals such as quartz (SiO_2), chamosite ($[\text{Fe}^{2+}, \text{Mg}]_2\text{Al}[\text{AlSi}_3\text{O}_{10}]$), calcite (CaCO_3), kaolinite ($\text{Al}_2\text{SiO}_5([\text{OH}]_4)$), and anatase (TiO_2) in Xuanwei coal have been found by Shao *et al.* (2015). Because of the relative low burning temperature, most of the mineral particles were emitted at its original elemental compositions. For example, some quartz (SiO_2), aluminates, calcite (CaCO_3) and anatase (TiO_2) (Figs. 3(c), 3(d), 3(i), and 3(l) from Xuanwei coal can be identified in burning-derived individual particles in this study. However, there were also some new particle formation through chemical reactions. For example, some Fe-rich particles partially resulted from the oxidized product of chamosite or pyrite ($4\text{FeS}_2 + 11\text{O}_2 = 2\text{Fe}_2\text{O}_3 + 8\text{SO}_2$) (Lu *et al.*, 2016). As shown in Fig. 6, the individual mineral particles were predominant in Si-rich (49%), followed by Ca-rich (25%), Fe-rich (14%), Ti-rich (7%) and other types (4%), which is consistent with the main mineral content of Xuanwei coal.

S-rich particles accounted for a small percentage throughout the whole burning process in this experiment.




Fig. 7. Frequency of certain elements present in individual non-carbonaceous particles in char burning stage.

Previous study showed that the emission of S-rich particles were related with the sulfur content of raw coals and the burning of high sulfur content coal can emitted more S-rich particles (Hou *et al.*, 2018). The total sulfur content of Xuanwei coal is extraordinary low, ranging from 0.06 to 0.64% (Shao *et al.*, 2015), resulting in the lower number percentage of S-rich particles.

Implications for Health Risk

Coal combustions can emit large amounts of polycyclic aromatic hydrocarbons (PAHs) which is harmful to human health (Downward *et al.*, 2014). Previous studies have shown that the epidemic diseases might be related to indoor PAHs released from the coal burning in Xuanwei (Mumford *et al.*, 1995; Liu *et al.*, 2017). However, Tian (2005) found there was no obvious relationship between PAHs and lung cancer in a geographical correlation study in Xuanwei. We find a large number of organic particles in the ignition and fierce burning stages in this experiments. Since part of the organic particles belongs to PAHs, further research regarding PAHs is needed in the future.

Mineral particles have been recognized as respiratory hazards (Donaldson and Borm, 2006). The crystalline fine mineral particles were more hazardous when compared with the amorphous mineral ones (Murphy *et al.*, 1998). Most of the mineral particles in household coal burning showed the crystalline structures due to the low burning temperature (Lu *et al.*, 2016).

The content of quartz in Xuanwei coal was almost ten times that found in the other coals (Large *et al.*, 2009). Tian (2005) and Tian *et al.* (2008) have found that most of the burning-derived Si-rich particles were quartz. Si-rich particles accounted for a large number percentage of all analyzed mineral particles, at 49%, much higher than the average (28%) of burning-derived particles from the coals of Zhijin, Datong, Dongsheng, Yinchuan and Jingxi (Hou *et al.*, 2018). In 1996, IARC have classified quartz as a Group 1 substance-carcinogen for humans (International Agency for Research on Cancer; <http://monographs.iarc.fr/ENG/Classification/index.php>). It should be mentioned that higher number percentage of Si-rich particles were also observed by Lu *et al.* (2016, 2017); but the Si-rich particles were not found by using TEM at Fuyuan County, the neighboring area of Xuanwei (Lu *et al.*, 2017).

Fe-rich particles accounted for 14.2%, which was nearly two times of the average (8%) coal burning-derived particles from the coals of Zhijin, Datong, Dongsheng, Yinchuan and Jingxi (Hou *et al.*, 2018). Also, 78% of analyzed mineral particles contained Fe. The reactive oxygen species (ROS) can be formed through chemical reactions: $(\text{Fe}^{2+} + \text{H}_2\text{O}_2 \rightarrow \text{Fe}^{3+} + \text{OH} + \text{OH}^-)$ (Valavanidis *et al.*, 2000; Ambroz *et al.*, 2001; Kim *et al.*, 2001; Breheny, 2014). Once these Fe-containing individual particles were inhaled into the lung, they may do harm to the lung tissues.

CONCLUSIONS

The detailed characteristics of individual particles emitted during the different stages of burning Xuanwei coal were

identified in this study. Four types of individual particles were classified, viz., organic particles, soot particles, S-rich particles, and mineral particles. Number-wise, our results showed that organic particles (66%), soot particles (71%), and mineral particles (73%) were predominant in the ignition stage, fierce burning stage, and char burning stage, respectively, while S-rich particles accounted only for a small percentage.

According to the elemental composition, the mineral portion comprised Si-rich, Ca-rich, Ti-rich, Fe-rich, and other types of particles. In the char burning stage, the particles that were rich in Si, partially identified as quartz or aluminum silicate, were dominant (49%); followed by ones that were Ca-rich (25%), which were identified as CaCO_3 or CaSO_4 ; Fe-rich (14%), which mainly consisted of Fe and O and were recognized as Fe_3O_4 or Fe_2O_3 ; Ti-rich (7%), which were dominated by Ti and O and recognized as TiO_2 ; and other types (4%).

ACKNOWLEDGEMENTS

This work was supported by the Projects of International Cooperation and Exchanges NSFC (Grant No. 41571130031) and National Natural Science Foundation of China (Grant No. 41572090 and No. 41807305).

REFERENCE

- Ambroz, H.B., Bradshaw, T.K., Kemp, T.J., Kornacka, E.M. and Przybytniak, G.K. (2001). Role of iron ions in damage to DNA: Influence of ionising radiation, UV light and H_2O_2 . *J. Photochem. Photobiol. A* 142: 9–18.
- Apicella, B., Senneca, O., Russo, C., Heuer, S., Cortese, L., Cerciello, F., Scherer, V., Schiemann, M. and Ciajolo, A. (2017). Separation and characterization of carbonaceous particulate (soot and char) produced from fast pyrolysis of coal in inert and CO_2 atmospheres. *Fuel* 201: 118–123.
- Barone-Adesi, F., Chapman, R.S., Silverman, D.T., He, X.Z., Hu, W., Vermeulen, R., Ning, B.F., Fraumeni, J.F., Rothman, N. and Lan, Q. (2012). Risk of lung cancer associated with domestic use of coal in Xuanwei, China: retrospective cohort study. *Br. Med. J.* 345: e5414.
- Breheny, D. (2014). Environmental reactive oxygen species (ROS) and cancer, In *Systems biology of free radicals and antioxidants*, Laher, I. (Ed.), Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 2853–2872.
- Cai, S., Li, Q., Wang, S., Chen, J., Ding, D., Zhao, B., Yang, D. and Hao, J. (2018). Pollutant emissions from residential combustion and reduction strategies estimated via a village-based emission inventory in Beijing. *Environ. Pollut.* 238: 230–237.
- Chen, Y.J., Tian, C.G., Feng, Y.L., Zhi, G.R., Li, J. and Zhang, G. (2015). Measurements of emission factors of $\text{PM}_{2.5}$, OC, EC, and BC for household stoves of coal combustion in China. *Atmos. Environ.* 109: 190–196.
- Donaldson, K. and Borm, P. (2006). *Particle toxicology*. CRC Press, Taylor & Francis Group.
- Downward, G.S., Hu, W., Large, D., Veld, H., Xu, J.,

- Reiss, B., Wu, G., Wei, F., Chapman, R.S. and Rothman, N. (2014). Heterogeneity in coal composition and implications for lung cancer risk in Xuanwei and Fuyuan counties, China. *Environ. Int.* 68: 94–104.
- Finkelman, R.B. and Tian, L.W. (2018). The health impacts of coal use in China. *Int. Geol. Rev.* 60: 579–589.
- Fletcher, T.H., Ma, J., Rigby, J.R., Brown, A.L. and Webb, B.W. (1997). Soot in coal combustion systems. *Prog. Energy Combust. Sci.* 23: 283–301.
- Fu, H., Zhang, M., Li, W., Chen, J., Wang, L., Quan, X. and Wang, W. (2012). Morphology, composition and mixing state of individual carbonaceous aerosol in urban Shanghai. *Atmos. Chem. Phys.* 12: 693–707.
- Hao, X., Yi, F., Ren, J., Zhang, R., Liu, D. and Lv, S. (2013). Characteristics of the fine quartz particles from the local coal combustion in Xuanwei, Yunnan Province. *Res. Environ. Sci.* 26: 1043–1049. (in Chinese with English Abstract)
- He, X.Z., Chen, W., Liu, Z.Y. and Chapman, R.S. (1991). An epidemiological study of lung cancer in Xuan Wei County, China: Current progress. Case-control study on lung cancer and cooking fuel. *Environ. Health Perspect.* 94: 9–13.
- Hosgood, H.D., Chapman, R.S., He, X.Z., Hu, W., Tian, L.W., Liu, L.Z., Lai, H., Chen, W., Rothman, N. and Lan, Q. (2013). History of lung disease and risk of lung cancer in a population with high household fuel combustion exposures in rural China. *Lung Cancer* 81: 343–346.
- Hou, C., Shao, L.Y., Zhao, C.M., Wang, J., Liu, J.X., Geng, C.M. (2018). Characterization of coal burning-derived individual particles emitted from an experimental domestic stove. *J. Environ. Sci.* 71: 45–55.
- Hu, W., Downward, G.S., Reiss, B., Xu, J., Bassig, B.A., Hosgood, H.D., Zhang, L.L., Seow, W.J., Wu, G.P., Chapman, R.S., Tian, L.W., Wei, F.S., Vermeulen, R. and Lan, Q. (2014). Personal and indoor PM_{2.5} exposure from burning solid fuels in vented and unvented stoves in a rural region of China with a high incidence of lung cancer. *Environ. Sci. Technol.* 48: 8456–8464.
- Kim, B.Y., Han, M.J. and Chung, A.S. (2001). Effects of reactive oxygen species on proliferation of chinese hamster lung fibroblast (V79) cells. *Free Radical Biol. Med.* 30: 686–698.
- Kim, C., Chapman, R.S., Hu, W., He, X.Z., Hosgood, H.D., Liu, L.Z., Lai, H., Chen, W., Silverman, D.T., Vermeulen, R., Tian, L.W., Bassig, B., Shen, M., Zhang, Y.W., Ma, S.G., Rothman, N. and Lan, Q. (2014). Smoky coal, tobacco smoking, and lung cancer risk in Xuanwei, China. *Lung Cancer* 84: 31–35.
- Lan, Q., He, X.Z., Shen, M., Tian, L.W., Liu, L.Z., Lai, H., Chen, W., Berndt, S.I., Hosgood, H.D., Lee, K.M., Zheng, T.Z., Blair, A. and Chapman, R.S. (2008). Variation in lung cancer risk by smoky coal subtype in Xuanwei, China. *Int. J. Cancer* 123: 2164–2169.
- Large, D.J., Kelly, S., Spiro, B., Tian, L.W., Shao, L.Y., Finkelman, R., Zhang, M., Somerfield, C., Plint, S., Ali, Y. and Zhou, Y.P. (2009). Silica-volatile interaction and the geological cause of the Xuan Wei Lung cancer epidemic. *Environ. Sci. Technol.* 43: 9016–9021.
- Li, Q., Jiang, J.K., Cai, S.Y., Zhou, W., Wang, S.X., Duan, L. and Hao, J.M. (2016a). Gaseous ammonia emissions from coal and biomass combustion in household stoves with different combustion efficiencies. *Environ. Sci. Technol. Lett.* 3: 98–103.
- Li, Q., Jiang, J.K., Zhang, Q., Zhou, W., Cai, S.Y., Duan, L., Ge, S. and Hao, J.M. (2016b). Influences of coal size, volatile matter content, and additive on primary particulate matter emissions from household stove combustion. *Fuel* 182: 780–787.
- Li, Q., Li, X.H., Jiang, J.K., Duan, L., Ge, S., Zhang, Q., Deng, J.G., Wang, S.X. and Hao, J.M. (2016c). Semi-coke briquettes: Towards reducing emissions of primary PM_{2.5}, particulate carbon, and carbon monoxide from household coal combustion in China. *Sci. Rep.* 6: 10.
- Li, Q., Jiang, J.K., Wang, S.X., Rumchev, K., Mead-Hunter, R., Morawska, L. and Hao, J.M. (2017). Impacts of household coal and biomass combustion on indoor and ambient air quality in China: Current status and implication. *Sci. Total Environ* 576: 347–361.
- Li, W.J., Shao, L.Y., and Buseck, P.R. (2010). Haze types in Beijing and the influence of agricultural biomass burning. *Atmos. Chem. Phys.* 10: 8119–8130.
- Li, W.J., Sun, J.X., Xu, L., Shi, Z.B., Riemer, N., Sun, Y.L., Fu, P.Q., Zhang, J.C., Lin, Y.T., Wang, X.F., Shao, L.Y., Chen, J.M., Zhang, X.Y., Wang, Z.F. and Wang, W.X. (2016d). A conceptual framework for mixing structures in individual aerosol particles. *J. Geophys. Res.* 121: 13784–13798.
- Lu, S.L., Hao, X.J., Liu, D.Y., Wang, Q.X., Zhang, W.C., Liu, P.W., Zhang, R.C., Yu, S., Pan, R.Q., Wu, M.H., Yonemochi, S. and Wang, Q.Y. (2016). Mineralogical characterization of ambient fine/ultrafine particles emitted from xuanwei c1 coal combustion. *Atmos. Res.* 169: 17–23.
- Lu, S.L., Tan, Z.Y., Liu, P.W., Zhao, H., Liu, D.Y., Yu, S., Cheng, P., Win, M.S., Hu, J.W., Tian, L.W., Wu, M.H., Yonemochi, S. and Wang, Q.Y. (2017). Single particle aerosol mass spectrometry of coal combustion particles associated with high lung cancer rates in Xuanwei and Fuyuan, China. *Chemosphere* 186: 278–286.
- Lui, K.H., Bandowe, B.A.M., Tian, L.W., Chan, C.S., Cao, J.J., Ning, Z., Lee, S.C. and Ho, K.F. (2017). Cancer risk from polycyclic aromatic compounds in fine particulate matter generated from household coal combustion in Xuanwei, China. *Chemosphere* 169: 660–668.
- Ma, J., Fletcher, T.H. and Webb, B.W. (1996). Conversion of coal tar to soot during coal pyrolysis in a post-flame environment. *Symp. (Int.) Combust.* 26: 3161–3167.
- Mansurov, Z.A. (2005). Soot formation in combustion processes (review). *Combust. Explos. Shock Waves* 41: 727–744.
- Mumford, J.L., Li, X., Hu, F., Lu, X.B. and Chuang, J.C. (1995). Human exposure and dosimetry of polycyclic aromatic hydrocarbons in urine from xuan wei, china with high lung cancer mortality, associated with exposure to unvented coal smoke. *Carcinogenesis* 16: 3031.
- Murphy, S.A., BeruBe, K.A., Pooley, F.D. and Richards,

- R.J. (1998). The response of lung epithelium to well characterised fine particles. *Life Sci.* 62: 1789–1799.
- Okada, K., Qin, Y. and Kai, K. (2005). Elemental composition and mixing properties of atmospheric mineral particles collected in Hohhot, China. *Atmos. Res.* 73: 45–67.
- Pian, W., Cheng, W., Niu, H. and Fan, J. (2016). TEM study of fine particles from coal-fired power plant ambient air. *World J. Eng.* 14: 311–316.
- Richter, H. and Howard, J.B. (2000). Formation of polycyclic aromatic hydrocarbons and their growth to soot - A review of chemical reaction pathways. *Prog. Energy Combust. Sci.* 26: 565–608.
- Shao, L.Y., Wang, J., Hou, H.H., Zhang, M.Q., Wang, H., Spiro, B., Large, D. and Zhou, Y.P. (2015). Geochemistry of the C1 coal of latest Permian during mass extinction in Xuanwei, Yunnan. *Acta Geol. Sin.* 89: 7. (in Chinese with English Abstract)
- Shao, L.Y., Hou, C., Geng, C.M., Liu, J.X., Hu, Y., Wang, J., Jones, T., Zhao, C.M. and Berube, K. (2016). The oxidative potential of PM₁₀ from coal, briquettes and wood charcoal burnt in an experimental domestic stove. *Atmos. Environ.* 127: 372–381.
- Tian, L. (2005). *Coal combustion emissions and lung cancer in Xuanwei, China*. Ph.D. Thesis. University of California, Berkeley USA.
- Tian, L.W., Dai, S.F., Wang, J.F., Huang, Y.C., Ho, S.C., Zhou, Y., Lucas, D. and Koshland, C.P. (2008). Nanoquartz in Late Permian C1 coal and the high incidence of female lung cancer in the Pearl River Origin Area: A retrospective cohort study. *BMC Public Health* 8: 12.
- Tiwari, M., Sahu, S.K. and Pandit, G.G. (2015). Inhalation risk assessment of PAH exposure due to combustion aerosols generated from household fuels. *Aerosol Air Qual. Res.* 15: 582–590.
- Valavanidis, A., Salika, A. and Theodoropoulou, A. (2000). Generation of hydroxyl radicals by urban suspended particulate air matter. the role of iron ions. *Atmos. Environ.* 34: 2379–2386.
- Wang, W.H., Shao, L.Y., Guo, M.L., Hou, C., Xing, J.P. and Wu, F. (2017). Physicochemical properties of individual airborne particles in Beijing during pollution periods. *Aerosol Air Qual. Res.* 17: 3209–3219.
- Wang, W.H., Shao, L.Y., Xing, J.P., Li, J., Chang, L.L. and Li, W.J. (2018). Physicochemical characteristics of individual aerosol particles during the 2015 China victory day parade in Beijing. *Atmosphere* 9: 40.
- Wang, X.F., Cotter, E., Iyer, K.N., Fang, J.X., Williams, B.J. and Biswas, P. (2015). Relationship between pyrolysis products and organic aerosols formed during coal combustion. *Proc. Combust. Inst.* 35: 2347–2354.
- Xiao, Y., Shao, Y., Yu, X. and Zhou, G. (2012). The epidemic status and risk factors of lung cancer in Xuanwei City, Yunnan Province, China. *Front. Med.* 6: 388–394.
- Xiao, Z.K., Tang, Y., Zhuo, J.K. and Yao, Q. (2017). Effect of the interaction between sodium and soot on fine particle formation in the early stage of coal combustion. *Fuel* 206: 546–554.
- Zhang, H.F., Zhu, T., Wang, S.X., Hao, J.M., Mestl, H.E.S., Alnes, L.W.H., Aunan, K., Dong, Z.Q., Ma, L.Y., Hu, Y., Zhang, M., Mellouki, A.W., Chai, F. and Wang, S.L. (2014). Indoor emissions of carbonaceous aerosol and other air pollutants from household fuel burning in Southwest China. *Aerosol Air Qual. Res.* 14: 1779–1288.
- Zhang, J.J. and Smith, K.R. (2007). Household air pollution from coal and biomass fuels in China: Measurements, health impacts, and interventions. *Environ. Health Perspect.* 115: 848–855.
- Zhang, Y.X., Schauer, J.J., Zhang, Y.H., Zeng, L.M., Wei, Y.J., Liu, Y. and Shao, M. (2008). Characteristics of particulate carbon emissions from real-world Chinese coal combustion. *Environ. Sci. Technol.* 42: 5068–5073.
- Zhou, W., Jiang, J.K., Duan, L. and Hao, J.M. (2016). Evolution of submicrometer organic aerosols during a complete residential coal combustion process. *Environ. Sci. Technol.* 50: 7861–7869.
- Zhu, C.S., Cao, J.J., Shen, Z.X., Liu, S.X., Zhang, T., Zhao, Z.Z., Xu, H.M. and Zhang, E.K. (2012). Indoor and outdoor chemical components of PM_{2.5} in the rural areas of northwestern China. *Aerosol Air Qual. Res.* 12: 1157–116.

Received for review, May 24, 2018

Revised, August 8, 2018

Accepted, August 27, 2018